

This is the first author's version of a work that was accepted for publication in the Journal of Algebraic Combinatorics. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *J. Algebraic Combin.* **44** (2016) 393–432.

<http://dx.doi.org/10.1007/s10801-016-0674-x>

The final publication is available at www.springerlink.com.

Generalised column removal for graded homomorphisms between Specht modules

Matthew Fayers

Queen Mary University of London,
Mile End Road,
London E1 4NS,
U.K.

m.fayers@qmul.ac.uk

Liron Speyer

Osaka University,
Suita,
Osaka 565-0871,
Japan

l.speyer@ist.osaka-u.ac.jp

2010 Mathematics subject classification: 20C30, 20C08, 05E10

Abstract

Let n be a positive integer, and let \mathcal{H}_n denote the affine KLR algebra in type A. Kleshchev, Mathas and Ram have given a homogeneous presentation for graded column Specht modules S_λ for \mathcal{H}_n . Given two multipartitions λ and μ , we define the notion of a *dominated* homomorphism $S_\lambda \rightarrow S_\mu$, and use the KMR presentation to prove a generalised column-removal theorem for graded dominated homomorphisms between Specht modules. In the process, we prove some useful properties of \mathcal{H}_n -homomorphisms between Specht modules which lead to an immediate corollary that, subject to a few demonstrably necessary conditions, every homomorphism $S_\lambda \rightarrow S_\mu$ is dominated, and in particular $\text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu) = 0$ unless λ dominates μ .

Brundan and Kleshchev show that certain cyclotomic quotients of \mathcal{H}_n are isomorphic to (degenerate) cyclotomic Hecke algebras of type A. Via this isomorphism, our results can be seen as a broad generalisation of the column-removal results of Fayers and Lyle and of Lyle and Mathas; generalising both into arbitrary level and into the graded setting.

1 Introduction

The KLR algebras, or quiver Hecke algebras, were constructed independently by Khovanov and Lauda [KL] and by Rouquier [R2], and have since received an abundance of interest. This is, in some part, due to the powerful result of Brundan and Kleshchev in [BK] that every (degenerate) Ariki–Koike algebra is isomorphic to a so-called cyclotomic quotient of a KLR algebra. The KLR algebras and their cyclotomic quotients are graded, and this allows us to study the graded representation theory of (degenerate) Ariki–Koike algebras, and in particular the graded representation theory of the symmetric groups. This motivates the study of KLR algebras, and in particular the study of their graded Specht modules. These were defined by Brundan, Kleshchev and Wang [BKW], and developed further by Kleshchev, Mathas and Ram [KMR], who gave a homogeneous presentation for each Specht module.

In trying to understand the (graded) structure of the Specht modules, the (graded) homomorphism spaces $\text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$ are of particular interest. In the ungraded setting, these

homomorphism spaces have received a great deal of attention in recent years. We concentrate in particular here on the row- and column-removal theorems for homomorphisms, proved by the first author and Lyle [FL] for the symmetric group and generalised to Hecke algebras of type A by Lyle and Mathas. In this paper, we provide graded versions of these theorems, while at the same time generalising them to higher levels so that they apply to all (degenerate) Ariki–Koike algebras.

In fact, our results apply not to all homomorphisms between two given Specht modules but only to those of a certain type, which we call *dominated* homomorphisms. However, in many cases (for example, for the symmetric group in odd characteristic) every homomorphism between two Specht modules is dominated, so our results apply generally; in particular, via the Brundan–Kleshchev isomorphism mentioned above, we recover the original row- and column-removal theorems of Lyle and Mathas.

We now summarise the structure of the paper. In Section 2, we introduce the combinatorics necessary for our purposes, as well as the set-up of the KLR algebras and their Specht modules. We proceed in Section 3 by introducing *dominated tableaux* and the corresponding dominated homomorphisms. Section 4 gives our main results pertaining to generalised column removal for homomorphisms. Finally, in Section 5 we provide an index of notation for the reader’s convenient reference.

Acknowledgements. The second author would like to thank Queen Mary University of London, without whose funding this work would not have been possible. The second author must also thank Professor Andrew Mathas, at the University of Sydney, with whom this work began. The visit to the University of Sydney was funded by the Eileen Colyer Prize and the Australian Research Council grant DP110100050 “Graded representations of Hecke algebras”. We are grateful to several anonymous referees for their extensive and helpful comments.

2 Background

In this section we recall some background and set up some notation. This varies from [KMR] in only a few details.

2.1 The symmetric group

Let \mathfrak{S}_n denote the symmetric group of degree n . Let s_1, \dots, s_{n-1} denote the standard Coxeter generators of \mathfrak{S}_n , i.e. s_i is the transposition $(i, i+1)$. Given $w \in \mathfrak{S}_n$, a *reduced expression* for w is an expression $w = s_{i_1} \dots s_{i_l}$ with l as small as possible; we call $l = l(w)$ the *length* of w .

We will need to use two natural partial orders on \mathfrak{S}_n . If $w, x \in \mathfrak{S}_n$, then we say that x is smaller than w in the *left order* (and write $x \leq_L w$) if $l(w) = l(wx^{-1}) + l(x)$; this is equivalent to the statement that there is a reduced expression for w which has a reduced expression for x as a suffix.

More important will be the *Bruhat order* on \mathfrak{S}_n : if $w, x \in \mathfrak{S}_n$, then we say that x is smaller than w in the Bruhat order (and write $x \preceq w$) if there is a reduced expression for w which has a (possibly non-reduced) expression for x as a subsequence. In fact [H, Theorem 5.10], if $x \preceq w$, then every reduced expression has a reduced expression for x as a subsequence.

The following proposition gives an alternative characterisation of the Bruhat order.

Proposition 2.1 [H, §5.9]. Suppose $w, x \in \mathfrak{S}_n$. Then $w \preceq x$ if and only if there are $w = w_0, w_1, \dots, w_r = x$ such that for each $1 \leq i \leq r$ we have $w_i = (u_i, v_i)w_{i-1}$, where $1 \leq u_i < v_i \leq n$ and $w_{i-1}^{-1}(u_i) < w_{i-1}^{-1}(v_i)$.

Later we shall need the following lemma; in fact, this is a special case of Deodhar's 'property Z' [D, Theorem 1.1].

Lemma 2.2. Suppose $w, x \in \mathfrak{S}_n$ with $x < w$. If $l(s_i w) < l(w)$ while $l(s_i x) > l(x)$, then $s_i x \preceq w$.


Proof. Since $l(s_i w) < l(w)$, w has a reduced expression s beginning with s_i . We can find a reduced expression for x as a subexpression of s , and this subexpression cannot include the first term s_i , since $l(s_i x) > l(x)$. So we can add the initial s_i to the subexpression to get a reduced expression for $s_i x$ as a subexpression of s . \square

We end this subsection by defining some very natural and useful homomorphisms. Suppose $1 \leq m \leq n$ and $0 \leq k \leq n - m$, and define the homomorphism $\text{shift}_k : \mathfrak{S}_m \rightarrow \mathfrak{S}_n$ by $s_i \mapsto s_{i+k}$ for every i . Note that if $k = 0$, this is the natural embedding.

2.2 Lie-theoretic notation

Throughout this paper e is a fixed element of the set $\{2, 3, 4, \dots\} \cup \{\infty\}$. If $e = \infty$ then we set $I := \mathbb{Z}$, while if $e < \infty$ then we set $I := \mathbb{Z}/e\mathbb{Z}$; we may identify I with the set $\{0, \dots, e-1\}$ when convenient. The Cartan matrix $(a_{ij})_{i,j \in I}$ is defined by $a_{ij} = 2\delta_{ij} - \delta_{i(j+1)} - \delta_{i(j-1)}$.

Let Γ be the quiver with vertex set I and an arrow from i to $i-1$ for each i . (Note that this convention is the same as that in [KMR], and opposite to that in [BK, BKW].) The quiver Γ is pictured below for some values of e .



In the relations we give below, we use arrows with reference to Γ ; thus we may write $i \rightarrow j$ to mean that $e \neq 2$ and $j = i-1$, or $i \rightleftarrows j$ to mean that $e = 2$ and $j = i-1$.

We adopt standard notation from Kac's book [K] for the Kac-Moody algebra associated to the Cartan matrix $(a_{ij})_{i,j \in I}$; in particular, we have fundamental dominant weights Λ_i and simple roots α_i for $i \in I$, and an invariant symmetric bilinear form (\mid) satisfying $(\Lambda_i \mid \alpha_j) = \delta_{ij}$ and $(\alpha_i \mid \alpha_j) = a_{ij}$ for $i, j \in I$. We let $Q^+ := \bigoplus_{i \in I} \mathbb{Z}_{\geq 0} \alpha_i$ be the *positive root lattice*. For $\alpha = \sum_{i \in I} c_i \alpha_i \in Q^+$, we define the *height* of α to be $\sum_{i \in I} c_i$. Given $\alpha, \beta \in Q^+$ with $\alpha = \sum_{i \in I} c_i \alpha_i$ and $\beta = \sum_{i \in I} d_i \alpha_i$, we write $\alpha \geq \beta$ if $c_i \geq d_i$ for each i .

Let I^l denote the set of all l -tuples of elements of I . We call an element of I^l an e -multicharge of level l . The symmetric group \mathfrak{S}_l acts on I^l on the left by place permutations. Given an e -multicharge $\kappa = (\kappa_1, \dots, \kappa_l)$, we define a corresponding dominant weight $\Lambda_\kappa := \Lambda_{\kappa_1} + \dots + \Lambda_{\kappa_l}$. For $\alpha \in Q^+$, we then define the *defect* of α (with respect to κ) to be

$$\text{def}(\alpha) = (\Lambda_\kappa \mid \alpha) - \frac{1}{2}(\alpha \mid \alpha).$$

2.3 Multicompositions and multipartitions

A *composition* is a sequence $\lambda = (\lambda_1, \lambda_2, \dots)$ of non-negative integers such that $\lambda_i = 0$ for sufficiently large i . We write $|\lambda|$ for the sum $\lambda_1 + \lambda_2 + \dots$. When writing compositions, we may omit trailing zeroes and group equal parts together with a superscript. We write \emptyset for the composition $(0, 0, \dots)$. A *partition* is a composition λ for which $\lambda_1 \geq \lambda_2 \geq \dots$.

Now suppose $l \in \mathbb{N}$. An l -*multicomposition* is an l -tuple $\lambda = (\lambda^{(1)}, \dots, \lambda^{(l)})$ of compositions, which we refer to as the *components* of λ . We write $|\lambda| = |\lambda^{(1)}| + \dots + |\lambda^{(l)}|$, and say that λ is an l -multicomposition of $|\lambda|$. If the components of λ are all partitions, then we say that λ is an l -*multipartition*. We write \mathcal{P}_n^l for the set of l -multipartitions of n . We abuse notation by using \emptyset also for the multipartition $(\emptyset, \dots, \emptyset)$.

If λ and μ are l -multicompositions of n , then we say that λ *dominates* μ , and write $\lambda \triangleright \mu$, if

$$|\lambda^{(1)}| + \dots + |\lambda^{(m-1)}| + \lambda_1^{(m)} + \dots + \lambda_r^{(m)} \geq |\mu^{(1)}| + \dots + |\mu^{(m-1)}| + \mu_1^{(m)} + \dots + \mu_r^{(m)}$$


for all $1 \leq m \leq l$ and $r \geq 0$.

If λ is an l -multicomposition, the *Young diagram* $[\lambda]$ is defined to be the set

$$\left\{ (r, c, m) \in \mathbb{N} \times \mathbb{N} \times \{1, \dots, l\} \mid c \leq \lambda_r^{(m)} \right\}.$$

We refer to the elements of $[\lambda]$ as the *nodes* of λ . We may also refer to (r, c, m) as the (r, c) -node of $\lambda^{(m)}$. If $\lambda \in \mathcal{P}_n^l$, a node of λ is *removable* if it can be removed from $[\lambda]$ to leave the Young diagram of a smaller l -multipartition, while a node not in $[\lambda]$ is *addable* if it can be added to $[\lambda]$ to leave the Young diagram of an l -multipartition.

We adopt an unusual (but in our view, extremely helpful) convention for drawing Young diagrams. We draw the nodes of each component as boxes in the plane, using the English convention, where the first coordinate increases down the page and the second coordinate increases from left to right. Then we arrange the diagrams for the components *in a diagonal line from top right to bottom left*. For example, if $\lambda = ((2^2), (2, 1^2), (3, 2)) \in \mathcal{P}_{13}^3$, then $[\lambda]$ is drawn as follows.



We shall use directions such as left and right with reference to this convention; for example, we shall say that a node (r, c, m) lies to the left of (r', c', m') if either $m > m'$ or $(m = m' \text{ and } c < c')$. Similarly, we say that (r, c, m) is above, or higher than, (r', c', m') if either $m < m'$ or $(m = m' \text{ and } r < r')$.

If λ is a partition, the *conjugate partition* λ' is defined by

$$\lambda'_i = \left| \left\{ j \geq 1 \mid \lambda_j \geq i \right\} \right|.$$

If λ is an l -multipartition, then the conjugate multipartition λ' is given by

$$\lambda' = (\lambda^{(l)'}, \dots, \lambda^{(1)'}).$$

Observe that with our convention, the Young diagram $[\lambda']$ may be obtained from $[\lambda]$ by reflecting in a diagonal line running from top left to bottom right.

2.4 Tableaux

If $\lambda \in \mathcal{P}_n^l$, a λ -tableau is a bijection $T : [\lambda] \rightarrow \{1, \dots, n\}$. We depict a λ -tableau T by drawing the Young diagram $[\lambda]$ and filling each box with its image under T . T is *row-strict* if its entries increase from left to right along each row of the diagram, and *column-strict* if its entries increase down each column. T is *standard* if it is both row- and column-strict. We write $\text{Std}(\lambda)$ for the set of standard λ -tableaux.

If T is a λ -tableau, then we define a λ' -tableau T' by

$$T'(r, c, m) = T(c, r, l + 1 - m)$$

for all $(r, c, m) \in [\lambda']$.

We import and modify some notation from [BKW] and [KMR]: given a tableau T and $1 \leq i, j \leq n$, we write $i \rightarrow_T j$ to mean that i and j lie in the same row of the same component, with j to the right of i . We write $i \nearrow_T j$ to mean that i and j lie in the same component of T , with j strictly higher and strictly to the right, and we write $i \nearrow_T j$ to mean that either $i \nearrow_T j$ or j lies in an earlier component than i . The notations $i \downarrow_T j$, $i \swarrow_T j$ and $i \searrow_T j$ are defined similarly.

There are two standard λ -tableaux of particular importance. The tableau T_λ is the standard tableau obtained by writing $1, \dots, n$ in order down successive columns from left to right, while T^λ is the tableau obtained by writing $1, \dots, n$ in order along successive rows from top to bottom. Note that we then have $T^\lambda = (T_\lambda)'$.

Example. With $\lambda = ((2^2), (2, 1^2), (3, 2))$ we have

$$T_\lambda = \begin{array}{|c|c|c|} \hline 10 & 12 \\ \hline 11 & 13 \\ \hline \end{array}, \quad T^\lambda = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline \end{array}$$

$$\begin{array}{|c|c|} \hline 6 & 9 \\ \hline 7 & \\ \hline 8 & \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 5 & 6 \\ \hline 7 & \\ \hline 8 & \\ \hline \end{array}$$

$$\begin{array}{|c|c|c|} \hline 1 & 3 & 5 \\ \hline 2 & 4 & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 9 & 10 & 11 \\ \hline 12 & 13 & \\ \hline \end{array}$$

The symmetric group \mathfrak{S}_n acts naturally on the left on the set of λ -tableaux. Given a λ -tableau T , we define the permutations w_T and w^T in \mathfrak{S}_n by

$$w_T T_\lambda = T = w^T T^\lambda.$$

Later we shall need the following lemma; recall that \leq_L denotes the left order on \mathfrak{S}_n .

Lemma 2.3. Suppose $\lambda \in \mathcal{P}_n^l$ and S, T are λ -tableaux with $w_S \leq_L w_T$. If T is standard, then S is standard.

Proof. Using induction on $l(w_T) - l(w_S)$, we may assume $l(w_T) = l(w_S) + 1$, which means in particular that $T = s_i S$ for some i . Since T is standard, the only way S could fail to be standard is if $i + 1$ occupies the node immediately below or immediately to the right of i in T . But either possibility means that i occurs before $i + 1$ in the ‘column reading word’ of T , i.e. the word obtained by reading the entries of T down successive columns from left to right. In other words, $w_T^{-1}(i) < w_T^{-1}(i + 1)$, but this means that $l(w_S) > l(w_T)$, a contradiction. \square

Now we introduce a dominance order on tableaux. If S, T are λ -tableaux, then we write $S \trianglerighteq T$ if and only if $w_S \succcurlyeq w_T$ (recall that \succcurlyeq denotes the Bruhat order on \mathfrak{S}_n). There should be no ambiguity in using the symbol \trianglerighteq for both the dominance order on multipartitions and the dominance order on tableaux.

There is an alternative description of the dominance order on tableaux which will be very useful. If T is a λ -tableau and $0 \leq m \leq n$, we define $T_{\downarrow m}$ to be the set of nodes of $[\lambda]$ whose entries are less than or equal to m . If T is row-strict, then $T_{\downarrow m}$ is the Young diagram of an l -multicomposition of m , which we call $\text{Shape}(T_{\downarrow m})$. If T is standard, then $\text{Shape}(T_{\downarrow m})$ is an l -multipartition of m .

Now we have the following proposition. This is proved in the case $l = 1$ in [M, Theorem 3.8] (where it is attributed to Ehresmann and James); in fact, the proof in [M] carries over to the case of arbitrary l without any modification.

Proposition 2.4. *Suppose $\lambda \in \mathcal{P}_n^l$ and S, T are row-strict λ -tableaux. Then $S \trianglelefteq T$ if and only if $\text{Shape}(S_{\downarrow m}) \trianglelefteq \text{Shape}(T_{\downarrow m})$ for $m = 1, \dots, n$.*

In this paper, we shall briefly consider a natural analogue of this notion for column-strict tableaux. Suppose $\lambda \in \mathcal{P}_n^l$ and T is a column-strict λ -tableau; define the diagram $T_{\downarrow m}$ as above, and define $T'_{\downarrow m}$ to be the ‘conjugate diagram’ to $T_{\downarrow m}$, that is

$$T'_{\downarrow m} = \{(c, r, l+1-k) \mid (r, c, k) \in T_{\downarrow m}\}.$$

Then $T'_{\downarrow m}$ is the Young diagram of an l -multicomposition of m , which we denote $\text{Shape}(T_{\downarrow m})'$. Now we have following statement, which can be deduced from Proposition 2.4 by conjugating tableaux.

Proposition 2.5. *Suppose $\lambda \in \mathcal{P}_n^l$ and S, T are column-strict λ -tableaux. Then $S \trianglelefteq T$ if and only if $\text{Shape}(S_{\downarrow m})' \trianglerighteq \text{Shape}(T_{\downarrow m})'$ for $m = 1, \dots, n$.*

2.5 Residues and degrees

In this section we connect the Lie-theoretic set-up above with multipartitions and tableaux. We fix an e -multicharge $\kappa = (\kappa_1, \dots, \kappa_l)$. We define the *residue* $\text{res } A = \text{res}^\kappa A$ of a node $A = (r, c, m) \in \mathbb{N} \times \mathbb{N} \times \{1, \dots, l\}$ by

$$\text{res } A = \kappa_m + (c - r) \pmod{e}.$$

We say that A is an i -node if it has residue i . Given $\lambda \in \mathcal{P}_n^l$, we define the *content* of λ to be the element

$$\text{cont}(\lambda) = \sum_{A \in [\lambda]} \alpha_{\text{res } A} \in Q^+.$$

We then define the *defect* $\text{def}(\lambda)$ of λ to be $\text{def}(\text{cont}(\lambda))$.

If T is a λ -tableau, we define its *residue sequence* to be the sequence $i(T) = (i_1, \dots, i_n)$, where i_r is the residue of the node $T^{-1}(r)$, for each r . The residue sequences of the tableaux T_λ and T^λ will be of particular importance, and we set $i_\lambda := i(T_\lambda)$ and $i^\lambda := i(T^\lambda)$.

Example. Take $\lambda = ((2^2), (2, 1^2), (3, 2))$ as in the last example, and suppose $e = 4$ and $\kappa = (1, 2, 0)$. Then the residues of the nodes of λ are given by the following diagram.

So we have

$$i_\lambda = (0, 3, 1, 0, 2, 2, 1, 0, 3, 1, 0, 2, 1), \quad i^\lambda = (1, 2, 0, 1, 2, 3, 1, 0, 0, 1, 2, 3, 0).$$

Now we recall from [BKW, §3.5] the degree and codegree of a standard tableau. Suppose $\lambda \in \mathcal{P}_n^l$ and A is an i -node of λ . Set

$d_A(\lambda) := |\{\text{addable } i\text{-nodes of } \lambda \text{ strictly below } A\}| - |\{\text{removable } i\text{-nodes of } \lambda \text{ strictly below } A\}|$,
and

$$d^A(\lambda) := |\{\text{addable } i\text{-nodes of } \lambda \text{ strictly above } A\}| - |\{\text{removable } i\text{-nodes of } \lambda \text{ strictly above } A\}|.$$

For $T \in \text{Std}(\lambda)$ we define the *degree* of T recursively, setting $\deg(T) := 0$ when T is the unique \emptyset -tableau. If $T \in \text{Std}(\lambda)$ with $|\lambda| > 0$, let $A = T^{-1}(n)$, let $T_{<n}$ be the tableau obtained by removing this node and set

$$\deg(T) := d_A(\lambda) + \deg(T_{<n}).$$

Similarly, define the *codegree* of T by setting $\text{codeg}(T) := 0$ if T is the unique \emptyset -tableau, and

$$\text{codeg}(\mathbf{T}) := d^A(\lambda) + \text{codeg}(T_{<n})$$

for $T \in \text{Std}(\lambda)$ with $|\lambda| > 0$. We note that the definitions of degree and codegree depend on the e -multicharge κ , and therefore we write \deg^κ and codeg^κ when we wish to emphasise κ .

Example. Suppose $e = 3$, $\kappa = (1, 1)$ and T is the $((2), (2, 1))$ -tableau

1	5
2	

3	4
---	---

which has residue sequence $i(T) = (1, 0, 1, 2, 2)$. Letting $A = T^{-1}(5) = (1, 2, 2)$, we find that $d_A(\lambda) = 1$ and $d^A(\lambda) = -1$. Recursively one finds that for the tableau

$$T_{<5} = \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 3 & 4 \\ \hline \end{array}$$

we have $\deg(T_{<5}) = 2$ and $\text{codeg}(T_{<5}) = 1$, so that $\deg(T) = 3$ and $\text{codeg}(T) = 0$.

The degree, codegree of a standard λ -tableau are related to the defect of λ by the following result.

Lemma 2.6 [BKW, Lemma 3.12]. *Suppose $\lambda \in \mathcal{P}_n^l$ and $T \in \text{Std}(\lambda)$. Then*

$$\deg(T) + \operatorname{codeg}(T) = \operatorname{def}(\lambda).$$

2.6 KLR algebras

We now give the definition of the algebras which will be our main object of study. Throughout this paper we fix a field \mathbb{F} .

Suppose $\alpha \in Q^+$ has height n , and set

$$I^\alpha = \{i \in I^n \mid \alpha_{i_1} + \cdots + \alpha_{i_n} = \alpha\}.$$

Now define \mathcal{H}_α to be the unital associative \mathbb{F} -algebra with generating set

$$\{e(i) \mid i \in I^\alpha\} \cup \{y_1, \dots, y_n\} \cup \{\psi_1, \dots, \psi_{n-1}\}$$

and relations

$$\begin{aligned} e(i)e(j) &= \delta_{i,j}e(i); \\ \sum_{i \in I^\alpha} e(i) &= 1; \\ y_re(i) &= e(i)y_r; \\ \psi_re(i) &= e(s_r i)\psi_r; \\ y_ry_s &= y_sy_r; \\ \psi_r y_s &= y_s \psi_r && \text{if } s \neq r, r+1; \\ \psi_r \psi_s &= \psi_s \psi_r && \text{if } |r-s| > 1; \\ y_r \psi_r e(i) &= (\psi_r y_{r+1} - \delta_{i_r, i_{r+1}})e(i); \\ y_{r+1} \psi_r e(i) &= (\psi_r y_r + \delta_{i_r, i_{r+1}})e(i); \\ \psi_r^2 e(i) &= \begin{cases} 0 & \text{if } i_r = i_{r+1}, \\ e(i) & \text{if } i_{r+1} \neq i_r, i_r \pm 1, \\ (y_{r+1} - y_r)e(i) & \text{if } i_r \rightarrow i_{r+1}, \\ (y_r - y_{r+1})e(i) & \text{if } i_r \leftarrow i_{r+1}, \\ (y_{r+1} - y_r)(y_r - y_{r+1})e(i) & \text{if } i_r \rightleftarrows i_{r+1}; \end{cases} \\ \psi_r \psi_{r+1} \psi_r e(i) &= \begin{cases} (\psi_{r+1} \psi_r \psi_{r+1} + 1)e(i) & \text{if } i_{r+2} = i_r \rightarrow i_{r+1}, \\ (\psi_{r+1} \psi_r \psi_{r+1} - 1)e(i) & \text{if } i_{r+2} = i_r \leftarrow i_{r+1}, \\ (\psi_{r+1} \psi_r \psi_{r+1} + y_r - 2y_{r+1} + y_{r+2})e(i) & \text{if } i_{r+2} = i_r \rightleftarrows i_{r+1}, \\ (\psi_{r+1} \psi_r \psi_{r+1})e(i) & \text{otherwise;} \end{cases} \end{aligned}$$

for all admissible r, s, i, j .

The *affine Khovanov–Lauda–Rouquier algebra* or *quiver Hecke algebra* \mathcal{H}_n is defined to be the direct sum $\bigoplus_\alpha \mathcal{H}_\alpha$, where the sum is taken over all $\alpha \in Q^+$ of height n .

Remarks.

1. We use the same notation for the generators ψ_r and y_s for different α ; when using these generators, we shall always make it clear which algebra \mathcal{H}_α these generators are taken from.
2. When $e < \infty$, we can modify the above presentation of \mathcal{H}_α to give a presentation for \mathcal{H}_n : we take generating set $\{e(i) \mid i \in I^n\} \cup \{y_1, \dots, y_n\} \cup \{\psi_1, \dots, \psi_{n-1}\}$, and replace the relation

$\sum_{i \in I^\alpha} e(i) = 1$ with $\sum_{i \in I^n} e(i) = 1$. The generator ψ_r in this presentation is just the sum of the corresponding generators ψ_r of the individual algebras \mathcal{H}_α in the direct sum $\bigoplus_\alpha \mathcal{H}_\alpha$, and similarly for y_s . When $e = \infty$ we cannot do this, since the set I^n is infinite (in fact, \mathcal{H}_n is non-unital in this case).

The following result can easily be checked from the definition of \mathcal{H}_α .

Lemma 2.7 [BK, Corollary 1]. *There is a \mathbb{Z} -grading on the algebra \mathcal{H}_α such that for all admissible r and i ,*

$$\deg(e(i)) = 0, \quad \deg(y_r) = 2, \quad \deg(\psi_r e(i)) = -a_{i_r i_{r+1}}.$$

Shift maps

Recall from §2.1 that $\text{shift}_k : \mathfrak{S}_m \rightarrow \mathfrak{S}_n$ denotes the homomorphism defined by $s_i \mapsto s_{i+k}$. We now define the corresponding maps for the algebras \mathcal{H}_α .

Definition. Suppose $1 \leq m \leq n$ and $0 \leq k \leq n - m$, $\alpha, \beta \in Q^+$ with α of height n and β of height m . Given $i \in I^\beta$, define $J_i := \{j \in I^\alpha \mid j_{s+k} = i_s \text{ for } 1 \leq s \leq m\}$, and let $e(i)^{+k} = \sum_{j \in J_i} e(j)$. Now define the homomorphism $\text{shift}_k : \mathcal{H}_\beta \rightarrow \mathcal{H}_\alpha$ by

$$e(i) \mapsto e(i)^{+k}, \quad \psi_r e(i) \mapsto \psi_{r+k} e(i)^{+k}, \quad y_r e(i) \mapsto y_{r+k} e(i)^{+k}.$$

It is easy to check from the definition of \mathcal{H}_α that shift_k is a degree-preserving (non-unital) homomorphism of algebras. Moreover, the PBW-type basis theorem for \mathcal{H}_α in [KL, Theorem 2.5] and [R2, Theorem 3.7] shows that if $\beta \leq \alpha$ then shift_k is injective (obviously shift_k is the zero map if $\beta \not\leq \alpha$).

Cyclotomic algebras and the Brundan–Kleshchev isomorphism theorem

Given $\alpha \in Q^+$ and an e -multicharge $\kappa = (\kappa_1, \dots, \kappa_l) \in I^l$, we define $\mathcal{H}_\alpha^\kappa$ to be the quotient of \mathcal{H}_α by the cyclotomic relations

$$y_1^{(\Lambda_\kappa \mid \alpha_{i_1})} e(i) = 0 \quad \text{for } i \in I^\alpha.$$

The cyclotomic KLR algebra \mathcal{H}_n^κ is then defined to be the sum $\bigoplus_\alpha \mathcal{H}_\alpha^\kappa$. Here we sum over all $\alpha \in Q^+$ of height n , though in fact only finitely many of the summands will be non-zero, so (even when $e = \infty$) \mathcal{H}_n^κ is a unital algebra.

Note that the embedding shift_0 passes naturally into the cyclotomic quotients.

A stunning result of Brundan and Kleshchev [BK, Main Theorem] is that if $e = \infty$ or if \mathbb{F} contains a primitive e th root of unity, then \mathcal{H}_n^κ is isomorphic to an Ariki–Koike algebra of level l , defined at an e th root of unity. Similarly, if $e = \text{char}(\mathbb{F})$, then \mathcal{H}_n^κ is isomorphic to a degenerate Ariki–Koike algebra; in particular, when $l = 1$, \mathcal{H}_n^κ is isomorphic to the group algebra $\mathbb{F}\mathfrak{S}_n$. As a consequence, these Hecke algebras are non-trivially \mathbb{Z} -graded. This theorem motivates our choice of notation \mathcal{H}_n for the KLR algebra.

2.7 Specht modules

We now recall the universal graded row and column Specht modules introduced by Kleshchev, Mathas and Ram; we refer the reader to [KMR, §§5,7] for further details.

Fix an e -multicharge κ . Suppose $\lambda \in \mathcal{P}_n^l$, and let $\alpha = \text{cont}(\lambda)$. Say that a node $A = (r, c, m) \in [\lambda]$ is a *column Garnir node* if $(r, c+1, m) \in [\lambda]$. The *column Garnir belt* \mathbf{B}_A is defined to be the set of nodes

$$\mathbf{B}_A = \{(s, c, m) \in [\lambda] \mid s \geq r\} \cup \{(s, c+1, m) \in [\lambda] \mid s \leq r\}.$$

This belt is used to define a *column Garnir element* $\mathbf{g}_A \in \mathcal{H}_\alpha$. The full definition of \mathbf{g}_A is quite complicated, and can be found in [KMR, Definition 7.10]. Here we just give \mathbf{g}_A explicitly in a special case which we will use in the proof of Proposition 3.10, and record some useful properties of \mathbf{g}_A which apply in general.

For our special case, we suppose that A is a Garnir node of λ of the form $(1, c, m)$. If a is the entry in node A of T_λ and b is the entry in node $(1, c+1, m)$, then $\mathbf{g}_A = \psi_a \psi_{a+1} \dots \psi_{b-1}$.

Now suppose $A = (r, c, m)$ is an arbitrary Garnir node of λ . Then in T_λ the nodes of \mathbf{B}_A are occupied by the integers $a, a+1, \dots, b$ for some $a < b$. The following facts can be distilled from [KMR, §7]:

- \mathbf{g}_A is a linear combination of products of the form $\psi_{i_1} \dots \psi_{i_d}$ where $a \leq i_1, \dots, i_d < b$;
- \mathbf{g}_A depends only on e, r, a and the length of the column containing A .

(In fact, as defined in [KMR], the column Garnir element \mathbf{g}_A also involves an idempotent $e(i)$ which depends on λ and makes \mathbf{g}_A homogeneous, but this term can be omitted without affecting the Garnir relation given below.)

Example. For example, let $\lambda = ((3, 3, 2, 2, 1), (2, 1))$ and let $A = (3, 1, 1)$. Then A is a column Garnir node, and T_λ (with the Garnir belt \mathbf{B}_A shaded) is as follows.

$$T_\lambda = \begin{array}{c} \begin{array}{|c|c|c|} \hline 4 & 9 & 13 \\ \hline 5 & 10 & 14 \\ \hline 6 & 11 & \\ \hline 7 & 12 & \\ \hline 8 & & \\ \hline \end{array} \\ \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \\ \hline \end{array} \end{array}.$$

The column Garnir element \mathbf{g}_A is then a linear combination of products of the generators $\psi_6, \psi_7, \psi_8, \psi_9, \psi_{10}$; the exact expression for \mathbf{g}_A depends on the choice of e .

Now define the *column Specht module* $S_{\lambda|\kappa}$ to be the graded \mathcal{H}_α -module generated by the vector z_λ of degree $\text{codeg}(T_\lambda)$ subject to the following relations:

1. $e(i_\lambda)z_\lambda = z_\lambda$;
2. $y_r z_\lambda = 0$ for all $r = 1, \dots, n$;
3. $\psi_r z_\lambda = 0$ for all $r = 1, \dots, n-1$ such that $r \downarrow_{T_\lambda} r+1$;
4. $\mathbf{g}_A z_\lambda = 0$ for all column Garnir nodes $A \in \lambda$.

We may relax notation and just write S_λ , if the e -multicharge κ is understood. We shall mostly consider S_λ as an \mathcal{H}_n -module, by setting $\mathcal{H}_\beta S_\lambda = 0$ for $\beta \neq \alpha$. Thus we have \mathcal{H}_n -modules $S_{\lambda|\kappa}$ for all e -multicharges κ and all $\lambda \in \mathcal{P}_n^l$. The main purpose of this paper is to study the space of \mathcal{H}_n -homomorphisms $S_\lambda \rightarrow S_\mu$, for $\lambda, \mu \in \mathcal{P}_n^l$. The following result is obvious from the definitions.

Lemma 2.8. *Suppose $\lambda, \mu \in \mathcal{P}_n^l$, and let $\alpha = \text{cont}(\lambda)$. If $\text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu) \neq 0$, then $\text{cont}(\mu) = \alpha$ (and in particular $\text{def}(\lambda) = \text{def}(\mu)$), and $\text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu) = \text{Hom}_{\mathcal{H}_\alpha}(S_\lambda, S_\mu)$.*

We shall also need to consider row Specht modules; for these, the definitions are largely obtained by ‘conjugating’ the definitions for column Specht modules. Fix κ, λ and α as above. Say that a node $A = (r, c, m) \in [\lambda]$ is a *row Garnir node* if $(r+1, c, m) \in [\lambda]$, and define the *row Garnir belt*

$$\mathbf{B}^A = \{(r, d, m) \in [\lambda] \mid d \geq c\} \cup \{(r+1, d, m) \in [\lambda] \mid d \leq c\}.$$

This belt is used to define a *row Garnir element* \mathbf{g}^A . We refer the reader to [KMR, Definition 5.8] for the definition of this; here we just note the following facts:

- in T^λ the nodes of \mathbf{B}^A are occupied by the integers $a, a+1, \dots, b$ for some $a < b$;
- \mathbf{g}^A is a linear combination of products of the form $\psi_{i_1} \dots \psi_{i_d}$ where $a \leq i_1, \dots, i_d < b$;
- \mathbf{g}^A depends only on e, c, a and the length of the row containing A .

Now we can define the *row Specht module* S^λ , which is the graded \mathcal{H}_α -module generated by the vector z^λ of degree $\deg(T^\lambda)$ subject to the relations

1. $e(i^\lambda)z^\lambda = z^\lambda$;
2. $y_r z^\lambda = 0$ for all $r = 1, \dots, n$;
3. $\psi_r z^\lambda = 0$ for all $r = 1, \dots, n-1$ such that $r \rightarrow_{T^\lambda} r+1$;
4. $\mathbf{g}^A z^\lambda = 0$ for all row Garnir nodes $A \in \lambda$.

We define basis elements for the row and column Specht modules as follows. For each $T \in \text{Std}(\lambda)$ we fix a *preferred reduced expression* $s_{r_1} \dots s_{r_a}$ for the permutation w_T , and define $\psi_T := \psi_{r_1} \dots \psi_{r_a}$ and $v_T := \psi_T z^\lambda$. Similarly, we fix a preferred reduced expression $s_{t_1} \dots s_{t_b}$ for w^T , and set $\psi^T := \psi_{t_1} \dots \psi_{t_b}$ and $v^T := \psi^T z^\lambda$.

Note that the elements v_T and v^T may depend on the choice of preferred reduced expressions, since the ψ_r do not satisfy the braid relations. However, the following results are independent of the choices made.

Lemma 2.9 [KMR, Propositions 5.14 & 7.14]. *Suppose $\lambda \in \mathcal{P}_n^l$ and $T \in \text{Std}(\lambda)$. Then $\deg(v^T) = \deg(T)$ and $\deg(v_T) = \text{codeg}(T)$.*

Lemma 2.10 [KMR, Corollaries 6.24 & 7.20]. *Suppose $\lambda \in \mathcal{P}_n^l$. Then $\{v^T \mid T \in \text{Std}(\lambda)\}$ is an \mathbb{F} -basis for S^λ , and $\{v_T \mid T \in \text{Std}(\lambda)\}$ is an \mathbb{F} -basis for S_λ .*

In spite of the dependence of these bases on the choices of preferred reduced expressions, we refer to the bases $\{v^T \mid T \in \text{Std}(\lambda)\}$ and $\{v_T \mid T \in \text{Std}(\lambda)\}$ as the *standard bases* for S^λ and S_λ respectively.

For the remainder of this section we summarise some basic results about the action of \mathcal{H}_α on S_λ . Many of these results are cited from [BKW], where they are stated for row Specht modules. In this paper we concentrate as far as possible on column Specht modules, so we translate all the results to this setting. Throughout we fix $\lambda \in \mathcal{P}_n^l$, and let $\psi_1, \dots, \psi_{n-1}$ refer to the generators of \mathcal{H}_α , where $\alpha = \text{cont}(\lambda)$. Recall that if S, T are standard λ -tableaux, then we write $S \triangleright T$ to mean that $w_S \succ w_T$.

Lemma 2.11 [BKW, Theorem 4.10(i)]. Suppose $T \in \text{Std}(\lambda)$, and $s_{j_1} \dots s_{j_r}$ is any reduced expression for w_T . Then $\psi_{j_1} \dots \psi_{j_r} v_T - v_T$ is a linear combination of basis elements v_U for $U \triangleleft T$.

Lemma 2.12 [BKW, Lemma 4.9]. Suppose $T \in \text{Std}(\lambda)$ and that $j-1 \rightarrow_T j$ or $j-1 \downarrow_T j$. Then $\psi_{j-1} v_T$ is a linear combination of basis elements v_U for $U \triangleleft T$.

Lemma 2.13 [BKW, Lemma 4.8]. Suppose $T \in \text{Std}(\lambda)$ and $1 \leq i \leq n$. Then $y_i v_T$ is a linear combination of basis elements v_U for $U \triangleleft T$.

We'll use Lemmas 2.11 and 2.13 to prove the following similar result, which is suggested but not proved in the proof of [BKW, Theorem 4.10].

Lemma 2.14. Suppose $T \in \text{Std}(\lambda)$ and $j-1 \not\rightarrow_T j$. Then $\psi_{j-1} v_T$ is a linear combination of basis elements v_U for $U \trianglelefteq T$.

We begin with the following simple observation.

Lemma 2.15. Suppose $T \in \text{Std}(\lambda)$. Then $j-1 \not\rightarrow_T j$ if and only if w_T has a reduced expression beginning with s_{j-1} .

Proof. Both conditions are equivalent to the condition that $w_T^{-1}(j-1) > w_T^{-1}(j)$. □

Proof of Lemma 2.14. By Lemma 2.15, w_T has a reduced expression of the form $s_{j-1} s_{k_1} \dots s_{k_r}$. Using Lemma 2.11 we have

$$v_T = \psi_{j-1} \psi_{k_1} \dots \psi_{k_r} v_\lambda + \sum_{\substack{U \in \text{Std}(\lambda) \\ U \triangleleft T}} a_U v_U$$

for some $a_U \in \mathbb{F}$. So

$$\psi_{j-1} v_T = \psi_{j-1}^2 \psi_{k_1} \dots \psi_{k_r} v_\lambda + \sum_{\substack{U \in \text{Std}(\lambda) \\ U \triangleleft T}} a_U \psi_{j-1} v_U. \quad (*)$$

Using the KLR relations (and moving the appropriate idempotent $e(i)$ through), the first term on the right-hand side becomes $g \psi_{k_1} \dots \psi_{k_r} v_\lambda$, where g is a polynomial in y_1, \dots, y_n . Now $s_{k_1} \dots s_{k_r}$ is a reduced expression for the standard tableau $S = s_{j-1} T$, so by Lemma 2.11 we have

$$\psi_{k_1} \dots \psi_{k_r} v_\lambda = v_S + \sum_{\substack{V \in \text{Std}(\lambda) \\ V \triangleleft S}} b_V v_V$$

for some $b_V \in \mathbb{F}$. So (since $S \triangleleft T$) the first term on the right-hand side of (*) is a linear combination of terms of the form gv_V for $V \in \text{Std}(\lambda)$ with $V \triangleleft T$. By Lemma 2.13 this reduces to a linear combination of basis elements v_V for $V \triangleleft T$.

Now consider each of the remaining terms $\psi_{j-1}v_U$ in (*). If $j-1 \not\leq_U j$, then by induction on the Bruhat order $\psi_{j-1}v_U$ is a linear combination of basis elements v_V for $V \trianglelefteq U \triangleleft T$, so we can ignore any such U . If $j-1 \rightarrow_U j$ or $j-1 \downarrow_U j$, then we apply Lemma 2.12 to get the same conclusion. If $j-1 \nearrow_U j$, let R be the tableau obtained by swapping $j-1$ and j in U ; then a reduced expression for w_R may be obtained by adding s_{j-1} at the start of a reduced expression for w_U , and we have $R \trianglelefteq T$ by Lemma 2.2. So by Lemma 2.11 again,

$$\psi_{j-1}v_U = v_R + \sum_{W \triangleleft R} c_W v_W$$

for some $c_W \in \mathbb{F}$, and we are done. \square

Lemma 2.16. *Suppose $\lambda \in \mathcal{P}_n^l$, and $T \in \text{Std}(\lambda)$. Suppose $j_1, \dots, j_r \in \{1, \dots, n-1\}$, and that when $\psi_{j_1} \dots \psi_{j_r} z_\lambda$ is expressed as a linear combination of standard basis elements, v_T appears with non-zero coefficient. Then the expression $s_{j_1} \dots s_{j_r}$ has a reduced expression for w_T as a subexpression.*

Proof. We proceed by induction on r , with the case $r = 0$ trivial. Let $j = j_1$. Then by assumption v_T appears with non-zero coefficient in $\psi_j v_S$, where $S \in \text{Std}(\lambda)$ and v_S appears with non-zero coefficient in $\psi_{j_2} \dots \psi_{j_r} z_\lambda$. By induction the expression $s_{j_2} \dots s_{j_r}$ has a subexpression which is a reduced expression for w_S , so if $w_T \leq w_S$ (i.e. if $T \trianglelefteq S$) then we are done. By Lemma 2.12 and Lemma 2.14, this happens if $j \rightarrow_S j+1$, $j \downarrow_S j+1$ or $j \nearrow_S j+1$. So we can assume that $j \nearrow_S j+1$. But in this case $w_T = s_j w_S$, with $l(w_T) = l(w_S) + 1$, so w_T has a reduced expression obtained by adding s_j at the start of a reduced expression for w_S . So again the result follows by induction. \square

2.8 Specht modules for \mathcal{H}_n^κ and homomorphisms

Throughout this paper we consider the Specht module S_λ as a module for the *affine* algebra \mathcal{H}_α (where $\alpha = \text{cont}(\lambda)$) and by extension for the algebra \mathcal{H}_n . In fact, it is not hard to show that S_λ is annihilated by the element $y_1^{(\Lambda_\kappa | \alpha_{i_1})} e(i)$ for every i , so that S_λ is a module for the cyclotomic algebra \mathcal{H}_n^κ introduced in §2.6. We shall almost entirely be studying the space of \mathcal{H}_n^κ -homomorphisms between two Specht modules S_λ and S_μ defined for the same e -multicharge κ , and clearly in this situation \mathcal{H}_n^κ -homomorphisms between these two modules are the same as \mathcal{H}_n^κ -homomorphisms. In view of the Brundan–Kleshchev isomorphism theorem mentioned above, our results can therefore be viewed as statements about homomorphisms between Specht modules for (degenerate) Ariki–Koike algebras, and so they generalise the results of the first author and Lyle for homomorphisms between Specht modules for the symmetric group [FL, Theorem 2.1], and of Lyle and Mathas for Hecke algebras of type A [LM, Theorem 1.1].

In this paper, however, we restrict attention entirely to the affine algebra \mathcal{H}_n . This is because we occasionally (in particular, in Theorem 3.16) need to compare Specht modules defined for different e -multicharges.

3 λ -dominated tableaux and dominated homomorphisms

In this paper we consider the space of homomorphisms between two given Specht modules. However, our results concerning row and column removal will only apply to homomorphisms of a certain kind, which we call *dominated* homomorphisms. But as we shall see in Theorem 3.6, in many cases all homomorphisms between Specht modules are dominated.

3.1 λ -dominated tableaux

Suppose $\lambda, \mu \in \mathcal{P}_n^l$ and $T \in \text{Std}(\mu)$. Given $0 \leq j \leq n$, we say that T is λ -column-dominated on $1, \dots, j$ if each $i \in \{1, \dots, j\}$ appears at least as far to the left in T as it does in T_λ . We say simply that T is λ -column-dominated if it is λ -column-dominated on $1, \dots, n$. We remind the reader of our unusual convention for drawing Young diagrams, in which a node (r, c, m) lies to the left of (r', c', m') if either $m > m'$ or $(m = m' \text{ and } c \leq c')$.

We write $\text{Std}_\lambda(\mu)$ for the set of λ -column-dominated standard μ -tableaux. It is easy to see that $\text{Std}_\lambda(\mu)$ is non-empty if and only if $\lambda \trianglerighteq \mu$, and that $\text{Std}_\mu(\mu) = \{T_\mu\}$.

We say T is *weakly* λ -column-dominated on $1, \dots, j$ if each $i \in \{1, \dots, j\}$ appears in a component at least as far to the left in T as it does in T_λ . We say that T is weakly λ -column-dominated if it is weakly λ -column-dominated on $1, \dots, n$.

We also introduce row-dominance. Say that $T \in \text{Std}(\mu)$ is λ -row-dominated if each $i \in \{1, \dots, n\}$ appears at least as high in T as it does in T^λ . We write $\text{Std}^\lambda(\mu)$ for the set of λ -row-dominated standard μ -tableaux, which is non-empty if and only if $\lambda \trianglelefteq \mu$.

Since we shall primarily be considering column Specht modules, we shall often simply say ‘ λ -dominated’ meaning ‘ λ -column-dominated’.

We give a helpful alternative characterisation of the λ -dominated and λ -row-dominated properties.

Lemma 3.1. *Suppose $\lambda, \mu \in \mathcal{P}_n^l$, and $S \in \text{Std}(\mu)$.*

1. *S is λ -column-dominated on $1, \dots, j$ if and only if $\text{Shape}((T_\lambda)_{\downarrow m}) \trianglerighteq \text{Shape}(S_{\downarrow m})$ for all $m = 1, \dots, j$.*
2. *S is λ -row-dominated on $1, \dots, j$ if and only if $\text{Shape}((T^\lambda)_{\downarrow m}) \trianglelefteq \text{Shape}(S_{\downarrow m})$ for all $m = 1, \dots, j$.*

Proof. We prove only (2); the proof of (1) is analogous. Suppose first that S is not λ -row-dominated on $1, \dots, j$. Choose an entry $m \leq j$ which appears strictly lower in S than in T^λ , and let $\tau = \text{Shape}((T^\lambda)_{\downarrow m})$ and $\sigma = \text{Shape}(S_{\downarrow m})$. Suppose that m appears in position (r, c, k) in T^λ . The construction of T^λ means that the entries $1, \dots, m-1$ all appear at least as high as m in T^λ , and so

$$|\tau^{(1)}| + \dots + |\tau^{(k-1)}| + \tau_1^{(k)} + \dots + \tau_r^{(k)} = m.$$

On the other hand, m appears below row r of component k in S , so

$$|\sigma^{(1)}| + \dots + |\sigma^{(k-1)}| + \sigma_1^{(k)} + \dots + \sigma_r^{(k)} < m.$$

Hence $\tau \not\trianglelefteq \sigma$.

Conversely, suppose $\text{Shape}(T_{\downarrow m}^\lambda) \not\trianglelefteq \text{Shape}(S_{\downarrow m})$ for some $m \leq j$; choose such an m , and let $\tau = \text{Shape}(T_{\downarrow m}^\lambda)$ and $\sigma = \text{Shape}(S_{\downarrow m})$. Since $\tau \not\trianglelefteq \sigma$, there are r, k such that

$$|\tau^{(1)}| + \dots + |\tau^{(k-1)}| + \tau_1^{(k)} + \dots + \tau_r^{(k)} > |\sigma^{(1)}| + \dots + |\sigma^{(k-1)}| + \sigma_1^{(k)} + \dots + \sigma_r^{(k)}.$$

If we let $d = |\tau^{(1)}| + \dots + |\tau^{(k-1)}| + \tau_1^{(k)} + \dots + \tau_r^{(k)}$, then $d \leq m$ and the integers $1, \dots, d$ all appear in row r of component k or higher in T^λ . Since $|\sigma^{(1)}| + \dots + |\sigma^{(k-1)}| + \sigma_1^{(k)} + \dots + \sigma_r^{(k)} < d$, at least one of the integers $1, \dots, d$ appears in S below row r of component k . So there is some $i \leq j$ which appears lower in S than in T^λ , so S is not λ -row-dominated on $1, \dots, j$. \square

Corollary 3.2. Suppose $\lambda, \mu \in \mathcal{P}_n^l$, and $S, T \in \text{Std}(\mu)$.

1. If S is λ -dominated on $1, \dots, j$ and $S \triangleright T$, then T is λ -dominated on $1, \dots, j$. In particular, if $S \in \text{Std}_\lambda(\mu)$ and $S \triangleright T$, then $T \in \text{Std}_\lambda(\mu)$.
2. If S is λ -row-dominated on $1, \dots, j$ and $S \trianglelefteq T$, then T is λ -row-dominated on $1, \dots, j$. In particular, if $S \in \text{Std}^\lambda(\mu)$ and $S \trianglelefteq T$, then $T \in \text{Std}^\lambda(\mu)$.

Lemma 3.3. Suppose $\lambda, \mu \in \mathcal{P}_n^l$, and $T, U \in \text{Std}(\mu)$ with $U \trianglelefteq T$. If T is weakly λ -dominated on $1, \dots, j$, then so is U .

Proof. The proof follows almost identically to that of Corollary 3.2(1). \square

3.2 Dominated homomorphisms

Given $\lambda, \mu \in \mathcal{P}_n^l$, we want to consider the space of \mathcal{H}_n -homomorphisms $\varphi : S_\lambda \rightarrow S_\mu$ with the property that $\varphi(z_\lambda)$ lies in the \mathbb{F} -span of $\{v_S \mid S \in \text{Std}_\lambda(\mu)\}$. But we need to show that this notion is well-defined.

Proposition 3.4. Suppose $\lambda, \mu \in \mathcal{P}_n^l$. Then the subspace $\langle v_S \mid S \in \text{Std}_\lambda(\mu) \rangle_{\mathbb{F}}$ of S_μ is independent of the choice of standard basis elements v_S .

Proof. Let V denote the space $\langle v_S \mid S \in \text{Std}_\lambda(\mu) \rangle_{\mathbb{F}}$, and take $T \in \text{Std}_\lambda(\mu)$. Let $s_{j_1} \dots s_{j_r}$ be a new reduced expression for w_T , and let $v'_T = \psi_{j_1} \dots \psi_{j_r} z_\mu$ (where $\psi_1, \dots, \psi_{n-1}$ are taken to lie in $\mathcal{H}_{\text{cont}(\lambda)}$). Let V' be the space obtained from V by replacing v_T with v'_T in the spanning set $\{v_S \mid S \in \text{Std}_\lambda(\mu)\}$; it suffices to show that $V = V'$. By Lemma 2.11,

$$v'_T = v_T + \sum_{U \triangleleft T} a_U v_U \quad \text{for some } a_U \in \mathbb{F}.$$

By Corollary 3.2(1), each v_U with $U \triangleleft T$ lies in V , and so $v'_T \in V$. Hence $V' \subseteq V$; but since the elements v_S are linearly independent, $\dim_{\mathbb{F}} V = \dim_{\mathbb{F}} V' = |\text{Std}_\lambda(\mu)|$. So $V' = V$. \square

In view of Proposition 3.4 and an analogue for row-dominated tableaux, the following definition makes sense.

Definition. Suppose $\lambda, \mu \in \mathcal{P}_n^l$. If $\varphi \in \text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$, we say that φ is (column-)dominated if $\varphi(z_\lambda) \in \langle v_S \mid S \in \text{Std}_\lambda(\mu) \rangle_{\mathbb{F}}$. We write $\text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$ for the space of dominated homomorphisms from S_λ to S_μ .

Similarly, if $\chi \in \text{Hom}_{\mathcal{H}_n}(S^\lambda, S^\mu)$, we say that χ is row-dominated if $\chi(z^\lambda) \in \langle v^S \mid S \in \text{Std}^\lambda(\mu) \rangle_{\mathbb{F}}$, and we write $\text{DHom}_{\mathcal{H}_n}(S^\lambda, S^\mu)$ for the space of row-dominated homomorphisms from S^λ to S^μ .

Proposition 3.5. $\text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$ and $\text{DHom}_{\mathcal{H}_n}(S^\lambda, S^\mu)$ are graded subspaces of $\text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$ and $\text{Hom}_{\mathcal{H}_n}(S^\lambda, S^\mu)$ respectively. That is, $\text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$ and $\text{DHom}_{\mathcal{H}_n}(S^\lambda, S^\mu)$ are spanned by homogeneous homomorphisms.

Proof. The proof proceeds almost identically to the proof of the fact that $\text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$ is graded, using the fact that $\langle v_S \mid S \in \text{Std}_\lambda(\mu) \rangle_{\mathbb{F}}$ is a graded subspace of S_μ (with a corresponding statement for S^μ). \square

The rest of this section is devoted to showing that in certain cases every Specht homomorphism is dominated. Specifically, we shall prove the following.

Theorem 3.6. *Suppose $e \neq 2$ and that $\kappa_1, \dots, \kappa_l$ are distinct. Then $\text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu) = \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$.*

Remark. The hypotheses that $e \neq 2$ and that $\kappa_1, \dots, \kappa_l$ are distinct are equivalent to the condition that \mathcal{H}_n^κ has exactly $2l$ isomorphism classes of one-dimensional modules. The following small examples show that these hypotheses are essential in Theorem 3.6; in fact, they show that Specht modules labelled by different multipartitions can be isomorphic without these assumptions.

1. Take $e = 2$, $\kappa = (0)$, $\lambda = ((1^2))$ and $\mu = ((2))$. Then there is a non-zero homomorphism $S_\lambda \rightarrow S_\mu$ defined by $z_\lambda \mapsto z_\mu$, though the tableau $T_\mu = \begin{bmatrix} 1 & 2 \end{bmatrix}$ is not λ -dominated. So $\text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu) \neq \{0\} = \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$.
2. For any e , take $\kappa = (0, 0)$, $\lambda = (\emptyset, (1))$ and $\mu = ((1), \emptyset)$. Then $z_\lambda \mapsto z_\mu$ again defines a non-zero homomorphism $S_\lambda \rightarrow S_\mu$, though T_μ is not λ -dominated.

The proof of Theorem 3.6 requires several preliminary results. We fix $\lambda, \mu \in \mathcal{P}_n^l$ and an e -multicharge κ of level l throughout. If $\text{cont}(\lambda) \neq \text{cont}(\mu)$, then by Lemma 2.8 $\text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu) = 0$, so that Theorem 3.6 is trivially true. So we assume that $\text{cont}(\lambda) = \text{cont}(\mu)$. In the results below, $\psi_1, \dots, \psi_{n-1}$ are elements of $\mathcal{H}_{\text{cont}(\lambda)}$.

Lemma 3.7. *Suppose $j \in \{2, \dots, n\}$ with $j-1 \downarrow_{T_\lambda} j$, and $T \in \text{Std}(\mu)$ is λ -dominated on $1, \dots, j$. Then $\psi_{j-1}v_T$ is a linear combination of basis elements v_U for standard tableaux U which are λ -dominated on $1, \dots, j$.*

Proof. If $j-1 \rightarrow_T j$ or $j-1 \downarrow_T j$ or $j-1 \nearrow_T j$, then the result follows from Corollary 3.2(1) together with either Lemma 2.12 or Lemma 2.14. The remaining possibility is that $j-1 \nearrow_T j$. But now if we let S be the standard tableau $s_{j-1}T$, then by Lemma 2.11 $\psi_{j-1}v_T = v_S + \sum_{U \prec S} b_U v_U$ for some $b_U \in \mathbb{F}$. Clearly since T is λ -dominated on $1, \dots, j$ and $j-1, j$ lie in the same column of T_λ , S is also λ -dominated on $1, \dots, j$. Corollary 3.2(1) completes the proof. \square

Proposition 3.8. *Suppose $e \neq 2$, and that $\varphi : S_\lambda \rightarrow S_\mu$ is a homomorphism, and write*

$$\varphi(z_\lambda) = \sum_{T \in \text{Std}(\mu)} a_T v_T \quad \text{for some } a_T \in \mathbb{F}.$$

Suppose $j \in \{2, \dots, n\}$ with $j-1 \downarrow_{T_\lambda} j$, and that each T for which $a_T \neq 0$ is λ -dominated on $1, \dots, j-1$. Then each T for which $a_T \neq 0$ is λ -dominated on $1, \dots, j$.

Proof. The fact that $j-1 \downarrow_{T_\lambda} j$ means that $\psi_{j-1}z_\lambda = 0$, so we must have $\sum_{T \in \text{Std}(\mu)} a_T \psi_{j-1}v_T = 0$. Assuming the proposition is false, there is at least one T which is not λ -dominated on $1, \dots, j$ such that $a_T \neq 0$; choose such a T which is \triangleright -maximal. Since T is λ -dominated on $1, \dots, j-1$, the entry j lies in a column strictly to the right of $j-1$ in T . We claim that we cannot have $j-1 \rightarrow_T j$. If this is the case, then the residue sequence $i(T)$ satisfies $i(T)_j = i(T)_{j-1} + 1$. However, since f

is a homomorphism and v_T appears with non-zero coefficient in $\varphi(z_\lambda)$, we must have $i(T) = i_\lambda$, and the fact that $j-1 \downarrow_{T_\lambda} j$ means that $(i_\lambda)_j = (i_\lambda)_{j-1} - 1$. Since $e \neq 2$, this is a contradiction.

Hence $j-1 \nearrow_T j$, so the tableau $S := s_{j-1}T$ is standard, and if we write $\psi_{j-1}v_T$ as a linear combination of standard basis elements, then v_S occurs with coefficient 1. We claim that v_S does not occur in any other $\psi_{j-1}v_{T'}$ when $a_{T'} \neq 0$: if T' is not λ -dominated on $1, \dots, j$, then (defining S' analogously to S) we have $\psi_{j-1}v_{T'} = v_{S'} + \sum_{U \triangleleft S'} c_U v_U$ for some $c_U \in \mathbb{F}$; but the fact that $T \not\triangleleft T'$ (by our choice of T being \triangleright -maximal) means that $S \not\triangleleft S'$, so v_S cannot occur. On the other hand, if T' is λ -dominated on $1, \dots, j$, then the result follows from Lemma 3.7, since S is not λ -dominated on $1, \dots, j$.

So v_S occurs with non-zero coefficient in $\sum_{T \in \text{Std}(\mu)} a_T \psi_{j-1}v_T$, a contradiction. \square

We now turn our attention to the case where j is in the top row of its component in T_λ .

Lemma 3.9. *Suppose $1 \leq a \leq j \leq n$, and that $j-1 \nearrow_{T_\lambda} j$ and $a \downarrow_{T_\lambda} a+i$ for all $i = 1, \dots, j-a-1$. If $T \in \text{Std}(\mu)$ is weakly λ -dominated on $1, \dots, j$ then $\psi_a \psi_{a+1} \dots \psi_{j-1}v_T$ is a linear combination of basis elements v_U for standard tableaux U which are weakly λ -dominated on $1, \dots, j$.*

Proof. We argue by induction on $l(s_a s_{a+1} \dots s_{j-1}) = j-a$. If $j-a = 0$, the result is trivial. So suppose $a < j$, and assume by induction that $\psi_{a+1} \dots \psi_{j-1}v_T$ is a linear combination of basis elements v_U which are weakly λ -dominated on $1, \dots, j$. We want to show that for each v_U , $\psi_a v_U$ is a linear combination of basis elements $v_{U'}$ for standard tableaux U' which are weakly λ -dominated on $1, \dots, j$.

If $a \rightarrow_U a+1$ or $a \downarrow_U a+1$ or $a \nearrow_U a+1$, then the result follows from Lemma 3.3 together with either Lemma 2.12 or Lemma 2.14. The remaining possibility is that $a \nearrow_U a+1$. Let S be the standard tableau $s_a U$. Then by Lemma 2.11, $\psi_a v_U = v_S + \sum_{U' \triangleleft S} a_{U'} v_{U'}$ for some $a_{U'} \in \mathbb{F}$.

Recalling that U is weakly λ -dominated on $1, \dots, j$ and that $a, a+1$ are in the same column (and therefore the same component) of T_λ , S is weakly λ -dominated on $1, \dots, j$ and Lemma 3.3 completes the proof. \square

Proposition 3.10. *Suppose $\varphi : S_\lambda \rightarrow S_\mu$ is a homomorphism with*

$$\varphi(z_\lambda) = \sum_{T \in \text{Std}(\mu)} a_T v_T \quad \text{for some } a_T \in \mathbb{F}.$$

Suppose $j \in \{2, \dots, n\}$ with either $j-1 \nearrow_{T_\lambda} j$ or $j-1 \rightarrow_{T_\lambda} j$, and that each T for which $a_T \neq 0$ is λ -dominated on $1, \dots, j-1$. Then each T for which $a_T \neq 0$ is λ -dominated on $1, \dots, j$.

Proof. The proof follows the same lines as Proposition 3.8. The condition that $j-1 \nearrow_{T_\lambda} j$ or $j-1 \rightarrow_{T_\lambda} j$ means that S_λ satisfies a Garnir relation $\psi_a \psi_{a+1} \dots \psi_{j-1} z_\lambda = 0$, where a is the entry immediately to the left of j in T_λ ; since f is a homomorphism, we therefore have $\sum_{T \in \text{Std}(\mu)} a_T \psi_a \dots \psi_{j-1}v_T = 0$. Assuming the result is false, there is at least one T which is not λ -dominated on $1, \dots, j$ such that $a_T \neq 0$; choose such a T which is \triangleright -maximal. Since T is λ -dominated on $1, \dots, j-1$ but not $1, \dots, j$, we have $j-1 \nearrow_T j$. In fact $j-1$ and j are in different components of T : if not, what is the entry immediately to the left of j in T ? It must be some $k < j$, since T is standard, but by assumption k is strictly left of j in T_λ and hasn't moved to the right in T .

Let S denote the standard tableau $s_a s_{a+1} \dots s_{j-1} T$. Then $l(w_S) = l(w_T) + j-a$, so that when we write $\psi_a \psi_{a+1} \dots \psi_{j-1}v_T$ as a linear combination of standard basis elements, v_S occurs with

coefficient 1. We claim that v_S does not occur with non-zero coefficient in $\psi_a \psi_{a+1} \dots \psi_{j-1} v_{T'}$ for any other T' with $a_{T'} \neq 0$: if T' is not λ -dominated on $1, \dots, j$, then (defining S' analogously to S) we have $\psi_a \psi_{a+1} \dots \psi_{j-1} v_{T'} = v_{S'} + \sum_{U \triangleleft S'} b_U v_U$ for some $b_U \in \mathbb{F}$; but the fact that $T \not\triangleleft T'$ (by our choice of T) means that $S \not\triangleleft S'$, so v_S cannot occur. On the other hand, if T' is λ -dominated on $1, \dots, j$, then the result follows from Lemma 3.9, since S is not weakly λ -dominated on $1, \dots, j$ as $j-1$ and j are in different components of T .

So v_S occurs with non-zero coefficient in $\sum_{T \in \text{Std}(\mu)} a_T \psi_a \psi_{a+1} \dots \psi_{j-1} v_T$, a contradiction. \square

The last thing we need for the proof of Theorem 3.6 is the following.

Lemma 3.11. *Suppose $\kappa_1, \dots, \kappa_l$ are distinct, and that $T \in \text{Std}(\mu)$ satisfies $i(T) = i_\lambda$. If T is λ -dominated on $1, \dots, j-1$ and j appears in the $(1,1)$ -position of its component in T_λ , then T is λ -dominated on $1, \dots, j$.*

Proof. Suppose not; then j appears in T strictly to the right of where it appears in T_λ . This means that j must appear in the $(1,1)$ -node of some component of T , since otherwise there would be a smaller entry immediately above or to the left of j , contradicting the assumption that T is λ -dominated on $1, \dots, j-1$.

So there are $1 \leq r < s \leq l$ such that $T_\lambda(1,1,s) = j = T(1,1,r)$. Hence $\kappa_s = (i_\lambda)_j = i(T)_j = \kappa_r$, contrary to assumption. \square

Proof of Theorem 3.6. Suppose $\varphi : S_\lambda \rightarrow S_\mu$ is a homomorphism, and write

$$\varphi(z_\lambda) = \sum_{T \in \text{Std}(\mu)} a_T v_T \quad \text{for some } a_T \in \mathbb{F}.$$

We must show that every T for which $a_T \neq 0$ is λ -dominated. In fact we show by induction on j that every such T is λ -dominated on $1, \dots, j$, with the case $j = 0$ being vacuous. So suppose $j \geq 1$, and assume by induction that T is λ -dominated on $1, \dots, j-1$. Note that since φ is a homomorphism, we have $i(T) = i_\lambda$.

If $j = 1$ or j lies in an earlier component of T_λ than $j-1$, then j lies in the $(1,1)$ -node of its component in T_λ . So by Lemma 3.11 T is λ -dominated on $1, \dots, j$. The remaining possibilities are that $j > 1$ and that one of

$$j-1 \downarrow_{T_\lambda} j, \quad j-1 \rightarrow_{T_\lambda} j, \quad j-1 \nearrow_{T_\lambda} j$$

occurs; these cases are dealt with in Propositions 3.8 and 3.10. \square

We immediately see the following interesting result.

Corollary 3.12. *Suppose $e \neq 2$ and that $\kappa_1, \dots, \kappa_l$ are distinct. If $\lambda, \mu \in \mathcal{P}_n^l$ with $\text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu) \neq \{0\}$, then $\lambda \triangleright \mu$. Furthermore (since $\text{Std}_\lambda(\lambda) = \{T_\lambda\}$) $\text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\lambda)$ is one-dimensional. In particular, S_λ is indecomposable.*

Remarks.

1. The authors thank an anonymous referee for pointing out that in the case where $e = \infty$ or \mathbb{F} contains a primitive e th root of unity, Corollary 3.12 can be deduced from results of Rouquier. In this case \mathcal{H}_n is isomorphic to an Ariki–Koike algebra defined at an e th root of unity, and [R1, Theorem 6.6] shows that under precisely the same hypotheses as

Corollary 3.12 there is a cyclotomic q -Schur algebra which is a 1-faithful quasi-hereditary cover of the Ariki–Koike algebra. This means that the Schur functor for these algebras maps the space of homomorphisms between Weyl modules bijectively to the space of homomorphisms between the corresponding Specht modules. Since the cyclotomic q -Schur algebra is quasi-hereditary, the conclusions of Corollary 3.12 hold for Weyl modules, and hence they hold for Specht modules too.

2. If $e = 2$ then S_λ may be decomposable. For example, when $l = 1$ and $\text{char}(\mathbb{F}) \neq 3$, the Specht module $S_{((5,12))}$ is decomposable; this was shown in [J, Example 23.10(iii)] in the case $\text{char}(\mathbb{F}) = 2$, and in [S, Theorem 6.8] in odd characteristic. Similarly, when $\kappa_i = \kappa_j$ for some $i \neq j$, we can have decomposable Specht modules: take $\kappa = (0, 0)$, $e = 3$ and $\text{char}(\mathbb{F}) \neq 2$; then $S_{((3),(3))}$ is decomposable.

In exactly the same way, we can prove the corresponding result for row Specht modules.

Theorem 3.13. *Suppose $e \neq 2$ and that $\kappa_1, \dots, \kappa_l$ are distinct, and $\lambda, \mu \in \mathcal{P}_n^l$. Then $\text{DHom}_{\mathcal{H}_n}(S^\lambda, S^\mu) = \text{Hom}_{\mathcal{H}_n}(S^\lambda, S^\mu)$. Hence $\text{Hom}_{\mathcal{H}_n}(S^\lambda, S^\mu) \neq \{0\}$ only if $\lambda \leq \mu$, $\text{Hom}_{\mathcal{H}_n}(S^\lambda, S^\lambda)$ is one-dimensional, and S^λ is indecomposable.*

3.3 Duality for dominated homomorphisms

In this section we consider the relationship between row and column Specht modules, as well as the relationship between Specht modules labelled by conjugate multipartitions. These relationships are encapsulated in [KMR, Theorems 7.25 and 8.5], from which it follows that a (generalised) column-removal theorem for homomorphisms between Specht modules is equivalent to the corresponding row-removal theorem. The main result of this section, which requires considerable additional work, is that the same is true for dominated homomorphisms.

Following [KMR, §3.2], let $\tau : \mathcal{H}_\alpha \rightarrow \mathcal{H}_\alpha$ denote the anti-automorphism which fixes all the generators $e(i), y_r, \psi_s$, and define $\tau : \mathcal{H}_n \rightarrow \mathcal{H}_n$ by combining these maps for all α . If $M = \bigoplus_{d \in \mathbb{Z}} M_d$ is a graded \mathcal{H}_n -module, let M^* denote the graded module with $M_d^* = \text{Hom}_{\mathbb{F}}(M_{-d}, \mathbb{F})$ for each d , with \mathcal{H}_n -action given by $(hf)m = f(\tau(h)m)$ for $m \in M$, $f \in M^*$ and $h \in \mathcal{H}_n$. Also, for $k \in \mathbb{Z}$, let $M\langle k \rangle$ denote the same module with the grading shifted by k , i.e. $M\langle k \rangle_d = M_{d-k}$. Finally, recall the defect $\text{def}(\lambda)$ of a multipartition from Section 2.5.

Theorem 3.14 [KMR, Theorem 7.25]. *Suppose $\lambda \in \mathcal{P}_n^l$. Then*

$$S^\lambda \cong (S_\lambda)^* \langle \text{def}(\lambda) \rangle \quad \text{and} \quad S_\lambda \cong (S^\lambda)^* \langle \text{def}(\lambda) \rangle.$$

Now suppose $\lambda, \mu \in \mathcal{P}_n^l$. Applying Theorem 3.14 to both λ and μ gives an isomorphism of graded vector spaces

$$\text{Hom}_{\mathcal{H}_n}(S^\mu, S^\lambda) \cong \text{Hom}_{\mathcal{H}_n}(S_\mu^* \langle \text{def}(\mu) \rangle, S_\lambda^* \langle \text{def}(\lambda) \rangle);$$

since by Lemma 2.8 $\text{def}(\lambda) = \text{def}(\mu)$ for any λ and μ with $\text{Hom}_{\mathcal{H}_n}(S^\mu, S^\lambda) \neq \{0\}$, this yields an isomorphism of graded vector spaces

$$\text{Hom}_{\mathcal{H}_n}(S^\mu, S^\lambda) \cong \text{Hom}_{\mathcal{H}_n}(S_\mu^*, S_\lambda^*).$$

τ is a homogeneous map of degree zero, so $\text{Hom}_{\mathcal{H}_n}(S_\mu^\otimes, S_\lambda^\otimes)$ is canonically isomorphic as a graded vector space to $\text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$, and hence we have an isomorphism of graded vector spaces

$$\Theta : \text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu) \xrightarrow{\sim} \text{Hom}_{\mathcal{H}_n}(S^\mu, S^\lambda).$$

Our aim is to prove the following.

Proposition 3.15. *Suppose $\lambda, \mu \in \mathcal{P}_n^l$, and let $\Theta : \text{Hom}_{\mathcal{H}_n}(S_\lambda, S_\mu) \rightarrow \text{Hom}_{\mathcal{H}_n}(S^\mu, S^\lambda)$ be the bijection above. Then $\Theta(\text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)) = \text{DHom}_{\mathcal{H}_n}(S^\mu, S^\lambda)$.*

We shall prove Proposition 3.15 below. First we examine the consequences for row and column removal. In order to be able to compare row and column removal, we combine Proposition 3.15 with a result which relates to an analogue of the sign representation of the symmetric group. Following [KMR, §3.3], let $\text{sgn} : \mathcal{H}_\alpha \rightarrow \mathcal{H}_\alpha$ denote the automorphism which maps $e(i) \mapsto e(-i)$, $y_r \mapsto -y_r$ and $\psi_s \mapsto -\psi_s$ for all i, r, s , and define $\text{sgn} : \mathcal{H}_n \rightarrow \mathcal{H}_n$ by combining these maps for all α . Given a graded \mathcal{H}_n -module M , let M^{sgn} denote the same graded vector space with the action of \mathcal{H}_n twisted by sgn .

Recall that if λ is a multipartition, then λ' denotes the conjugate multipartition to λ , and that if $S \in \text{Std}(\lambda)$, then $S' \in \text{Std}(\lambda')$ denotes the conjugate tableau to S . Also define the *conjugate e -multicharge* $\kappa' := (-\kappa_l, \dots, -\kappa_1)$. Now the following is immediate from the construction of row and column Specht modules.

Theorem 3.16 [KMR, Theorem 8.5]. *Suppose $\lambda \in \mathcal{P}_n^l$. Then there is an isomorphism $(S^{\lambda|\kappa})^{\text{sgn}} \cong S_{\lambda'|\kappa'}$ of \mathcal{H}_n -modules, given by $v^S \mapsto v_{S'}$.*

Remark. Theorem 3.16 is one place where it is essential that we consider Specht modules as modules for \mathcal{H}_n , rather than its cyclotomic quotients, since the two modules involved are defined relative to different e -multicharges.

Now suppose $\lambda, \mu \in \mathcal{P}_n^l$. Since sgn is a homogeneous automorphism of \mathcal{H}_n , we have an equality of graded vector spaces

$$\text{Hom}_{\mathcal{H}_n}((S^{\mu|\kappa})^{\text{sgn}}, (S^{\lambda|\kappa})^{\text{sgn}}) = \text{Hom}_{\mathcal{H}_n}(S^{\mu|\kappa}, S^{\lambda|\kappa}), \quad (*)$$

Combining this with Theorem 3.16, we have an isomorphism of graded vector spaces

$$\text{Hom}_{\mathcal{H}_n}(S_{\mu'|\kappa'}, S_{\lambda'|\kappa'}) \cong \text{Hom}_{\mathcal{H}_n}(S^{\mu|\kappa}, S^{\lambda|\kappa}). \quad (\dagger)$$

Applying Theorem 3.14 yields an isomorphism of graded vector spaces

$$\text{Hom}_{\mathcal{H}_n}(S_{\mu'|\kappa'}, S_{\lambda'|\kappa'}) \cong \text{Hom}_{\mathcal{H}_n}(S_{\lambda|\kappa}, S_{\mu|\kappa}). \quad (\ddagger)$$

We want to show that the same holds for dominated homomorphisms; this is immediate when $e > 2$ and $\kappa_1, \dots, \kappa_l$ are distinct, by Theorem 3.6. In general, we observe that $(*)$ remains true with Hom replaced by DHom , and the explicit form of the isomorphism in Theorem 3.16 shows that (\dagger) does too, since $S \in \text{Std}_{\mu'}(\lambda')$ if and only if $S' \in \text{Std}^\mu(\lambda)$. Finally, Proposition 3.15 shows that (\ddagger) remains true for DHom too. So we have the following theorem.

Theorem 3.17. *Suppose $\lambda, \mu \in \mathcal{P}_n^l$. Then there is an isomorphism of graded vector spaces*

$$\text{DHom}_{\mathcal{H}_n}(S_{\lambda|\kappa}, S_{\mu|\kappa}) \cong \text{DHom}_{\mathcal{H}_n}(S_{\mu'|\kappa'}, S_{\lambda'|\kappa'}).$$

It remains to prove Proposition 3.15; for the remainder of this section, all Specht modules are defined for the e -multicharge κ .

We begin by recalling how the isomorphism $S^\lambda \cong S_\lambda^\otimes \langle \text{def}(\lambda) \rangle$ in Theorem 3.14 is constructed. Given the standard basis $\{v_T \mid T \in \text{Std}(\lambda)\}$ for S_λ , let $\{f^T \mid T \in \text{Std}(\lambda)\}$ be the dual basis for S_λ^\otimes ; although the elements f^T in general depend on the choice of the elements v_T (i.e. on the choice of preferred reduced expressions), it is an easy exercise to show that the element f^{T^λ} does not. The isomorphism $S^\lambda \rightarrow S_\lambda^\otimes \langle \text{def}(\lambda) \rangle$ is defined by $z^\lambda \mapsto f^{T^\lambda}$.

Lemma 3.18. *Suppose $\lambda \in \mathcal{P}_n^l$, and let $\theta^\lambda : S^\lambda \rightarrow S_\lambda^\otimes \langle \text{def}(\lambda) \rangle$ be the isomorphism constructed above.*

1. *For any $S \in \text{Std}(\lambda)$ we have $\theta^\lambda(v^S) \in \langle f^T \mid T \in \text{Std}(\lambda), T \triangleright S \rangle_{\mathbb{F}}$.*
2. *θ^λ maps the space $\langle v^S \mid S \in \text{Std}^\mu(\lambda) \rangle_{\mathbb{F}}$ bijectively to the space $\langle f^S \mid S \in \text{Std}^\mu(\lambda) \rangle_{\mathbb{F}}$.*

Proof.

1. For each $T \in \text{Std}(\lambda)$, write $\tau(\psi^S)v_T = \sum_{U \in \text{Std}(\lambda)} a_{TU}v_U$. Then one can check that the definitions give $\theta^\lambda(v^S) = \sum_{T \in \text{Std}(\lambda)} a_{TT^\lambda} f^T$. So it suffices to show that $a_{TT^\lambda} = 0$ when $T \not\triangleright S$. Clearly to prove this it is sufficient to show this in the case where $\mathbb{F} = \mathbb{C}$, and so (as in the proof of [KMR, Theorem 7.25]) we can invoke the proof of [HM, Proposition 6.19]; here θ^λ is given in the form $x \mapsto \{x, -\}$, for a bilinear form $\{, \} : S^\lambda \times S_\lambda \langle \text{def}(\lambda) \rangle \rightarrow \mathbb{C}$ satisfying $\{v^S, v_T\} = 0$ unless $T \triangleright S$, which is exactly what we want.
2. From (1) and Corollary 3.2(2) we have $\theta^\lambda(v^S) \in \langle f^T \mid T \in \text{Std}^\mu(\lambda) \rangle_{\mathbb{F}}$ whenever $S \in \text{Std}^\mu(\lambda)$, so $\theta^\lambda(\langle v^S \mid S \in \text{Std}^\mu(\lambda) \rangle) \subseteq \langle f^S \mid S \in \text{Std}^\mu(\lambda) \rangle$. But θ^λ is an isomorphism of vector spaces and

$$\dim_{\mathbb{F}} \langle v^S \mid S \in \text{Std}^\mu(\lambda) \rangle_{\mathbb{F}} = |\text{Std}^\mu(\lambda)| = \dim_{\mathbb{F}} \langle f^S \mid S \in \text{Std}^\mu(\lambda) \rangle_{\mathbb{F}},$$

so in fact $\theta^\lambda(\langle v^S \mid S \in \text{Std}^\mu(\lambda) \rangle) = \langle f^S \mid S \in \text{Std}^\mu(\lambda) \rangle_{\mathbb{F}}$. □

Lemma 3.19. *Suppose $\lambda, \mu \in \mathcal{P}_n^l$. Suppose $S \in \text{Std}(\lambda)$ and U is a λ -tableau such that $w_S \succ w_U$ and that for every $1 \leq i \leq n$ the number i appears in U weakly to the right of where it appears in T^μ . Then $S \in \text{Std}^\mu(\lambda)$.*

Proof. Using Lemma 3.1(2) we just need to show that $\text{Shape}(S_{\downarrow m}) \triangleright \text{Shape}(T_{\downarrow m}^\mu)$ for all m . Let U_c be the column-strict tableau which is column-equivalent to U . Then by Proposition 2.1 $w_U \succ w_{U_c}$. By Proposition 2.5, we have that $\text{Shape}((U_c)_{\downarrow m})' \triangleright \text{Shape}(S_{\downarrow m})'$ for all m . Furthermore, the condition that every entry in U_c lies weakly to the right of where it lies in T^μ is equivalent to every entry in $(U_c)'$ lying weakly below where it lies in $(T^\mu)'$, so we necessarily have that $\text{Shape}((U_c)_{\downarrow m})' \trianglelefteq \text{Shape}(T_{\downarrow m}^\mu)'$ for all m . Reapplying Proposition 2.5, we have $w_{U_c} \succ w_{T^\mu}$. □

Lemma 3.20. *Suppose $\lambda, \mu \in \mathcal{P}_n^l$, $S \in \text{Std}(\lambda) \setminus \text{Std}^\mu(\lambda)$ and $T \in \text{Std}_\lambda(\mu)$. Then when $\psi_S v_T$ is expressed in terms of the standard basis $\{v_U \mid U \in \text{Std}(\mu)\}$, the coefficient of v_{T^μ} is zero.*

Proof. Suppose to the contrary that v_{T^μ} does appear with non-zero coefficient in $\psi_S v_T = \psi_S \psi_T z_\mu$. Let $s_{i_1} \dots s_{i_a}$ and $s_{j_1} \dots s_{j_b}$ be the preferred reduced expressions for w_S and w_T respectively. Then by Lemma 2.16 there is a reduced expression for w_{T^μ} occurring as a subexpression of

$s_{i_1} \dots s_{i_a} s_{j_1} \dots s_{j_b}$. If we separate this reduced expression into two parts, which occur as subexpressions of $s_{i_1} \dots s_{i_a}$ and $s_{j_1} \dots s_{j_b}$ respectively, and let w, x denote the corresponding elements of \mathfrak{S}_n , then we have

$$w \preceq w_S, x \preceq w_T, wx = w_{T^\mu}, l(w) + l(x) = l(w_{T^\mu}).$$

Putting $V = xT_\mu$, we have $V \in \text{Std}(\mu)$ by Lemma 2.3, and in fact $V \in \text{Std}_\lambda(\mu)$ (using Corollary 3.2(1), because $w_V \preceq w_T$ and $T \in \text{Std}_\lambda(\mu)$). If we let $U = wT_\lambda$ then, as functions $[\mu] \rightarrow [\lambda]$,

$$U^{-1}T^\mu = T_\lambda^{-1}xT_\mu = T_\lambda^{-1}V.$$

The fact that V is λ -dominated can be expressed as saying that the map $T_\lambda^{-1}V : [\mu] \rightarrow [\lambda]$ maps any node of μ to a node weakly to the right. So each entry of U appears weakly to the right of where it appears in T^μ , i.e. U satisfies the hypotheses of Lemma 3.19. Hence by Lemma 3.19 $S \in \text{Std}^\mu(\lambda)$, contrary to hypothesis. \square

Proof of Proposition 3.15. We shall prove that $\Theta(\text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)) \subseteq \text{DHom}_{\mathcal{H}_n}(S^\mu, S^\lambda)$; the same argument with λ and μ interchanged and with row and column Specht modules interchanged proves the opposite containment.

Suppose $\varphi \in \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$, and write $\varphi(z_\lambda) = \sum_{T \in \text{Std}_\lambda(\mu)} a_T v_T$ for some $a_T \in \mathbb{F}$. Let $\varphi^* : S_\mu^* \rightarrow S_\lambda^*$ denote the dual map. We want to show that the homomorphism $\Theta(\varphi)$ which corresponds to φ^* via Theorem 3.14 is row-dominated, i.e. $\Theta(\varphi)(z^\mu) \in \langle v^S \mid S \in \text{Std}^\mu(\lambda) \rangle_{\mathbb{F}}$. By the construction of the isomorphism $S^\mu \rightarrow S_\mu^*$ and by Lemma 3.18, this is the same as saying that $\varphi^*(f^{T^\mu}) \in \langle f^S \mid S \in \text{Std}^\mu(\lambda) \rangle_{\mathbb{F}}$; in other words, $\varphi^*(f^{T^\mu})(v_S) = 0$ when $S \in \text{Std}(\lambda) \setminus \text{Std}^\mu(\lambda)$.

φ^* is given by $f \mapsto f \circ \varphi$. In particular, $\varphi^*(f^{T^\mu}) = f^{T^\mu} \circ \varphi$, which maps v_S to the coefficient of v_{T^μ} in $\varphi(v_S) = \sum_{T \in \text{Std}_\lambda(\mu)} a_T \psi_S v_T$. By Lemma 3.20 this coefficient is zero when $S \notin \text{Std}^\mu(\lambda)$, and the result follows. \square

4 Column removal for homomorphisms

Now we come to the main results of the paper, which give row- and column-removal theorems for dominated homomorphisms between Specht modules.

4.1 Generalised column removal for multipartitions


Definition. Suppose $\lambda = (\lambda^{(1)}, \dots, \lambda^{(l)}) \in \mathcal{P}_n^l$. For any $1 \leq m \leq l$ and any $c \geq 0$, define $\lambda_L^{(m),c}$ to be the partition consisting of all nodes in the first c columns of $\lambda^{(m)}$, and $\lambda_R^{(m),c}$ the partition consisting of all nodes after the first c columns of $\lambda^{(m)}$. That is,

$$(\lambda_L^{(m),c})_i = \min\{\lambda_i^{(m)}, c\}, \quad (\lambda_R^{(m),c})_i = \max\{\lambda_i^{(m)} - c, 0\} \quad \text{for all } i \geq 1.$$

Now define

$$\begin{aligned} \lambda_R &= \lambda_R(c, m) = (\lambda^{(1)}, \dots, \lambda^{(m-1)}, \lambda_R^{(m),c}), \\ \lambda_L &= \lambda_L(c, m) = (\lambda_L^{(m),c}, \lambda^{(m+1)}, \dots, \lambda^{(l)}). \end{aligned}$$

Here is an enlightening pictorial representation of this construction, with $l = 3$, $m = 2$ and $c = 3$.



Now we consider tableaux. Suppose λ_L, λ_R are as above, and let $n_L = |\lambda_L|$ and $n_R = |\lambda_R|$. Given $T_L \in \text{Std}(\lambda_L)$ and $T_R \in \text{Std}(\lambda_R)$, define $T_L \# T_R$ to be the λ -tableau obtained by filling in the entries $1, \dots, n_L$ as they appear in T_L , and then filling in the entries $n_L + 1, \dots, n$ as $1, \dots, n_R$, respectively, appear in T_R . Observe that if $T \in \text{Std}(\lambda)$ and the integers $1, \dots, n_L$ all appear in T in column c of component m or further to the left, then T has the form $T_L \# T_R$ for some $T_L \in \text{Std}(\lambda_L)$ and $T_R \in \text{Std}(\lambda_R)$. We write $\text{Std}_{LR}(\lambda)$ for the set of $T \in \text{Std}(\lambda)$ with this property.

Example. Take $l = 3$ and $\lambda = ((3), (2^2), (2, 1))$. Taking $m = 2$ and $c = 1$, we get

$$\lambda_L = ((1^2), (2, 1)), \quad \lambda_R = ((3), (1^2)).$$

If we choose

$$T_L = \begin{array}{|c|c|} \hline 1 & \\ \hline 3 & \\ \hline \end{array}, \quad T_R = \begin{array}{|c|c|c|} \hline 2 & 3 & 5 \\ \hline 1 & & \\ \hline 4 & & \\ \hline \end{array}$$

then we obtain

$$T_L \# T_R = \begin{array}{|c|c|c|} \hline 7 & 8 & 10 \\ \hline 1 & 6 & \\ \hline 3 & 9 & \\ \hline \end{array}$$

4.2 Simple row and column removal

Theorem 4.1 (Graded Column Removal). *Suppose $\lambda, \mu \in \mathcal{P}_n^l$ and $1 \leq m \leq l$. Suppose that $\lambda^{(m+1)} = \dots = \lambda^{(l)} = \mu^{(m+1)} = \dots = \mu^{(l)} = \emptyset$, and $k := (\lambda^{(m)'})_1 = (\mu^{(m)'})_1$. Let $\lambda_R = \lambda_R(1, m)$,*

$\mu_R = \mu_R(1, m)$ and $\kappa_R = (\kappa_1, \dots, \kappa_{m-1}, \kappa_m + 1)$. Then

$$\mathrm{DHom}_{\mathcal{H}_n}(S_{\lambda|\kappa}, S_{\mu|\kappa}) \cong \mathrm{DHom}_{\mathcal{H}_{n-k}}(S_{\lambda_R|\kappa_R}, S_{\mu_R|\kappa_R})$$

as graded vector spaces over \mathbb{F} .

Remark. Recalling Theorem 3.6, this result in fact implies that $\mathrm{Hom}_{\mathcal{H}_n}(S_{\lambda|\kappa}, S_{\mu|\kappa}) \cong \mathrm{Hom}_{\mathcal{H}_{n-k}}(S_{\lambda_R|\kappa_R}, S_{\mu_R|\kappa_R})$ when $e \neq 2$ and $\kappa_1, \dots, \kappa_l$ are distinct.

Proof. We construct the isomorphism explicitly in the KLR setting. First note that we may assume $\lambda \supseteq \mu$, since otherwise $\mathrm{Std}_{\lambda_R}(\mu_R) = \mathrm{Std}_{\lambda}(\mu) = \emptyset$ and the result is immediate. We also observe that $\mathrm{cont}(\lambda) = \mathrm{cont}(\mu)$ if and only if $\mathrm{cont}(\lambda_R) = \mathrm{cont}(\mu_R)$; if these conditions do not hold then the result is trivial since both homomorphism spaces are zero, so we assume $\mathrm{cont}(\lambda) = \mathrm{cont}(\mu)$, and set $\alpha := \mathrm{cont}(\lambda)$, $\beta := \mathrm{cont}(\lambda_R)$.

For this proof we make an assumption about the choice of preferred reduced expressions defining the standard bases for $S_{\mu_R|\kappa_R}$ and $S_{\mu|\kappa}$. Given $T \in \mathrm{Std}_{\lambda_R}(\mu_R)$, we define $T^+ := T_{\mu_L} \# T$, where

$$\mu_L = \mu_L(1, m) = ((1^k), \emptyset, \dots, \emptyset) \in \mathcal{D}_k^{l-m+1}.$$

In other words, T^+ is obtained from T by increasing each entry by k , adding the column $\begin{bmatrix} 1 \\ \vdots \\ k \end{bmatrix}$ at

the left of component m , and then adding $l - m$ empty components at the end. Now recall the maps (both denoted shift_k) from \mathfrak{S}_{n-k} to \mathfrak{S}_n and from \mathcal{H}_β to \mathcal{H}_α . Observe that for $T \in \mathrm{Std}_{\lambda_R}(\mu_R)$ we have $w_{T^+} = \mathrm{shift}_k(w_T)$. By choosing compatible reduced expressions for w_{T^+} and w_T , we may assume that $\psi_{T^+} = \mathrm{shift}_k(\psi_T)$ as well.

Now let $c = (\lambda_R^{(m)})'_1$. Then the entries $1, \dots, c$ all appear in the first column of component m in T_{λ_R} , and hence if $T \in \mathrm{Std}_{\lambda_R}(\mu_R)$ these entries all appear in the first column of component m of T . In particular, w_T fixes $1, \dots, c$, so ψ_T only involves terms ψ_j for $j > c$; hence ψ_{T^+} only involves terms ψ_j for $j > k + c$.

Now suppose $\varphi_R \in \mathrm{DHom}_{\mathcal{H}_{n-k}}(S_{\lambda_R|\kappa_R}, S_{\mu_R|\kappa_R})$. Then

$$\varphi_R(z_{\lambda_R}) = \sum_{T \in \mathrm{Std}_{\lambda_R}(\mu_R)} a_T v_T \quad \text{for some } a_T \in \mathbb{F}.$$

We define $\varphi : S_{\lambda|\kappa} \rightarrow S_{\mu|\kappa}$ by

$$\varphi(z_\lambda) = \sum_{T \in \mathrm{Std}_{\lambda_R}(\mu_R)} a_T v_{T^+}.$$

We must verify that this does indeed define a homomorphism, i.e. that $h\varphi(z_\lambda) = 0$ whenever $h \in \mathrm{Ann}(z_\lambda)$. (Here and henceforth we write $\mathrm{Ann}(z_\lambda)$ for the annihilator of z_λ .) Firstly, note that if $T \in \mathrm{Std}_{\lambda_R}(\mu_R)$ with $a_T \neq 0$, then T has residue sequence i_{λ_R} ; this implies that T^+ has residue sequence i_λ , so that $e(i_\lambda)\varphi(z_\lambda) = \varphi(z_\lambda)$, as required. For the other relations, observe from the defining relations for the column Specht module that $\mathrm{shift}_k(\mathrm{Ann}(z_{\lambda_R})) \subseteq \mathrm{Ann}(z_\lambda)$ (and similarly for μ_R and μ). Now for $k < j \leq n$ we have $y_{j-k} \in \mathrm{Ann}(z_{\lambda_R})$, so (since φ_R is a homomorphism) $y_{j-k} \sum_T a_T \psi_T \in \mathrm{Ann}(z_{\mu_R})$. Hence

$$\mathrm{Ann}(z_\mu) \ni \mathrm{shift}_k \left(y_{j-k} \sum_T a_T \psi_T \right) = y_j \sum_T a_T \psi_{T^+},$$

so that $y_j\varphi(z_\lambda) = 0$. A similar statement applies to ψ_j whenever $k < j < n$ with $j \downarrow_{T_\lambda} j+1$, and to any Garnir element \mathbf{g}_A where A does not lie in the first column of component m .

It remains to check the generators of $\text{Ann}(z_\lambda)$ which do not lie in $\text{shift}_k(\text{Ann}(z_{\lambda_R}))$, i.e. the elements $y_1, \dots, y_k, \psi_1, \dots, \psi_{k-1}$ and \mathbf{g}_A for A of the form $(j, 1, m)$ with $1 \leq j \leq c$. Let h denote any of these elements, and observe that since each ψ_{T^+} is a product of terms ψ_i with $i > k+c$, h commutes with ψ_{T^+} (note that if $h = \mathbf{g}_{(j,1,m)}$, then h only involves terms ψ_i for $i < k+c$). Hence

$$h\varphi(z_\lambda) = h \sum_{\mathbf{T}} a_{\mathbf{T}} \psi_{\mathbf{T}^+} z_\mu = \sum_{\mathbf{T}} a_{\mathbf{T}} \psi_{\mathbf{T}^+} h z_\mu = 0,$$

since $h \in \text{Ann}(z_\mu)$.

So $\text{Ann}(z_\lambda)\varphi(z_\lambda) = 0$, and φ is a well-defined homomorphism. So we have a map $\Phi : \text{DHom}_{\mathcal{H}_{n-k}}(S_{\lambda_R|\kappa_R}, S_{\mu_R|\kappa_R}) \rightarrow \text{DHom}_{\mathcal{H}_n}(S_{\lambda|\kappa}, S_{\mu|\kappa})$ given by $\varphi_R \mapsto \varphi$, and Φ is obviously linear. To show that Φ is bijective, we construct its inverse. Any $S \in \text{Std}_\lambda(\mu)$ must have entries $1, \dots, k$ in order down the first column of its m th component; that is, $S = T^+$ for some $T \in \text{Std}_{\lambda_R}(\mu_R)$. So given $\theta \in \text{DHom}_{\mathcal{H}_n}(S_{\lambda|\kappa}, S_{\mu|\kappa})$, we can write

$$\theta(z_\lambda) = \sum_{T \in \text{Std}_{\lambda_R}(\mu_R)} a_T v_{T^+} \quad \text{for some } a_T \in \mathbb{F}.$$

Applying (a simpler version of) the above argument in reverse, we see that we have a homomorphism $\theta_R : S_{\lambda_R} \rightarrow S_{\mu_R}$ given by

$$\theta_R(z_{\lambda_R}) = \sum_{T \in \text{Std}_{\lambda_R}(\mu_R)} a_T v_T.$$

So we get a linear map $\text{DHom}_{\mathcal{H}_n}(S_{\lambda|\kappa}, S_{\mu|\kappa}) \rightarrow \text{DHom}_{\mathcal{H}_{n-k}}(S_{\lambda_R|\kappa_R}, S_{\mu_R|\kappa_R})$ which is a two-sided inverse to Φ , and hence Φ is a bijection.

Finally, to show that we have an isomorphism of graded vector spaces, we show that Φ is homogeneous of degree 0. That is, if $0 \neq \varphi_R \in \text{DHom}_{\mathcal{H}_{n-k}}(S_{\lambda_R|\kappa_R}, S_{\mu_R|\kappa_R})$ is homogeneous, then φ is also homogeneous with $\deg(\varphi) = \deg(\varphi_R)$. To see this, we write

$$\varphi_R(z_{\lambda_R}) = \sum_{T \in \text{Std}_{\lambda_R}(\mu_R)} a_T v_T \quad \text{for some } a_T \in \mathbb{F}.$$

Then

$$\varphi(z_\lambda) = \sum_{T \in \text{Std}_{\lambda_R}(\mu_R)} a_T v_{T^+},$$

and for each T with $a_T \neq 0$ we have

$$\text{codeg}^K(T^+) - \text{codeg}^K(T_\lambda) = \text{codeg}^{KR}(T) - \text{codeg}^{KR}(T_{\lambda_R}) = \deg(\varphi_R).$$

Hence φ is homogeneous of degree $\deg(\varphi_R)$. □

Now we make corresponding definitions for row removal.

Definition. Suppose $\lambda \in \mathcal{P}_n^l$. For any $1 \leq m \leq l$ and any $r \geq 0$, define

$$\lambda_T^{(m),r} = (\lambda_1^{(m)}, \dots, \lambda_r^{(m)}, 0, 0, \dots), \quad \lambda_B^{(m),r} = (\lambda_{r+1}^{(m)}, \lambda_{r+2}^{(m)}, \dots).$$

Now let

$$\begin{aligned} \lambda_T &= \lambda_T(r, m) = (\lambda^{(1)}, \dots, \lambda^{(m-1)}, \lambda_T^{(m),r}), \\ \lambda_B &= \lambda_B(r, m) = (\lambda_B^{(m),r}, \lambda^{(m+1)}, \dots, \lambda^{(l)}), \end{aligned}$$

and set $n_T = |\lambda_T|$ and $n_B = |\lambda_B|$.

Corollary 4.2 (Graded Row Removal). *Suppose $\lambda, \mu \in \mathcal{P}_n^l$ and $1 \leq m \leq l$. Suppose that $\lambda^{(1)} = \dots = \lambda^{(m-1)} = \mu^{(1)} = \dots = \mu^{(m-1)} = \emptyset$, and $k := \lambda_1^{(m)} = \mu_1^{(m)}$. Let $\lambda_B = \lambda_B(1, m)$, $\mu_B = \mu_B(1, m)$ and $\kappa_B = (\kappa_m - 1, \kappa_{m+1}, \dots, \kappa_l)$. Then*

$$\mathrm{DHom}_{\mathcal{H}_n}(S_{\lambda|\kappa}, S_{\mu|\kappa}) \cong \mathrm{DHom}_{\mathcal{H}_{n-k}}(S_{\lambda_B|\kappa_B}, S_{\mu_B|\kappa_B})$$

as graded vector spaces over \mathbb{F} .

Proof. $\mathrm{DHom}_{\mathcal{H}_n}(S_{\lambda|\kappa}, S_{\mu|\kappa}) \cong \mathrm{DHom}_{\mathcal{H}_n}(S_{\mu'|\kappa'}, S_{\lambda'|\kappa'})$ by Theorem 3.17,
 $\cong \mathrm{DHom}_{\mathcal{H}_{n-k}}(S_{(\mu_B)'|(\kappa_B)'}, S_{(\lambda_B)'|(\kappa_B)'})$ by Theorem 4.1,
 $\cong \mathrm{DHom}_{\mathcal{H}_{n-k}}(S_{\lambda_B|\kappa_B}, S_{\mu_B|\kappa_B})$ by Theorem 3.17 again. \square

Now we prove a ‘final-column removal’ theorem, where we assume that the rightmost non-empty columns of λ and μ are in the same place and of the same length.

Theorem 4.3 (Final Column Removal). *Suppose $\lambda, \mu \in \mathcal{P}_n^l$ and $1 \leq m \leq l$. Suppose $\lambda^{(1)} = \dots = \lambda^{(m-1)} = \mu^{(1)} = \dots = \mu^{(m-1)} = \emptyset$, $d := \lambda_1^{(m)} = \mu_1^{(m)}$ and $k := (\lambda^{(m)})'_d = (\mu^{(m)})'_d$. Let $\lambda_L = \lambda_L(d-1, m)$, $\mu_L = \mu_L(d-1, m)$ and $\kappa_L = (\kappa_m, \dots, \kappa_l)$. Then*

$$\mathrm{DHom}_{\mathcal{H}_n}(S_{\lambda|\kappa}, S_{\mu|\kappa}) \cong \mathrm{DHom}_{\mathcal{H}_{n-k}}(S_{\lambda_L|\kappa_L}, S_{\mu_L|\kappa_L})$$

as graded vector spaces over \mathbb{F} .

Proof. We first use Corollary 4.2 to remove the first k rows of length d from both $\lambda^{(m)}$ and $\mu^{(m)}$. We obtain

$$\mathrm{DHom}_{\mathcal{H}_n}(S_{\lambda|\kappa}, S_{\mu|\kappa}) \cong \mathrm{DHom}_{\mathcal{H}_{n-dk}}(S_{\lambda_B|\kappa_B}, S_{\mu_B|\kappa_B})$$

where $\lambda_B = \lambda_B(k, m)$, $\mu_B = \mu_B(k, m)$ and $\kappa_B = (\kappa_m - k, \kappa_{m+1}, \dots, \kappa_l)$. We then use Corollary 4.2 again to add k rows of length $d-1$ to the top of both $\lambda_B^{(m)}$ and $\mu_B^{(m)}$. We obtain

$$\mathrm{DHom}_{\mathcal{H}_{n-dk}}(S_{\lambda_B|\kappa_B}, S_{\mu_B|\kappa_B}) \cong \mathrm{DHom}_{\mathcal{H}_{n-k}}(S_{\lambda_L|\kappa_L}, S_{\mu_L|\kappa_L})$$

which gives the result. \square

It will be helpful below to be able to give a direct construction for final-column removal, as done in the proof of Theorem 4.1 for first-column removal. We assume the hypotheses and notation of Theorem 4.3, and for ease of notation we assume that S_λ and S_μ are defined

using the e -multicharge κ , while S_{λ_L} and S_{μ_L} are defined using κ_L . We can also assume that $\text{cont}(\lambda) = \text{cont}(\mu) =: \alpha$, and hence $\text{cont}(\lambda_L) = \text{cont}(\mu_L) =: \beta$.

We identify \mathfrak{S}_{n-k} with its image under the map $\text{shift}_0 : \mathfrak{S}_{n-k} \rightarrow \mathfrak{S}_n$, and similarly for \mathcal{H}_β and \mathcal{H}_α . As in the proof of Theorem 4.1 we make an assumption on preferred reduced expressions: given a standard μ_L -tableau T , we define T^+ to be the standard μ -tableau obtained by adding a column with entries $n-k+1, \dots, n$ at the right of component m ; then we have $w_{T^+} = w_T$, and we assume that our preferred reduced expressions have been chosen in such a way that $\psi_{T^+} = \psi_T$.

Lemma 4.4. *With the above notation, we have $\text{Ann}(z_{\lambda_L}) = \text{Ann}(z_\lambda) \cap \mathcal{H}_\beta$.*

Proof. It follows directly from the presentation for column Specht modules that $\text{Ann}(z_{\lambda_L}) \subseteq \text{Ann}(z_\lambda) \cap \mathcal{H}_\beta$, so we must show the opposite containment. Consider the \mathcal{H}_β -submodule $\mathcal{H}_\beta z_\lambda$ of S_λ generated by z_λ . For any $T \in \text{Std}(\lambda_L)$ we have $v_{T^+} = \psi_{T^+} z_\lambda = \psi_T z_\lambda \in \mathcal{H}_\beta z_\lambda$, and the v_{T^+} are linearly independent, so $\dim_{\mathbb{F}} \mathcal{H}_\beta z_\lambda \geq |\text{Std}(\lambda_L)| = \dim_{\mathbb{F}} S_{\lambda_L}$. So we have

$$\dim_{\mathbb{F}} \mathcal{H}_\beta z_\lambda \geq \dim_{\mathbb{F}} \mathcal{H}_\beta z_{\lambda_L},$$

i.e.

$$\dim_{\mathbb{F}} \frac{\mathcal{H}_\beta}{\text{Ann}(z_\lambda) \cap \mathcal{H}_\beta} \geq \dim_{\mathbb{F}} \frac{\mathcal{H}_\beta}{\text{Ann}(z_{\lambda_L})},$$

and so $\text{Ann}(z_{\lambda_L}) \supseteq \text{Ann}(z_\lambda) \cap \mathcal{H}_\beta$. □

Now we consider dominated homomorphisms. Observe that since λ and μ have the same last column, $\text{Std}_\lambda(\mu) = \{T^+ \mid T \in \text{Std}_{\lambda_L}(\mu_L)\}$. So if $\varphi \in \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$, then we can write

$$\varphi(z_\lambda) = \sum_{T \in \text{Std}_{\lambda_L}(\mu_L)} a_T v_{T^+} \quad \text{with } a_T \in \mathbb{F}.$$

Then we can define a homomorphism

$$\begin{aligned} \varphi^- : S_{\lambda_L|\kappa_L} &\longrightarrow S_{\mu_L|\kappa_L} \\ z_{\lambda_L} &\longmapsto \sum_{T \in \text{Std}_{\lambda_L}(\mu_L)} a_T v_T. \end{aligned}$$

To see that this definition yields a well-defined homomorphism, we must show that $h \sum_T a_T v_T = 0$ whenever $h \in \text{Ann}(z_{\lambda_L})$. By Lemma 4.4 we have $h \in \text{Ann}(z_\lambda)$, and hence (since φ is a homomorphism) $h \sum_T a_T v_{T^+} = 0$; in other words, $h \sum_T a_T \psi_T \in \text{Ann}(z_\mu)$. We also have $h \sum_T a_T \psi_T \in \mathcal{H}_\beta$, so by Lemma 4.4 again (with λ replaced by μ) $h \sum_T a_T \psi_T \in \text{Ann}(z_{\mu_L})$, as required.

So we have a map $\varphi \mapsto \varphi^- : \text{DHom}_{\mathcal{H}_n}(S_{\lambda|\kappa}, S_{\mu|\kappa}) \rightarrow \text{DHom}_{\mathcal{H}_{n-k}}(S_{\lambda_L|\kappa_L}, S_{\mu_L|\kappa_L})$. This is obviously an injective map of degree 0, and hence (by Theorem 4.3) a graded isomorphism.

4.3 Generalised column removal

Armed with first-column removal and final-column removal, we can now consider generalised column removal. In what follows, we fix $c \geq 0$ and $1 \leq m \leq l$, and for any $\nu \in \mathcal{P}_n^l$ we write $\nu_L = \nu_L(c, m)$ and $\nu_R = \nu_R(c, m)$. We suppose $\lambda, \mu \in \mathcal{P}_n^l$, and assume that $|\lambda_L| = |\mu_L| =: n_L$,

so that $|\lambda_R| = |\mu_R| = n - n_L =: n_R$. We also assume that $\lambda \triangleright \mu$. This assumption implies that $\lambda_L \triangleright \mu_L$ and $\lambda_R \triangleright \mu_R$, which in particular gives

$$(\lambda^{(m)})'_c \geq (\mu^{(m)})'_c \geq (\mu^{(m)})'_{c+1}$$

so that it is possible to define a multipartition $\lambda_L \# \mu_R \in \mathcal{P}_n^l$ with $(\lambda_L \# \mu_R)_L = \lambda_L$ and $(\lambda_L \# \mu_R)_R = \mu_R$.

We write $\kappa_L = (\kappa_m, \dots, \kappa_l)$, $\kappa_R = (\kappa_1, \dots, \kappa_m + c)$, $\mathcal{H}_L = \mathcal{H}_{n_L}$ and $\mathcal{H}_R = \mathcal{H}_{n_R}$. For ease of notation, we will assume throughout the following that the Specht modules S_λ , S_μ and $S_{\lambda_L \# \mu_R}$ are defined using the e -multicharge κ , while S_{λ_L} and S_{μ_L} are defined using κ_L and S_{λ_R} and S_{μ_R} are defined using κ_R .

Suppose $\varphi_L \in \text{DHom}_{\mathcal{H}_L}(S_{\lambda_L}, S_{\mu_L})$ and $\varphi_R \in \text{DHom}_{\mathcal{H}_R}(S_{\lambda_R}, S_{\mu_R})$, and write

$$\varphi_L(z_{\lambda_L}) = \sum_{S \in \text{Std}_{\lambda_L}(\mu_L)} a_S v_S, \quad \varphi_R(z_{\lambda_R}) = \sum_{T \in \text{Std}_{\lambda_R}(\mu_R)} b_T v_T$$

with coefficients $a_S, b_T \in \mathbb{F}$. If there is a homomorphism $\varphi : S_\lambda \rightarrow S_\mu$ satisfying

$$\varphi(z_\lambda) = \sum_{\substack{S \in \text{Std}_{\lambda_L}(\mu_L) \\ T \in \text{Std}_{\lambda_R}(\mu_R)}} a_S b_T v_{S \# T},$$

then we write $\varphi = \varphi_L \# \varphi_R$, and say that φ is a *product homomorphism*.

Lemma 4.5. *Every product homomorphism $S_\lambda \rightarrow S_\mu$ factors through $S_{\lambda_L \# \mu_R}$.*

Proof. Suppose that $\varphi = \varphi_L \# \varphi_R$ is a product homomorphism, and as above write

$$\varphi_L(z_{\lambda_L}) = \sum_{S \in \text{Std}_{\lambda_L}(\mu_L)} a_S v_S, \quad \varphi_R(z_{\lambda_R}) = \sum_{T \in \text{Std}_{\lambda_R}(\mu_R)} b_T v_T.$$

Now define

$$\begin{aligned} \varphi_L \# \text{id} : S_{\lambda_L \# \mu_R} &\longrightarrow S_\mu & \text{id} \# \varphi_R : S_\lambda &\longrightarrow S_{\lambda_L \# \mu_R} \\ z_{\lambda_L \# \mu_R} &\longmapsto \sum_{S \in \text{Std}_{\lambda_L}(\mu_L)} a_S v_{S \# T_{\mu_R}} & z_\lambda &\longmapsto \sum_{T \in \text{Std}_{\lambda_R}(\mu_R)} b_T v_{T_{\lambda_L} \# T}. \end{aligned}$$

Then $\varphi_L \# \text{id}$ and $\text{id} \# \varphi_R$ are both \mathcal{H}_n -homomorphisms; this follows from the direct constructions of column-removal homomorphisms in the proof of Theorem 4.1 and following the proof of Theorem 4.3. Clearly $(\varphi_L \# \text{id}) \circ (\text{id} \# \varphi_R) = \varphi$, so φ factors through $S_{\lambda_L \# \mu_R}$. \square

Proposition 4.6. *Assume the hypotheses and notation above. Then every $\varphi \in \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$ is a sum of product homomorphisms.*

Proof. We may assume that $\text{cont}(\lambda) = \text{cont}(\mu)$ (since otherwise there are no non-zero homomorphisms $S_\lambda \rightarrow S_\mu$). So for this proof we write $\alpha := \text{cont}(\lambda)$ and define shift_0 to be the map from \mathcal{H}_L to \mathcal{H}_α obtained by combining the maps $\text{shift}_0 : \mathcal{H}_\beta \rightarrow \mathcal{H}_\alpha$ for all $\beta \in Q^+$ of height n_L ; similarly, shift_{n_L} denotes the map from \mathcal{H}_R to \mathcal{H}_α obtained by combining the maps $\text{shift}_{n_L} : \mathcal{H}_\beta \rightarrow \mathcal{H}_\alpha$ for all $\beta \in Q^+$ of height n_R .

For this proof we make an assumption about the choice of preferred reduced expressions similar to that in the proof of Theorem 4.3. Specifically, we assume that these expressions have been chosen in such a way that if $S \in \text{Std}_{\lambda_L}(\mu_L)$ and $T \in \text{Std}_{\lambda_R}(\mu_R)$, then the preferred expression for $w_{S\#T}$ is just the concatenation of the preferred expression for w_S with the expression obtained by applying shift_{h_L} to every term in the preferred expression for w_T . Hence $\psi_{S\#T} = \psi_S \text{shift}_{h_L}(\psi_T)$.

Now we show that every dominated homomorphism $S_\lambda \rightarrow S_\mu$ is a sum of product homomorphisms. To do this, we first discuss dominated tableaux. Note that the conditions on λ and μ imply that $\text{Std}_\lambda(\mu) = \{T_L \# T_R \mid T_L \in \text{Std}_{\lambda_L}(\mu_L), T_R \in \text{Std}_{\lambda_R}(\mu_R)\}$. Choose a total order \blacktriangleright on $\text{Std}_\lambda(\mu)$ with the property that if $R, S \in \text{Std}_{\lambda_L}(\mu_L)$ and $T, U \in \text{Std}_{\lambda_R}(\mu_R)$, then

$$R\#T \blacktriangleright R\#U \iff S\#T \blacktriangleright S\#U \quad \text{and} \quad R\#T \blacktriangleright S\#T \iff R\#U \blacktriangleright S\#U.$$

(For example, we could do this by choosing total orders $\blacktriangleright_L, \blacktriangleright_R$ on $\text{Std}_{\lambda_L}(\mu_L), \text{Std}_{\lambda_R}(\mu_R)$ and setting $V \blacktriangleright W$ if and only if $V_L \blacktriangleright_L W_L$ or $(V_L = W_L \text{ and } V_R \blacktriangleright_R W_R)$.)

Now suppose $\varphi : S_\lambda \rightarrow S_\mu$ is a non-zero dominated homomorphism, and write $\varphi(z_\lambda) = \sum_{T \in \text{Std}_\lambda(\mu)} a_T v_T$ with each $a_T \in \mathbb{F}$. Let U be the largest tableau (with respect to \blacktriangleright) such that $a_U \neq 0$, and proceed by induction on U .

Claim. Let \mathcal{U} denote the set of tableaux $T \in \text{Std}_\lambda(\mu)$ such that $T_R = U_R$. Then there is an \mathcal{H}_L -homomorphism

$$\begin{aligned} \varphi_L^U : S_{\lambda_L} &\longrightarrow S_{\mu_L} \\ z_{\lambda_L} &\longmapsto \sum_{T \in \mathcal{U}} a_T v_{T_L}. \end{aligned}$$

Proof. First we make an observation, which follows from the construction of Specht modules and our assumptions on preferred reduced expressions. If $W \in \text{Std}_\lambda(\mu)$ and $h \in \mathcal{H}_L$, and we write $h v_{W_L} = \sum_{T \in \text{Std}(\mu_L)} b_T v_T$, then $\text{shift}_0(h) v_W = \sum_{T \in \text{Std}(\mu_L)} b_T v_{T\#W_R}$. In particular, $\text{shift}_0(h) v_W$ is a linear combination of basis elements v_S for $S \in \text{Std}_{LR}(\mu)$ with $S_R = W_R$.

Now take $h \in \text{Ann}(z_{\lambda_L})$. Then $\text{shift}_0(h) \in \text{Ann}(z_\lambda)$, so $\text{shift}_0(h) \sum_{T \in \text{Std}_\lambda(\mu)} a_T v_T = 0$ (because φ is a homomorphism). If we look just at $\text{shift}_0(h) \sum_{T \in \mathcal{U}} a_T v_T$, then by the previous paragraph this lies in $\langle v_T \mid T \in \text{Std}_{LR}(\mu), T_R = U_R \rangle_{\mathbb{F}}$, while $\text{shift}_0(h) \sum_{T \notin \mathcal{U}} a_T v_T$ lies in $\langle v_T \mid T \in \text{Std}_{LR}(\mu), T_R \neq U_R \rangle_{\mathbb{F}}$. The v_T are linearly independent, and hence

$$\langle v_T \mid T \in \text{Std}_{LR}(\mu), T_R = U_R \rangle_{\mathbb{F}} \cap \langle v_T \mid T \in \text{Std}_{LR}(\mu), T_R \neq U_R \rangle_{\mathbb{F}} = 0.$$

Hence $\text{shift}_0(h) \sum_{T \in \mathcal{U}} a_T v_T = 0$.

Define a linear map $\#U_R : S_{\mu_L} \rightarrow S_\mu$ by $v_T \mapsto v_{T\#U_R}$ for $T \in \text{Std}(\mu_L)$. Then, from above, we have

$$(hm)\#U_R = h(m\#U_R)$$

for any $h \in \mathcal{H}_L$ and any $m \in S_{\mu_L}$. So for each $h \in \text{Ann}(z_{\lambda_L})$, we have $h \sum_{T \in \mathcal{U}} a_T v_{T_L} = 0$.

We can do essentially the same thing left and right interchanged; that is, if we let $\mathcal{U}' = \{T \in \text{Std}_\lambda(\mu) \mid T_L = U_L\}$, then we have an \mathcal{H}_R -homomorphism

$$\begin{aligned} \varphi_R^U : S_{\lambda_R} &\longrightarrow S_{\mu_R} \\ z_{\lambda_R} &\longmapsto \sum_{T \in \mathcal{U}'} a_T v_{T_R}. \end{aligned}$$

As in the proof of Lemma 4.5, we construct homomorphisms

$$\varphi_L^U \# \text{id} : S_{\lambda_L \# \mu_R} \longrightarrow S_\mu \quad \text{id} \# \varphi_R^U : S_\lambda \longrightarrow S_{\lambda_L \# \mu_R},$$

whose composition is the product homomorphism $\varphi_L^U \# \varphi_R^U : S_\lambda \rightarrow S_\mu$. v_U appears with non-zero coefficient (namely a_U^2) in $\varphi_L^U \# \varphi_R^U$, and U is maximal (with respect to the order \blacktriangleright) with this property. So if we consider the homomorphism $\xi := \varphi - \frac{1}{a_U} \varphi_L \# \varphi_R$, then (if $\xi \neq 0$) the most dominant tableau occurring with non-zero coefficient in ξ is smaller than U . By induction ξ is a sum of product homomorphisms, and hence so is φ . \square

Now we can prove our main result.

Theorem 4.7 (Generalised graded column removal). *Suppose $\lambda, \mu \in \mathcal{P}_n^l$, $c \geq 0$ and $1 \leq m \leq l$ and define $\lambda_L, \lambda_R, \mu_L, \mu_R$ as in Section 4.1. Assume $|\lambda_L(c, m)| = |\mu_L(c, m)| =: n_L$ and $|\lambda_R(c, m)| = |\mu_R(c, m)| =: n_R$ for some fixed $c \geq 0$ and $1 \leq m \leq l$ and define $\mathcal{H}_L = \mathcal{H}_{n_L}$ and $\mathcal{H}_R = \mathcal{H}_{n_R}$.*

1. *For any $\varphi_L \in \text{DHom}_{\mathcal{H}_L}(S_{\lambda_L}, S_{\mu_L})$ and $\varphi_R \in \text{DHom}_{\mathcal{H}_R}(S_{\lambda_R}, S_{\mu_R})$, there is a product homomorphism $\varphi_L \# \varphi_R \in \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$.*
2. *The map $\varphi_L \otimes \varphi_R \mapsto \varphi_L \# \varphi_R$ defines an isomorphism of graded \mathbb{F} -vector spaces*

$$\text{DHom}_{\mathcal{H}_L}(S_{\lambda_L}, S_{\mu_L}) \otimes \text{DHom}_{\mathcal{H}_R}(S_{\lambda_R}, S_{\mu_R}) \cong \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu).$$

Proof. First suppose $\lambda \not\triangleright \mu$. Then $\text{Std}_\lambda(\mu) = \emptyset$, so $\text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu) = 0$. Furthermore, we have either $\lambda_L \not\triangleright \mu_L$ or $\lambda_R \not\triangleright \mu_R$, so that either $\text{DHom}_{\mathcal{H}_L}(S_{\lambda_L}, S_{\mu_L}) = 0$ or $\text{DHom}_{\mathcal{H}_R}(S_{\lambda_R}, S_{\mu_R}) = 0$. So the result follows.

So we can assume that $\lambda \triangleright \mu$, which allows us to define the multipartition $\lambda_L \# \mu_R$ as above. Applying Theorem 4.1 repeatedly, we have

$$\text{DHom}_{\mathcal{H}_n}(S_\lambda, S_{\lambda_L \# \mu_R}) \cong \text{DHom}_{\mathcal{H}_R}(S_{\lambda_R}, S_{\mu_R}).$$

Similarly, by Theorem 4.3 applied repeatedly we have

$$\text{DHom}_{\mathcal{H}_n}(S_{\lambda_L \# \mu_R}, S_\mu) \cong \text{DHom}_{\mathcal{H}_L}(S_{\lambda_L}, S_{\mu_L}).$$

Combining these isomorphisms, and using the explicit constructions given above, we have an isomorphism of graded vector spaces

$$\begin{aligned} \text{DHom}_{\mathcal{H}_L}(S_{\lambda_L}, S_{\mu_L}) \otimes \text{DHom}_{\mathcal{H}_R}(S_{\lambda_R}, S_{\mu_R}) &\xrightarrow{\sim} \text{DHom}_{\mathcal{H}_n}(S_{\lambda_L \# \mu_R}, S_\mu) \otimes \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_{\lambda_L \# \mu_R}) \\ \varphi_L \otimes \varphi_R &\mapsto (\varphi_L \# \text{id}) \otimes (\text{id} \# \varphi_R). \end{aligned}$$

Composition of homomorphisms yields a map

$$\omega : \text{DHom}_{\mathcal{H}_n}(S_{\lambda_L \# \mu_R}, S_\mu) \otimes \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_{\lambda_L \# \mu_R}) \longrightarrow \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$$

which is homogeneous of degree zero, and by Lemma 4.5 and Proposition 4.6 ω is surjective. So we have a surjective map

$$\begin{aligned} \text{DHom}_{\mathcal{H}_L}(S_{\lambda_L}, S_{\mu_L}) \otimes \text{DHom}_{\mathcal{H}_R}(S_{\lambda_R}, S_{\mu_R}) &\longrightarrow \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu) \\ \varphi_L \otimes \varphi_R &\mapsto \varphi_L \# \varphi_R. \end{aligned}$$

This map is easily seen to be injective, and the result follows. \square

Remark. Analogous results for graded decomposition numbers have recently been obtained by the second author and Bowman [BS], in the more general setting of the ‘diagrammatic Cherednik algebras’ of Webster.

4.4 Generalised row removal

Now we consider generalised row removal for homomorphisms between column Specht modules. Fix $1 \leq m \leq l$ and $r \geq 0$, and for any $\nu \in \mathcal{P}_n^l$ write $\nu_T = \nu_T(r, m)$, $\nu_B = \nu_B(r, m)$. Suppose $\lambda, \mu \in \mathcal{P}_n^l$ with $|\lambda_T| = |\mu_T| =: n_T$, so that $|\lambda_B| = |\mu_B| = n - n_T =: n_B$. Set $\kappa_T = (\kappa_1, \dots, \kappa_m)$ and $\kappa_B = (\kappa_m - r, \kappa_{m+1}, \dots, \kappa_l)$, and write $\mathcal{H}_T = \mathcal{H}_{n_T}$ and $\mathcal{H}_B = \mathcal{H}_{n_B}$. In what follows we shall take S_λ and S_μ to be defined with respect to the e -multicharge κ , S_{λ_T} and S_{μ_T} with respect to κ_T , and S_{λ_B} and S_{μ_B} with respect to κ_B .

With this notation in place, we can state a generalised row-removal theorem for homomorphisms. This follows from Theorem 4.7 using Theorem 3.17 in the same way that Corollary 4.2 is deduced from Theorem 4.1.

Theorem 4.8 (Generalised graded row removal). *Suppose $\lambda, \mu \in \mathcal{P}_n^l$, $r \geq 0$ and $1 \leq m \leq l$ and define $\lambda_T, \lambda_B, \mu_T, \mu_B, n_T, n_B, \mathcal{H}_T, \mathcal{H}_B$ as above. Then there is an isomorphism of graded \mathbb{F} -vector spaces*

$$\mathrm{DHom}_{\mathcal{H}_T}(S_{\lambda_T}, S_{\mu_T}) \otimes \mathrm{DHom}_{\mathcal{H}_B}(S_{\lambda_B}, S_{\mu_B}) \cong \mathrm{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu).$$

Our proof of Theorem 4.7 gives a direct construction of the column-removal isomorphism, but a direct construction for row removal seems to be hard to obtain, especially using the standard bases for column Specht modules.

Example. Take $e = 2$ and $\kappa = (0, 1, 0)$. Let $\lambda = ((1^2), (2, 1^3), (1))$ and $\mu = ((1), (3, 1), (3))$, and take $(m, r) = (2, 1)$, so that $\lambda_T = ((1^2), (2))$, $\lambda_B = ((1^3), (1))$ and $\mu_T = \mu_B = ((1), (3))$. Set $\kappa_T = (0, 1)$ and $\kappa_B = (0, 0)$. Then (regardless of the field \mathbb{F}) the graded dimensions of $\mathrm{DHom}_{\mathcal{H}_4}(S_{\lambda_T|\kappa_T}, S_{\mu_T|\kappa_T})$ and $\mathrm{DHom}_{\mathcal{H}_4}(S_{\lambda_B|\kappa_B}, S_{\mu_B|\kappa_B})$ are v and 1 respectively. So by Theorem 4.8 the graded dimension of $\mathrm{DHom}_{\mathcal{H}_8}(S_{\lambda|\kappa}, S_{\mu|\kappa})$ is v . The unique (up to scaling) homomorphisms

$$S_{\lambda_T} \longrightarrow S_{\mu_T}, \quad S_{\lambda_B} \longrightarrow S_{\mu_B}, \quad S_\lambda \longrightarrow S_\mu$$

are given by

$$z_{\lambda_T} \mapsto v_S, \quad z_{\lambda_B} \mapsto v_T, \quad z_\lambda \mapsto v_U + 2v_V,$$

where

$$S = \begin{array}{|c|c|c|} \hline & & 3 \\ \hline 1 & 2 & 4 \\ \hline \end{array}, \quad T = \begin{array}{|c|c|c|} \hline & & 2 \\ \hline 1 & 3 & 4 \\ \hline \end{array}, \quad U = \begin{array}{|c|c|c|} \hline & & 7 \\ \hline 2 & 6 & 8 \\ 3 & & \\ \hline \end{array}, \quad V = \begin{array}{|c|c|c|} \hline & & 7 \\ \hline 4 & 6 & 8 \\ 5 & & \\ \hline \end{array}$$

$$\begin{array}{|c|c|c|} \hline 1 & 4 & 5 \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline \end{array}$$

It seems hard to reconcile these homomorphisms when expressed in this form, except perhaps in characteristic 2. (Note that the incompatibility of these expressions is not an artefact of the choice of preferred reduced expressions – the standard basis elements appearing in this example are independent of the choice of reduced expressions.)

In order to obtain an explicit row-removal construction, it seems to be necessary to use a different basis for the Specht module. Suppose we have $\lambda_B, \lambda_T, \mu_B$ and μ_T as above, with

$|\mu_T| = n_T = |\lambda_T|$. Partition the set $\{1, \dots, n\}$ into two sets S_B and S_T , by defining S_B to be the set of integers in the bottom part of T_λ and S_T the set of integers in the top part; that is,

$$\begin{aligned} S_B &= \{T_\lambda(s, c, k) \mid (s, c, k) \in [\lambda] \text{ and either } k > m \text{ or } k = m \text{ and } s > r\}, \\ S_T &= \{T_\lambda(s, c, k) \mid (s, c, k) \in [\lambda] \text{ and either } k < m \text{ or } k = m \text{ and } s \leq r\}. \end{aligned}$$

Let $\text{lab}_B : \{1, \dots, n_B\} \rightarrow S_B$ and $\text{lab}_T : \{1, \dots, n_T\} \rightarrow S_T$ be the unique order-preserving bijections.

Now given a μ_B -tableau T and a μ_T -tableau S , define a μ -tableau $T\#S$ by composing lab_B with T and lab_T with S and ‘gluing’ in the natural way.

Lemma 4.9. *Suppose λ and μ satisfy the conditions above. If $T \in \text{Std}_{\lambda_B}(\mu_B)$ and $S \in \text{Std}_{\lambda_T}(\mu_T)$, then $T\#S \in \text{Std}_\lambda(\mu)$.*

Proof. First we show that $T\#S$ is standard. Suppose A and B are nodes in the same component of $[\mu]$, with B either immediately to the right of A or immediately below A ; then we require $T\#S(B) > T\#S(A)$. This is clear from the fact that S and T are standard and the functions lab_T and lab_B are order-preserving, except in the case where $A = (r, b, m)$ and $B = (r+1, b, m)$ for some $1 \leq b \leq \mu_{r+1}^{(m)}$. So assume we are in this situation.

Let $k = \lambda_{r+1}^{(m)}$. Then the first k columns of $\lambda_T^{(m)}$ all have length r . Since $\text{Std}_{\lambda_T}(\mu_T)$ is non-empty we have $\lambda_T \geq \mu_T$, and hence the first k columns of $\mu_T^{(m)}$ all have length r also. Hence (since S is λ_T -dominated) S agrees with T_{λ_T} on these columns. So we have $T\#S(A) = \text{lab}_T(T_{\lambda_T}(A)) = T_\lambda(A)$.

We also have $\lambda_B \geq \mu_B$ since $\text{Std}_{\lambda_B}(\mu_B) \neq \emptyset$, so that $k \geq \mu_{r+1}^{(m)} \geq b$ (and in particular $B \in [\lambda]$). Since T is λ_B -dominated, we have $T(1, b, 1) \geq T_{\lambda_B}(1, b, 1)$, so that

$$T\#S(B) = \text{lab}_B(T(1, b, 1)) \geq \text{lab}_B(T_{\lambda_B}(1, b, 1)) = T_\lambda(B).$$

So $T\#S(A) = T_\lambda(A) < T_\lambda(B) \leq T\#S(B)$, as required.

To see that $T\#S$ is λ -dominated, it suffices to note that since $S \in \text{Std}_{\lambda_T}(\mu_T)$, every element of S_T appears in $\text{lab}_T(S)$ at least as far to the left as it appears in T_λ , and likewise for $T \in \text{Std}_{\lambda_B}(\mu_B)$ and elements of S_B . \square

Now we can give a conjectured explicit construction for the generalised row-removal isomorphism for homomorphisms. Recall from Section 3.3 the basis $\{f_T \mid T \in \text{Std}(\mu)\}$ for $(S^\mu)^\otimes$; using Theorem 3.14 and shifting the degree of each f_T by $\text{def}(\mu)$, we can regard $\{f_T \mid T \in \text{Std}(\mu)\}$ as a basis for S_μ . Note that by the analogue of Lemma 3.18(2) for column Specht modules, any $\varphi \in \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$ can be written as

$$\varphi(z_\lambda) = \sum_{T \in \text{Std}_\lambda(\mu)} a_T f_T \quad \text{for some } a_T \in \mathbb{F}.$$

Conjecture 4.10. *Suppose $\lambda, \mu \in \mathcal{P}_n^l$, $r \geq 0$ and $1 \leq m \leq n$. Define $\lambda_T, \lambda_B, \mu_T, \mu_B, n_T, n_B, \mathcal{H}_T, \mathcal{H}_B$ as above, and assume $|\mu_T| = n_T$. Suppose $\varphi_T \in \text{DHom}_{\mathcal{H}_T}(S_{\lambda_T}, S_{\mu_T})$ and $\varphi_B \in \text{DHom}_{\mathcal{H}_B}(S_{\lambda_B}, S_{\mu_B})$, and write*

$$\varphi_B(z_{\lambda_B}) = \sum_{T \in \text{Std}_{\lambda_B}(\mu_B)} a_T f_T, \quad \varphi_T(z_{\lambda_T}) = \sum_{S \in \text{Std}_{\lambda_T}(\mu_T)} b_S f_S$$

with $a_T, b_S \in \mathbb{F}$. Then there is an \mathcal{H}_n -homomorphism $\varphi_B \bar{\#} \varphi_T : S_\lambda \rightarrow S_\mu$ satisfying

$$\varphi_B \bar{\#} \varphi_T(z_\lambda) = \sum_{\substack{T \in \text{Std}_{\lambda_B}(\mu_B) \\ S \in \text{Std}_{\lambda_T}(\mu_T)}} a_T b_S f_{T \bar{\#} S}.$$

Example. Retaining the notation from the last example, we have

$$T_\lambda = \begin{array}{c} \boxed{7} \\ \boxed{8} \\ \\ \boxed{2} \boxed{6} \\ \boxed{3} \\ \boxed{4} \\ \boxed{5} \\ \boxed{1} \end{array}$$

so that $S_T = \{2, 6, 7, 8\}$ and $S_B = \{1, 3, 4, 5\}$. Taking S , T and U as in the last example, we get $T \bar{\#} S = U$. It is easy to check that

$$f_S = v_S, \quad f_T = v_T, \quad f_U = v_U + 2v_V,$$

so the conjecture holds in this case.

Remark. If Conjecture 4.10 is true, then we have a map of graded \mathbb{F} -vector spaces

$$\begin{aligned} \text{DHom}_{\mathcal{H}_B}(S_{\lambda_B}, S_{\mu_B}) \otimes \text{DHom}_{\mathcal{H}_T}(S_{\lambda_T}, S_{\mu_T}) &\longrightarrow \text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu) \\ \varphi_B \otimes \varphi_T &\longmapsto \varphi_B \bar{\#} \varphi_T. \end{aligned}$$

This map is obviously linear, and (since the f_T are linearly independent) injective. Hence by Theorem 4.8 it is a bijection. So we have an explicit construction for the generalised row-removal isomorphism.

5 Index of notation

For the reader's convenience we conclude with an index of the notation we use in this paper. We provide references to the relevant subsections.

\mathbb{F}	a field	
\mathbb{N}	the set of positive integers	
\mathfrak{S}_n	the symmetric group of degree n	2.1
s_1, \dots, s_{n-1}	the Coxeter generators of \mathfrak{S}_n	2.1
l	the Coxeter length function on \mathfrak{S}_n	2.1
\leq_L	the left order on \mathfrak{S}_n	2.1
\preceq	the Bruhat order on \mathfrak{S}_n	2.1
shift_k	the shift homomorphism $\mathfrak{S}_m \rightarrow \mathfrak{S}_n$	2.1
I	the set $\mathbb{Z}/e\mathbb{Z}$ (or \mathbb{Z} , if $e = \infty$)	2.2
Γ	a quiver with vertex set I	2.2
$i \rightarrow j$	there is an arrow from i to j (but no arrow from j to i) in Γ	2.2

$i \rightrightarrows j$	there are arrows from i to j and from j to i in Γ	2.2
α_i	simple root labelled by $i \in I$	2.2
Λ_i	fundamental dominant weight labelled by $i \in I$	2.2
(\mid)	invariant bilinear form	2.2
Q^+	the positive root lattice	2.2
Λ_κ	the dominant weight $\Lambda_{\kappa_1} + \cdots + \Lambda_{\kappa_l}$	2.2
$\text{def}(\alpha)$	$(\Lambda_k \mid \alpha) - \frac{1}{2}(\alpha \mid \alpha)$	2.2
$ \lambda $	the number of nodes of a (multi)partition λ	2.3
\mathcal{P}_n^l	the set of l -multipartitions of n	2.3
\emptyset	the unique partition or l -multipartition of 0	2.3
\supseteq	the dominance order on multipartitions or tableaux	2.3
$[\lambda]$	the Young diagram of a multipartition λ	2.3
λ'	the conjugate (multi)partition to λ	2.3
$\text{Std}(\lambda)$	the set of standard λ -tableaux	2.4
T'	the conjugate tableau to T	2.4
$i \downarrow_T j$	i and j lie in the same column of T , with j lower than i	2.4
$i \swarrow_T j$	i and j lie in the same component of T , with j strictly lower and to the left of i	2.4
$i \not\swarrow_T j$	$i \swarrow_T j$ or i lies in an earlier component of T than j	2.4
T_λ	the λ -tableau obtained by writing $1, \dots, n$ in order down successive columns	2.4
T^λ	the λ -tableau obtained by writing $1, \dots, n$ in order along successive rows	2.4
w_T	the permutation for which $w_T T_\lambda = T$	2.4
w^T	the permutation for which $w^T T^\lambda = T$	2.4
$\text{Shape}(T_{\downarrow m})$	the l -multicomposition formed from the nodes of T whose entries are less than or equal to m	2.4
$\text{res } A$	the residue of a node A	2.5
$\text{cont}(\lambda)$	the content of a multipartition λ	2.5
$\text{def}(\lambda)$	the defect of a multipartition λ	2.5
$i(T)$	the residue sequence of a tableau T	2.5
i_λ	$i(T_\lambda)$	2.5
i^λ	$i(T^\lambda)$	2.5
$\deg(T)$	the degree of a tableau T	2.5
$\text{codeg}(T)$	the codegree of a tableau T	2.5
\mathcal{H}_n	the KLR algebra of degree n	2.6
shift_κ	the shift homomorphism $\mathcal{H}_\beta \rightarrow \mathcal{H}_\alpha$	2.6
\mathcal{H}_n^κ	the cyclotomic KLR algebra determined by κ	2.6
\mathbf{B}_A	the Garnir belt corresponding to a Garnir node A	2.7
\mathbf{g}_A	the Garnir element corresponding to a Garnir node A	2.7
S_λ	the column Specht module corresponding to a multipartition λ	2.7
S^λ	the row Specht module corresponding to a multipartition λ	2.7
z_λ	the standard generator of S_λ	2.7
ψ_T	$\psi_{t_1} \dots \psi_{t_b}$, where $s_{t_1} \dots s_{t_b}$ is the preferred reduced expression for w_T	2.7
v_T	$\psi_T z_\lambda$	2.7
$\text{Std}_\lambda(\mu)$	the set of λ -dominated standard μ -tableaux	3.1

$\text{Std}^\lambda(\mu)$	the set of λ -row-dominated standard μ -tableaux	3.1
$\text{DHom}_{\mathcal{H}_n}(S_\lambda, S_\mu)$	the space of dominated homomorphisms from S_λ to S_μ	3.2
$\text{DHom}_{\mathcal{H}_n}(S^\lambda, S^\mu)$	the space of dominated homomorphisms from S^λ to S^μ	3.2
M^\circledast	the graded dual of a graded module M	3.3
$M\langle k \rangle$	the graded module M with the grading shifted by k	3.3
$\text{Std}_{\text{LR}}(\lambda)$	the set of λ -tableaux in which the entries $1, \dots, n_L$ appear strictly to the left of the entries $n_L + 1, \dots, n$	4
$\lambda_L \# \lambda_R$	the multipartition obtained by joining the left and right parts λ_L, λ_R together	4.3
$T_L \# T_R$	the tableau obtained by joining the left and right parts T_L, T_R together	4.3

References

- [BS] C. Bowman & L. Speyer, ‘An analogue of row removal for diagrammatic Cherednik algebras’, arXiv:1601.05543. 30
- [BK] J. Brundan & A. Kleshchev, ‘Blocks of cyclotomic Hecke algebras and Khovanov–Lauda algebras’, *Invent. Math.* **178** (2009), 451–484. cited on pp. 1, 3, 9
- [BKW] J. Brundan, A. Kleshchev & W. Wang, ‘Graded Specht modules’, *J. Reine Angew. Math.* **655** (2011), 61–87. 1, 3, 5, 7, 12
- [D] V. Deodhar, ‘Some characterizations of Bruhat ordering on a Coxeter group and determination of the relative Möbius function’, *Invent. Math.* **39** (1977), 187–198. 3
- [FL] M. Fayers & S. Lyle, ‘Row and column removal theorems for homomorphisms between Specht modules’, *J. Pure Appl. Algebra* **185** (2003), 147–164. 2, 13
- [HM] J. Hu & A. Mathas, ‘Graded cellular bases for the cyclotomic Khovanov–Lauda–Rouquier algebras of type A' ’, *Adv. Math.* **225** (2010), 598–642. 21
- [H] J. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Studies in Advanced Mathematics **29**, Cambridge University Press, 1990. 2, 3
- [J] G. James, *The representation theory of the symmetric groups*, Lecture notes in mathematics **682**, Springer-Verlag, New York/Berlin, 1978. 19
- [K] V. Kac, *Infinite dimensional Lie algebras*, 3rd Edition, Cambridge University Press 1990. 3
- [KL] M. Khovanov & A. Lauda, ‘A diagrammatic approach to categorification of quantum groups I’, *Represent. Theory* **13** (2009), 309–347. 1, 9
- [KMR] A. Kleshchev, A. Mathas & A. Ram, ‘Universal graded Specht modules for cyclotomic Hecke algebras’, *Proc. London Math. Soc.* **105** (2012), 1245–1289. 1, 2, 3, 5, 10, 11, 19, 20, 21
- [LM] S. Lyle & A. Mathas, ‘Row and column removal theorems for homomorphisms of Specht modules and Weyl modules’, *J. Algebraic Combin.* **22** (2005), 151–179. 13
- [M] A. Mathas, *Iwahori–Hecke Algebras and Schur Algebras of the Symmetric Group*, University Lecture Series **15**, American Mathematical Society, Providence, RI, 1999. 6
- [R1] R. Rouquier, ‘ q -Schur algebras and complex reflection groups’, *Mosc. Math. J.* **8** (2008), 119–158. 18
- [R2] R. Rouquier, ‘2-Kac–Moody algebras’, arXiv:0812.5023. 1, 9
- [S] L. Speyer, ‘Decomposable Specht modules for the Iwahori–Hecke algebra $\mathcal{H}_{\mathbb{R}, -1}(\mathfrak{S}_n)$ ’, *J. Algebra* **418** (2014), 227–264. 19