

Laplacian eigenvalues and optimality: II. The Laplacian of a graph

R. A. Bailey and Peter Cameron



London Taught Course Centre, June 2012




The Laplacian of a graph

This lecture will be about the Laplacian matrix of a graph and its eigenvalues, and their relation to some graph parameters. This is not a complete account of the theory, but concentrates mainly on the things that are most relevant for experimental design.

For further reading, we recommend

- ▶ B. Bollobás, *Modern Graph Theory*, Springer, (especially chapters II and IX)
- ▶ B. Mohar, Some applications of Laplace eigenvalues of graphs, pp.227–275 in *Graph Symmetry: Algebraic Methods and Applications* (ed. G. Hahn and G. Sabidussi), Kluwer, 1997.

Which graph is best?



Of course the question is not well defined. But which would you choose for a network, if you were concerned about connectivity, reliability, etc.?

What makes a good network?

- ▶ No two vertices should be too far apart.
- ▶ There should be several alternative routes between two vertices. (But should these routes be disjoint?)
- ▶ There should be no “bottlenecks”.
- ▶ Loss of a small part of the network should not result in disconnection.

Of course, we are resource-limited, else we would just put multiple edges between any two nodes.

Which graph is best connected?

Here are some ways of measuring the “connectivity” of a graph.

- ▶ How many spanning trees does it have? The more spanning trees, the better connected. The first graph has 2000 spanning trees, the second has 576.
- ▶ Electrical resistance. Imagine that the graph is an electrical network with each edge being a 1-ohm resistor. Now calculate the resistance between each pair of terminals, and sum over all pairs; the lower the total, the better connected. In the first graph, the sum is 33; in the second, it is 206/3.

Which graph is best connected?

- ▶ Isoperimetric number. This is defined to be

$$i(G) = \min \left\{ \frac{|\partial S|}{|S|} : S \subseteq V(G), 0 < |S| \leq n/2 \right\},$$

where $n = |V(G)|$ and for a set S of vertices, ∂S is the set of edges from S to its complement. Large isoperimetric number means that there are many edges out of any set of vertices. The isoperimetric number for the first graph is 1 (there are just five edges between the inner and outer pentagons), that of the second graph is 1/5 (there is just one edge between the top and bottom pieces).

The Laplacian of a graph

Let G be a graph on n vertices. (Multiple edges are allowed but loops are not.)

The **Laplacian matrix** of G is the $n \times n$ matrix $L = L(G)$ whose (i, i) entry is the number of edges containing vertex i , while for $i \neq j$ the (i, j) entry is the negative of the number of edges joining vertices i and j .

This is a real symmetric matrix; its eigenvalues are the **Laplacian eigenvalues** of G . Note that its row sums are zero.

The weighted Laplacian

Suppose that we have positive weights $w(e)$ on the edges of G . Then the **weighted Laplacian** has the (i, i) entry the sum of weights of edges containing i , and whose (i, j) entry for $i \neq j$ is minus the sum of the weights of edges joining i to j . If the weights are rational, then we may multiply them by the least common multiple of the denominators to make them integers. Then replace an edge of weight w by w edges, to obtain a multigraph with the same Laplacian. For general real weights, replace them first by rational approximations. We will not consider weighted Laplacians.

Relation to classical Laplacian

The classical Laplacian is a second-order differential operator defined on functions on a manifold, closely related to potential theory, the wave equation, etc. A manifold can be **triangulated**, that is, approximated by a graph drawn in it. If we take the weight of an edge to be inversely proportional the square of its length, then the weighted Laplacian of the graph is an approximation to the Laplacian of the manifold. In the other direction, given a graph, we can build a manifold reflecting its structure. Given a d -valent vertex, take a sphere with d holes; glue spheres corresponding to the vertices of an edge together along the corresponding holes. We won't pursue this any further.

Adjacency matrix and Laplacian

The usual **adjacency matrix** $A(G)$ of a graph G on n vertices has rows and columns indexed by vertices; the (i, j) entry is the number of edges connecting i to j . Note that we can allow loops here (though it is not clear whether a loop contributes 1 or 2 to the corresponding diagonal entry!) For the Laplacian, we forbid loops. If G is a regular graph with valency d , then $L(G) = dI - A(G)$; so the Laplacian eigenvalues are obtained by subtracting the adjacency matrix eigenvalues from d . If G is not regular, there is no such simple relationship between the eigenvalues of the two matrices.

Positive semi-definiteness

Theorem

The Laplacian of a graph is positive semidefinite.

For L is the sum of submatrices $\begin{pmatrix} +1 & -1 \\ -1 & +1 \end{pmatrix}$, one for each edge (this 2×2 matrix in the positions indexed by the two vertices of the edge, with zeros elsewhere). This matrix is positive semidefinite (its eigenvalues are 2 and 0.) We'll see another argument for this later. It follows that the eigenvalues of L are all non-negative.

The multiplicity of zero

Theorem

The multiplicity of 0 as an eigenvalue of L is equal to the number of connected components of the graph.

An eigenvector with zero eigenvalue is a function on the vertices whose value at i is the weighted average of its values on the neighbours of i , each neighbour weighted by the number of edges joining it to i . (If you know about harmonic functions, you will recognise this!) Considering a vertex where the maximum modulus is achieved, we see that the same value occurs on all neighbours, so the function is constant on connected components. In particular, if the graph is connected (as we always assume), the zero eigenvalue (called "trivial") has multiplicity 1; the other eigenvalues are **nontrivial**. The eigenvectors for the trivial eigenvalue are the constant vectors.

On average

Note that the sum of the eigenvalues is the trace of L , which is the sum of the vertex valencies, or twice the number of edges. So the average of the non-trivial eigenvalues is $2|E(G)| / (|V(G)| - 1)$; it depends just on the numbers of vertices and edges, and the detailed structure of the graph has no effect. We'll see that other means, in particular the geometric and harmonic means, of the non-trivial eigenvalues, give us important information!

Examples

The Petersen graph is strongly regular; its adjacency matrix A satisfies $A^2 + A - 2I = J$, where J is the all-1 matrix; its eigenvalues are 3, 1 and -2 , and so the Laplacian eigenvalues are 0, 2 and 5, with multiplicities 1, 5 and 4 respectively. For the other graph in our introductory example, the Laplacian eigenvalues are 0, 2, 3 (multiplicity 2), 4 (multiplicity 2), 5, and the roots of $x^3 - 9x^2 + 20x - 4$ (which are approximately 0.2215, 3.2892, and 5.4893).

The Rayleigh principle

Recall that eigenvectors corresponding to distinct eigenvalues of a symmetric matrix are orthogonal.

Theorem

Let λ_1, λ_2 be the smallest and second-smallest eigenvalues of the symmetric matrix A , and suppose that λ_1 is a simple eigenvalue with eigenvector u . Let v be any non-zero vector orthogonal to u . Then

$$\frac{vAv^T}{vv^T} \geq \lambda_2,$$

with equality if and only if v is an eigenvector with eigenvalue λ_2 .

This is obvious when v is expressed as a linear combination of eigenvectors of A .

There is an extension to any eigenvalue.

The cutset lemma

Theorem

Let G be a connected graph on n vertices, and E a set of m edges whose removal disconnects G into vertex sets of sizes n_1 and n_2 , with $n_1 + n_2 = n$. Let μ be the smallest non-trivial eigenvalue of L . Then $m \geq \mu n_1 n_2 / (n_1 + n_2)$.

For let V_1 and V_2 be the vertex sets in the theorem, and let v be the vector with value n_2 on vertices in V_1 , and $-n_1$ in vertices in V_2 . These values are chosen so that v is orthogonal to the all-1 vector (the trivial eigenvector). Clearly,

$$vv^T = n_1 n_2^2 + n_2 n_1^2 = (n_1 + n_2) n_1 n_2.$$

I claim that $v^T L v = m(n_1 + n_2)^2$. Recall that L is the sum of submatrices corresponding to edges; we have to add the contributions of these. An edge within one of the parts contributes 0; one between the parts contributes $(n_1 + n_2)^2$. The claim follows.

The theorem follows from the Rayleigh principle.

Isoperimetric number

Theorem

Let G be a connected graph whose smallest nontrivial Laplacian eigenvalue is μ . Then the isoperimetric number $i(G)$ is at least $\mu/2$.

For let S be a set of at most half the vertices, and let $|S| = n_1$, $|V \setminus S| = n_2$, and $|\partial(S)| = m$. Then by the cutset lemma,

$$\frac{|\partial(S)|}{|S|} = \frac{m}{n_1} \geq \frac{\mu n_2}{n_1 + n_2} \geq \frac{\mu}{2}.$$

So, on one of our criteria, a good network is one whose smallest nontrivial Laplacian eigenvalue is as large as possible.

Examples

In our two examples, the smallest nontrivial Laplacian eigenvalues are 2 (for the Petersen graph) and 0.2215 (for the other graph).

Note that the Petersen graph has isoperimetric number 1, meeting the bound of half the least non-trivial eigenvalue. So the vector which is $+1$ on the outer pentagon and -1 on the inner pentagon is an eigenvector.

In the other graph, the true value is a bit more than half the smallest eigenvalue.

Expanders

Loosely, an expander is a regular connected graph whose smallest non-trivial eigenvalue is large. The above result shows that expanders have large isoperimetric numbers.

More precisely, a sequence $(G_n : n \in \mathbb{N})$ is a sequence of **expanders** if there is a constant $c > 0$ such that the smallest non-trivial Laplacian eigenvalue of every graph G_n is at least c .

It is known that a random regular graph is an expander with high probability; but explicit constructions are more difficult.

The first constructions were given by Margulis and by Lubotzky, Phillips and Sarnak, and depend on substantial number-theoretic and group-theoretic background.

Cheeger's inequality

Cheeger's inequality is a result about Laplacians of manifolds; it has a discrete analogue. It gives a bound in the other direction between the isoperimetric number and the smallest nontrivial Laplacian eigenvalue.

Theorem


Let G be a connected graph; let Δ be the maximum valency of G , and μ the smallest nontrivial Laplacian eigenvalue. Then

$$i(G) \leq \sqrt{2\Delta\mu}.$$

Mohar improved the upper bound to $\sqrt{(2\Delta - \mu)\mu}$ if the graph is connected but not complete.

Incidence matrix

Choose a fixed but arbitrary orientation of the edges of the graph G . Define the vertex-edge **incidence matrix** Q to have rows indexed by vertices, columns by edges, and (v, e) entry $+1$ if v is the head of the edge e , -1 if v is the tail of e , and 0 otherwise.



Incidence matrix and Laplacian

Theorem

Let G have incidence matrix Q and Laplacian L . Then $QQ^T = L$.

For the (v, v) entry of QQ^T is the number of edges with either head or tail at v ; and the (v, w) entry is the sum of -1 for all edges with head at v and tail at w or *vice versa*.

This shows, again, that L is positive semidefinite. And note that the orientation doesn't matter.

The Moore–Penrose inverse

Let A be a real symmetric matrix. Then we have a **spectral decomposition** of A :

$$A = \sum_{\lambda \in \Lambda} \lambda P_\lambda,$$

where Λ is the set of eigenvalues of A , and P_λ is the orthogonal projector onto the space of eigenvectors with eigenvalue λ .

We define the **Moore–Penrose inverse** of A by

$$A^- = \sum_{\lambda \neq 0} \lambda^{-1} P_\lambda.$$

In other words, we invert A where we can.

The Moore–Penrose inverse is a **quasi-inverse** of A in the sense of ring theory: that is,

$$A^- A A^- = A^-, \quad A A^- A = A.$$

The Moore–Penrose inverse of the Laplacian

We will see a lot of the matrix L^- , where L is the Laplacian of a graph G on n vertices.

If G is connected, then the projector onto the trivial eigenspace is J/n , where J is the all-1 matrix. So adding this to L changes the trivial eigenvalue to 1, and subtracting it takes the 1 off again.

In other words,

$$L^- = (L + J/n)^{-1} - J/n.$$

Electrical networks

As mentioned earlier, we regard the graph G as an electrical network, where we regard each edge as a one-ohm resistor. Given any two vertices i and j , the **effective resistance** between i and j is the voltage of a battery which, when connected to the two vertices, causes a current of 1 ampere to flow.

Theorem

Let G be a connected graph with Laplacian L . Then the effective resistance between i and j is

$$L_{ii}^- + L_{jj}^- - L_{ij}^- - L_{ji}^-,$$

where L^- is the Moore–Penrose inverse of L .

Ohm's Law and Kirchhoff's Laws

If we apply a voltage between two vertices, the flow of current in edges and the potentials at vertices are governed by three laws:

- ▶ **Ohm's Law:** the potential drop in each edge is the product of the current and the resistance (and so is equal to the current since we have set all resistances to 1).
- ▶ **Kirchhoff's Voltage Law:** the sum of the potential drops on any path between vertices i and j is independent of the choice of path.
- ▶ **Kirchhoff's Current Law:** if vertex i is not connected to the battery, then the sum of the currents flowing into i is equal to the sum of the currents flowing out.

Proof of the theorem

Kirchhoff's voltage law and Ohm's Law are taken care of if we take a vector p of potentials with components indexed by vertices, and require that the current on the edge e is equal to the potential difference between its ends. (As before, we take a fixed ordering of each edge, and take the current to be negative if it flows from head to tail of the edge.) Note that p is defined up to adding a constant vector. This is expressed by the requirement that pQ is the vector of currents in the edges, where Q is the vertex-edge incidence matrix.

$pQQ^T = pL$ is a vector whose i th entry is the sum of the signed currents into the vertex i . So Kirchhoff's current law says that pQQ^T has all entries zero except at the two vertices connected to the battery. If the current is 1 ampere, the entries in $pL = pQQ^T$ are +1 and -1 on these two vertices. Let us write $pL = f_i - f_j$, where f_i is the unit basis vector corresponding to v . Now $f_i - f_j$ is orthogonal to the all-1 vector, so $(f_i - f_j)L^- = p$. This gives the vector of potentials. The potential difference between v and w is the dot product of this vector with $f_i - f_j$, that is, xL^-x^T , where $x = f_i - f_j$. This is the potential difference required to make a current of 1 ampere flow; hence it is the effective resistance between i and j . This can be written

$$R_{ij} = L_{ii}^- + L_{jj}^- - L_{ij}^- - L_{ji}^-,$$

as required.

The average pairwise resistance

One of our criteria for a good network is that the average pairwise resistance between two vertices should be small. The next theorem shows that this is equivalent to maximizing the **harmonic mean** of the nontrivial Laplacian eigenvalues.

Theorem

The average pairwise resistance is equal to 2 divided by the harmonic mean of the nontrivial Laplacian eigenvalues.

Proof of the resistance theorem

The sum of the resistances between all ordered pairs of vertices is

$$\sum_{i \neq j} R_{ij} = 2(n-1) \text{Trace}(L^-) - 2 \sum_{i \neq j} L_{ij}^- = 2n \text{Trace}(L^-),$$

since the sum of all elements of L^- is zero (as the all-1 vector is an eigenvector with eigenvalue 0).

So the average pairwise resistance is $2 \text{Trace}(L^-) / (n-1)$.

Now the trace of L^- is the sum of the reciprocals of the non-zero eigenvalues of L , and so we are done.

Examples

For the Petersen graph, the harmonic mean of the non-trivial eigenvalues is

$$(((5 \cdot 1/2) + (4 \cdot 1/5))/9)^{-1} = 30/11,$$

so the average resistance is 11/15.

For the other graph, a similar calculation gives 135/103, so the average resistance is 206/135.

Example: Petersen graph

For the Petersen graph, we can exploit symmetry to calculate the resistance between two terminals. Two vertices equivalent under a symmetry fixing the terminals must be at the same potential, and so edges between them can be neglected. If the terminals i and i are joined, the graph reduces to a pentagon $i = i_0, i_1, i_2, i_3, i_4 = j$, with one edge from i to j , two from i_0 to i_1 and from i_3 to i_4 , and four from i_1 to i_2 and i_2 to i_3 . So the resistance of the path $(i_0, i_1, i_2, i_3, i_4)$ is $1/2 + 1/4 + 1/4 + 1/2 = 3/2$. This is in parallel with a single edge, so the overall resistance is $1/(1 + 2/3) = 3/5$. Similar but slightly more complicated arguments give the resistance between non-adjacent terminals as $4/5$. So the total is $15 \cdot 3/5 + 30 \cdot 4/5 = 33$, and the average is $33/45 = 11/15$, agreeing with the eigenvalue calculation.

The Matrix-Tree Theorem

Theorem

Let G be a connected graph on n vertices. Then the following three quantities are equal:

1. the number of spanning trees of G ;
2. $(\lambda_2 \cdots \lambda_n) / n$, where $\lambda_2, \dots, \lambda_n$ are the nontrivial Laplacian eigenvalues of G ;
3. any cofactor of $L(G)$ (that is, the determinant of the matrix obtained by deleting row i and column j , multiplied by $(-1)^{i+j}$).

Since one of our criteria for a good network is a large number of spanning trees, this is equivalent to maximizing the geometric mean of the non-trivial Laplacian eigenvalues.




THEOREM OF THE DAY



The Matrix Tree Theorem (Kirchoff) Let G be a graph with n vertices and let $L(G)$ be the $n \times n$ matrix whose entry in row i and column j is defined to be

$$-(\text{the number of edges joining vertex } i \text{ to vertex } j) \quad \text{if } i \neq j, \quad \text{and} \\ \text{the number of edges incident with vertex } i \quad \text{if } i = j.$$

Then the number of spanning trees of G is given by $\det L(G)(i|1)$, where $L(G)(i|1)$ is the matrix obtained by deleting the 1st row and 1st column of $L(G)$.



A	B	C	D	E	F	G
2	-1	-1	0	0	0	0
-1	4	-2	-1	0	0	0
-1	-2	3	-1	0	0	0
0	-1	-1	4	0	0	0
0	0	0	-1	4	0	0
0	0	0	0	0	5	0

The graph on the left has 5 edges on $n = 4$ vertices. Its spanning trees, in the center, are those subsets of $n - 1 = 3$ edges which contain no cyclic paths. The determinant function, \det , yields a single figure from a square matrix or table. It is available in standard spreadsheet applications as the **=DETERM** function. As shown here in the **OpenOffice Calc** package, the calculation will produce the answer 5, since there are exactly 5 spanning trees for the given graph. In fact, any row of $L(G)$, not just the first, and any column, may be deleted in the statement of the theorem without changing the absolute value of the result.

This calculation was first devised by Gustav Kirchhoff in 1847 as a way of obtaining values of current flow in electrical networks. Matrices were first emerging as a powerful mathematical tool about the same time.

Web link: www.math.ku.edu/~jmartin/mc2994/graph1.pdf

Further reading: *Introduction to Graph Theory*, 5th Ed., by Robin J. Wilson, Prentice Hall, 2010, Chapter 4.

From www.themathaday.org by Robin Wherry. This file hosted by London South University

The Cauchy–Binet formula

The proof depends on the **Cauchy–Binet formula**, which says the following:

Theorem

Let A be an $m \times n$ matrix, and B an $n \times m$ matrix, where $m < n$. Then

$$\det(AB) = \sum_X \det(A(X)) \det(B(X)),$$

where X ranges over all m -element subsets of $\{1, \dots, n\}$. Here $A(X)$ is the $m \times m$ matrix whose columns are the columns of A with index in X , and $B(X)$ is the $m \times m$ matrix whose rows are the rows of B with index in X .

Proof of the Matrix-Tree Theorem

Let Q be the incidence matrix of G , so that $QQ^T = L$. Let i be any vertex of G , and let $N = Q_i$ be the matrix obtained by deleting the row of Q indexed by i . It can be shown that, if X is a set of $n - 1$ edges, then $\det(N(X))$ is ± 1 if X is the edge set of a spanning tree, and is 0 otherwise. By the Cauchy–Binet formula, $\det(NN^T)$ is equal to the number of spanning trees. But NN^T is the principal cofactor of $L(G)$ obtained by deleting the row and column indexed by i .

Matrices with row and column sum zero

To finish the proof, let A be any matrix with row and column sums zero, and let $B = A + J$, where J is the all-1 matrix. We evaluate $\det(B)$.

- Replace the first row by the sum of all the rows; this makes the entries in the first row n and doesn't change the other entries; the determinant is unchanged.
- Replace the first column by the sum of all the columns. This makes the first entry n^2 , and the other entries in this column n , and doesn't change the other entries of the matrix; the determinant is unchanged.
- Subtract $1/n$ times the first row from each other row. The elements of the first column, other than the first, become 0; we subtract 1 from all elements not in the first row or column of B , leaving the entries of A ; and the determinant is unchanged.

We conclude $\det(B)$ is n^2 times the $(1,1)$ cofactor of A . It is easily checked that the argument works for any cofactor of A . So all cofactors of A are equal. Finally, the all-1 vector is an eigenvector of B with eigenvalue n , while its other eigenvalues are the same as those of A . Thus $\det(B)$ is n times the product of the nontrivial eigenvalues of A .

Cayley's formula

The Matrix-Tree theorem gives us a simple proof of the famous formula of Cayley:

Theorem

The number of spanning trees in the complete graph on n vertices is equal to n^{n-2} .

For the Laplacian of the complete graph is $nI - J$, where J is the all-1 matrix; its non-trivial Laplacian eigenvalues are all equal to n , and so the number of spanning trees is $n^{n-1}/n = n^{n-2}$.

In our two examples, the number of spanning trees are 2000 and 576 respectively.

Markov chains

A *Markov chain* on a finite state space S is a sequence of random variables with values in S which has no memory: the state at time $n + 1$ depends only on the state at time n .

A Markov chain is defined by a **transition matrix** P , with rows and columns indexed by S , where p_{ij} is the probability of moving from state i to state j in one time step.

As usual, the entries of P are non-negative and the row sums are equal to 1.

Random walks

An important example of a Markov chain is the **random walk** on a graph G . The state space is the vertex set $V(G)$. At time n , if the process is at vertex i , it chooses at random (with equal probabilities) an edge containing i , and at stage $n + 1$ moves to the other end of this edge.

If the graph has no loops, then the probability of moving from i to j is $-L_{ij}/L_{ii}$, where L is the Laplacian. In particular, if the graph is regular with degree d , then $P = I - L/d$.

More generally, $P = I - D^{-1}L$, where D is the diagonal matrix whose (i, i) entry is the number of edges incident with i .

Theory of Markov chains

If a Markov chain has transition matrix P , then the (i, j) entry of P^m is the probability of moving from i to j in m steps.

The Markov chain is **connected** if, for any i and j , there exists m such that $(P^m)_{ij} \neq 0$; it is **aperiodic** if the greatest common divisors of the values of m for which $(P^m)_{ii} \neq 0$ for some i is 1.

A random walk on a graph G is connected if and only if G is connected, and is aperiodic if and only if G is not bipartite.

Theorem

A **connected aperiodic Markov chain** has a unique **limiting distribution**, to which it converges from any starting distribution.

Since the row sums of P are all 1, we see that $Pp^\top = p^\top$, where p is the all-1 vector; our assumptions imply that the multiplicity of 1 as eigenvalue is 1. Now left and right eigenvalues are equal, so there is a vector $q \neq 0$ such that $qP = q$. It can be shown that the entries of q are non-negative; we can normalise it so that their sum is 1. Then q is a probability distribution which is fixed by P , so it is the unique stationary distribution.

Convergence

Suppose that P is symmetric. Then we can write $P = \sum \lambda P_\lambda$ where λ runs over the eigenvalues, and P_λ is the projection onto the λ eigenspace. Then $P^m = \sum \lambda^m P_\lambda$.

It is also true, by the **Perron–Frobenius theorem**, that every eigenvalue λ satisfies $|\lambda| \leq 1$. If the Markov chain is irreducible and aperiodic, then 1 is a simple eigenvalue, and all other eigenvalues have multiplicities strictly less than 1.

Now let x be any non-negative vector whose coordinates sum to 1. We can regard x as the initial probability distribution.

Then we have

$$xP^m = \sum \lambda^m xP_\lambda \rightarrow xP_1$$

as $m \rightarrow \infty$. So $xP_1 = q$ is the limiting distribution, and the convergence to q is like μ^m where μ is the second-largest modulus of an eigenvalue. So the convergence is exponential if μ is not close to 1.

Random walks revisited

For a random walk, we have $P = I - D^{-1}L$. Then

$$D^{1/2}PD^{-1/2} = I - D^{-1/2}LD^{-1/2}.$$

This matrix is symmetric, and is similar to P ; so P is indeed diagonalizable. However, the analysis is a bit more complicated, and not given here.

Its eigenvalues are $1 - \lambda$, where λ is an eigenvalue of the positive semidefinite matrix $D^{-1/2}LD^{-1/2}$, so for rapid convergence we require that the smallest positive eigenvalue of this matrix should be as large as possible.

Thus the problem is a twisted version of the usual problem about the smallest non-trivial Laplacian eigenvalue. If the graph is regular, so that $D = dI$, then it reduces exactly to the former problem.

Other results

The smallest nontrivial Laplacian eigenvalue μ of a graph G is an important parameter which occurs in many other situations. For example, a recent result of Krivelevich and Sudakov asserts that, in a regular graph of valency d on n vertices, if μ is sufficiently large in terms of n and d , then the graph is Hamiltonian.

Summing up

We saw that three important parameters of a connected graph, which are determined by its Laplacian spectrum, are:

- ▶ the harmonic mean of the non-trivial Laplacian eigenvalues, which tells us about the average resistance between pairs of vertices;
- ▶ the geometric mean of the non-trivial Laplacian eigenvalues, which tells us about the number of spanning trees;
- ▶ the smallest non-trivial Laplacian eigenvalue, which is related to the isoperimetric number and the rate of convergence of the random walk on the graph.

In the next lecture, we will see that these are also important parameters in experimental design!