

DECOMPOSABLE FUNCTORS AND THE EXPONENTIAL PRINCIPLE, II

PETER J. CAMERON[†], CHRISTIAN KRATTENTHALER[‡], and THOMAS W. MÜLLER[†]

[†] School of Mathematical Sciences,
Queen Mary & Westfield College, University of London,
Mile End Road, London E1 4NS, United Kingdom.

WWW: <http://www.maths.qmw.ac.uk/~pjc/>

WWW: <http://www.maths.qmw.ac.uk/~twm/>

[‡] Fakultät für Mathematik, Universität Wien,
Nordbergstraße 15, A-1090 Vienna, Austria.

WWW: <http://www.mat.univie.ac.at/~kratt>

Dédié à la mémoire de Pierre Leroux

ABSTRACT. We develop a new setting for the exponential principle in the context of multisort species, where indecomposable objects are generated intrinsically instead of being given in advance. Our approach uses the language of functors and natural transformations (composition operators), and we show that, somewhat surprisingly, a single axiom for the composition already suffices to guarantee validity of the exponential formula. We provide various illustrations of our theory, among which are applications to the enumeration of (semi-)magic squares.

1. INTRODUCTION

One of the corner stones of combinatorial enumeration is a theory which runs under several different names, for instance *theory of species* [7], *theory of exponential structures* [27, Ch. 5], *theory of exponential families* [28], *symbolic method* [12, Part A], *théorie du composé partitionnel* [13], *theory of prefabs* [5], all of which are more or less equivalent. It is probably fair to say that the most elaborate of these theories is the theory of species, as formulated by Joyal [16] (with the functorial concept of species of structures going back to Ehresmann [11]) and further developed by many other authors. It provides the most general framework for such a theory, at the expense of employing a rather abstract language, namely the language of category theory.

A fundamental theorem in each of these theories is the so-called *exponential formula*. Roughly speaking, given a family \mathcal{G} of labelled combinatorial objects (“components”), one produces a larger family \mathcal{F} (“composite objects”) whose objects are obtained by

2000 *Mathematics Subject Classification*. Primary 05A15; Secondary 05A16 05A19 05C30.

Key words and phrases. labelled combinatorial structures, multisort species, exponential principle, functor decomposition, magic squares.

[‡]Research partially supported by the Austrian Science Foundation FWF, grants Z130-N13 and S9607-N13, the latter in the framework of the National Research Network “Analytic Combinatorics and Probabilistic Number Theory”.

putting together various elements of \mathcal{G} . The theorem then states that the (exponential) generating function for \mathcal{F} equals the exponential of the (exponential) generating function for \mathcal{G} .

The aim of the present article is to develop a setting, where one starts with a family \mathcal{F} of labelled combinatorial objects (“composite objects”) and a composition of such objects, and then identifies, in an intrinsic way, a subfamily \mathcal{G} of indecomposable objects (“components”), such that each element of \mathcal{F} can be decomposed into objects from \mathcal{G} , and such that the exponential formula holds for \mathcal{F} and \mathcal{G} . The main point here is that, in contrast to the usual set-up for the exponential formula, indecomposable objects are *not* given in advance, but are defined *inherently* via the composition operation. In particular, our theory leads to a uniform definition of the property to be *indecomposable* for arbitrary labelled combinatorial objects equipped with a composition operator. Interestingly, we show that a single axiom for the composition operator suffices to guarantee validity of the exponential formula. The natural language for formulating a corresponding theory is that of functors and natural transformations. Consequently, our presentation will be in the context of species theory.

For “ordinary” species (1-sort species), such a theory has been presented in [10] on the basis of two axioms for the composition operator. In the present article, we extend this approach to *weighted multisort species*. Moreover, we show that, actually, one of the axioms in [10] can be derived from the other, and that also in our multivariate setting a single axiom suffices. Strictly speaking, our presentation does not cover weighted species (in the sense of [16, Sec. 6], [7, p. 104]) in full generality; rather, we restrict ourselves to the case where the defining functor maps to a category of finite sets, thus avoiding unnecessary technicalities. However, extension to the general case of weighted multisort species is completely straightforward, and is left to the interested reader (see also Footnote 3).

In the next section, we develop the general set-up for our theory. It is formulated within the theory of multisort species, for which we define certain composition operators $\boldsymbol{\eta}$ that are subject to a single axiom, which, in order to be consistent with [10], we call (D1). Furthermore, in the same section, we present our main results. These are two exponential formulae, see Theorems 1 and 4. Theorem 4 refines Theorem 1 by introducing another variable whose powers keep track of the number of “components.” The proof of Theorem 4 requires two general facts about our composition operators $\boldsymbol{\eta}$, which are presented in Propositions 2 and 3. (It is the latter, which, in the less general context of [10], had been assumed as a separate axiom, (D2). As our proof of Proposition 3 shows, this was actually not necessary since, within the general framework, (D2) follows from (D1).) The proofs of Theorems 1 and 4, and of Propositions 2 and 3, are given in Sections 4 and 6, respectively. They require a number of auxiliary results, which are established in Sections 3 and 5, respectively.

Sections 7 and 8 offer illustrations for the theory developed in Sections 2–6. Section 7 presents three simple examples highlighting different aspects of combinatorial situations covered by Theorems 1 and 4. In Section 8, we show how to apply our results to obtain generating function identities for (semi-)magic squares, thereby generalising previous results in the literature.

The final section, Section 9, aims to throw light on the question how far a family \mathcal{F} of labelled combinatorial objects, equipped with a composition operator in the sense indicated above, can be from the species $E(\mathcal{G})$ of sets of indecomposable objects in \mathcal{F} . We find that, roughly speaking, whenever the composition operator is pointwise associative

and commutative, it can be re-constructed, in a sense made precise in Theorem 22, from the standard operation of forming the disjoint union in $E(\mathcal{G})$.

After the present paper was completed, Menni kindly made us aware of his paper [19] (see [20] for further work in this direction), which develops a framework for the exponential principle in the context of symmetric monoidal categories. Menni's results overlap with those of our paper. More precisely, according to [19, Ex. 3.5], a weight-free version of our Theorem 1 can be obtained by applying [19, Prop.1.10] to the discrete groupoid with r elements. It is conceivable that [19, Prop.1.10] also implies a weight-free version of Theorem 4, since the case of $r = 1$ is indeed derived in [19, Ex. 3.5]. For this case, Menni also observes that Axiom (D2) is implied by the set-up. Notwithstanding, we believe our approach to have its own merits. While Menni's arguments make heavy use of categorical notions and constructions (symmetric monoidal categories, Day convolution, Kleisli categories, etc.), our exposition is merely based on the concepts of category, (covariant) functor, and natural transformation. Moreover, our approach readily lends itself to a more general weighted version, an idea not pursued in [19].

2. SET-UP AND MAIN RESULTS

Denote by $\widehat{\mathbf{Set}}$ the category of finite sets and injective mappings, and by \mathbf{Set} the subcategory consisting of finite sets and bijective maps. Moreover, for a positive integer r , let \mathfrak{D}_r be the full subcategory of $\mathbf{Set}^r \times \mathbf{Set}^r$ whose objects are given by

$$\mathrm{Ob}(\mathfrak{D}_r) = \left\{ (\Omega_1, \Omega_2) \in \mathrm{Ob}(\mathbf{Set}^r \times \mathbf{Set}^r) : \Omega_1 \cap \Omega_2 = \emptyset \right\},$$

where $\emptyset = (\emptyset, \dots, \emptyset)$ is the element of $\mathrm{Ob}(\mathbf{Set}^r)$ all of whose components are empty, and intersection is componentwise.

The ingredients needed for our theory are *r-sort species* and certain *composition operators* defined on them. Recall from [16] (or see [7, Def. 4 on p. 102] for a definition avoiding the language of category theory) that, for a positive integer r , an *r-sort species* is a covariant functor $F : \mathbf{Set}^r \rightarrow \mathbf{Set}$. Given r and an *r-sort species* F , the composition operators we have in mind are certain natural transformations η from the functor¹

$$F \times F : \mathfrak{D}_r \xrightarrow{(F,F)} \mathbf{Set} \times \mathbf{Set} \xrightarrow{\times} \mathbf{Set} \xrightarrow{\iota} \widehat{\mathbf{Set}}$$

to the functor

$$F \circ \amalg : \mathfrak{D}_r \xrightarrow{\amalg} \mathbf{Set}^r \xrightarrow{F} \mathbf{Set} \xrightarrow{\iota} \widehat{\mathbf{Set}};$$

that is, families $\eta = (\eta_{(\Omega_1, \Omega_2)})_{(\Omega_1, \Omega_2) \in \mathrm{Ob}(\mathfrak{D}_r)}$ of injective maps,

$$\eta_{(\Omega_1, \Omega_2)} : F[\Omega_1] \times F[\Omega_2] \hookrightarrow F[\Omega_1 \amalg \Omega_2],$$

such that, for every morphism $f : (\Omega_1, \Omega_2) \rightarrow (\tilde{\Omega}_1, \tilde{\Omega}_2)$ of \mathfrak{D}_r , the diagram

$$\begin{array}{ccc} F[\Omega_1] \times F[\Omega_2] & \xrightarrow{\eta_{(\Omega_1, \Omega_2)}} & F[\Omega_1 \amalg \Omega_2] \\ (F \times F)[f] \downarrow & & \downarrow (F \circ \amalg)[f] \\ F[\tilde{\Omega}_1] \times F[\tilde{\Omega}_2] & \xrightarrow{\eta_{(\tilde{\Omega}_1, \tilde{\Omega}_2)}} & F[\tilde{\Omega}_1 \amalg \tilde{\Omega}_2] \end{array} \quad (2.1)$$

commutes. Here, \times is the natural product (Cartesian product) in the category of sets, \amalg is the natural coproduct (componentwise disjoint union) in the category \mathbf{Set}^r and in the category \mathbf{Set} (relying on the context to clarify the intended meaning), and $\iota : \mathbf{Set} \rightarrow \widehat{\mathbf{Set}}$ is the inclusion functor. In what follows, the set-theoretic operations $\cap, \cup, -$ as well as

¹The introduction of the category \mathfrak{D}_r corrects a slight imprecision in the set-up of [10].

the inclusion relation \subseteq and $|$ (restriction of morphisms) in \mathbf{Set}^r are all understood to be componentwise.² We shall most of the time drop the indices of η -maps when they are clear from the context, thus writing $\eta(F[\Omega_1] \times F[\Omega_2])$ instead of $\eta_{(\Omega_1, \Omega_2)}(F[\Omega_1] \times F[\Omega_2])$, for example. We shall think of the elements of a set $\eta(F[\Omega_1] \times F[\Omega_2])$ as *composite objects* within $F[\Omega_1 \amalg \Omega_2]$.

Given an r -sort species F and a composition operator η as above, the next step is to identify the subset $F_\eta[\Omega]$ of “*indecomposable*” elements of a set $F[\Omega]$. It is most natural to define $F_\eta : \text{Ob}(\mathbf{Set}^r) \rightarrow \text{Ob}(\mathbf{Set})$ via

$$F_\eta[\Omega] := \begin{cases} F[\Omega] - \bigcup_{\substack{(\mathbf{I}, \mathbf{J}) \in \text{Ob}(\mathfrak{D}_r) \\ \mathbf{I} \amalg \mathbf{J} = \Omega \\ \mathbf{I} \neq \emptyset \neq \mathbf{J}}} \eta(F[\mathbf{I}] \times F[\mathbf{J}]), & \Omega \neq \emptyset \\ \emptyset, & \Omega = \emptyset \end{cases}, \quad \Omega \in \text{Ob}(\mathbf{Set}^r).$$

At this point, F_η is just defined as a *map* from $\text{Ob}(\mathbf{Set}^r)$ to $\text{Ob}(\mathbf{Set})$. In Lemma 14 in Section 3 we shall show that F_η is in fact a functor, that is, an r -sort species.

Not every natural transformation η is suited for giving rise to an exponential principle. We present the single axiom which is needed for this purpose next. Given F , we call a natural transformation $\eta : F \times F \rightarrow F \circ \amalg$ a *composition operator* of F , if Axiom (D1) below holds.

(D1) For each $\Omega \in \text{Ob}(\mathbf{Set}^r)$ and any two partitions $(\Omega_1, \Omega_2), (\tilde{\Omega}_1, \tilde{\Omega}_2) \in \text{Ob}(\mathfrak{D}_r)$ of Ω into disjoint parts,

$$\Omega_1 \amalg \Omega_2 = \Omega = \tilde{\Omega}_1 \amalg \tilde{\Omega}_2,$$

we have that

$$\eta(F[\Omega_1] \times F[\Omega_2]) \cap \eta(F[\tilde{\Omega}_1] \times F[\tilde{\Omega}_2]) = \eta(\eta(F[\Omega_{11}] \times F[\Omega_{12}]) \times \eta(F[\Omega_{21}] \times F[\Omega_{22}])), \quad (2.2)$$

where $\Omega_{ij} := \Omega_i \cap \tilde{\Omega}_j$ for $i, j \in \{1, 2\}$.

An r -sort species F will be called *decomposable*, if $F \neq \emptyset$ (that is, $F[\Omega] \neq \emptyset$ for some $\Omega \in \text{Ob}(\mathbf{Set}^r)$), and if F admits some composition operator η .

Next, we define *weights* on (F, η) . Fix a commutative ring Λ which contains the rational numbers. A family $\mathbf{w} = (w_\Omega)_{\Omega \in \text{Ob}(\mathbf{Set}^r)}$ of maps $w_\Omega : F[\Omega] \rightarrow \Lambda$ is termed a Λ -*weight* on (F, η) , if the following three conditions hold:

(W0) For all $x \in F[\emptyset]$, we have $w_\emptyset(x) = 1$.

(W1) For each morphism $\mathbf{f} : \Omega_1 \rightarrow \Omega_2$ of \mathbf{Set}^r , the diagram

$$\begin{array}{ccc} F[\Omega_1] & \xrightarrow{w_{\Omega_1}} & \Lambda \\ F[\mathbf{f}] \downarrow & & \downarrow \text{id}_\Lambda \\ F[\Omega_2] & \xrightarrow{w_{\Omega_2}} & \Lambda \end{array}$$

commutes.

²Throughout this paper, we use the symbol $-$ to denote the difference of sets.

(W2) For each pair $(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_r)$, the diagram

$$\begin{array}{ccc} F_\eta[\Omega_1] \times F[\Omega_2] & \xrightarrow{\eta_{(\Omega_1, \Omega_2)}|_{F_\eta[\Omega_1] \times F[\Omega_2]}} & F[\Omega_1 \amalg \Omega_2] \\ w_{\Omega_1}|_{F_\eta[\Omega_1]} \times w_{\Omega_2} \downarrow & & \downarrow w_{\Omega_1 \amalg \Omega_2} \\ \Lambda \times \Lambda & \xrightarrow{\text{multiplication in } \Lambda} & \Lambda \end{array}$$

commutes.

Here, (W0) and (W1) make F a *weighted* r -sort species (cf. [7, p. 104]), whereas (W2) demands (in a weak form) the Λ -weight \mathbf{w} to be compatible with the composition operator η .³ In Section 9, we shall also need the concept of a *weak* Λ -weight, by which we mean a collection $\mathbf{w} = (w_\Omega)_{\Omega \in \text{Ob}(\text{Set}^r)}$ of mappings as above satisfying (W0) and (W1), but not necessarily (W2).

Given a Λ -weight $\mathbf{w} = (w_\Omega)_{\Omega \in \text{Ob}(\text{Set}^r)}$ on (F, η) , we define the corresponding *exponential generating functions* for F and F_η , respectively, by⁴

$$\begin{aligned} \text{GF}_F(z_1, \dots, z_r) &:= \sum_{n_1, \dots, n_r \geq 0} \sum_{x \in F([n_1], \dots, [n_r])} w_{([n_1], \dots, [n_r])}(x) \frac{z_1^{n_1} \cdots z_r^{n_r}}{n_1! \cdots n_r!}, \\ \text{GF}_{F_\eta}(z_1, \dots, z_r) &:= \sum_{n_1, \dots, n_r \geq 0} \sum_{x \in F_\eta([n_1], \dots, [n_r])} w_{([n_1], \dots, [n_r])}(x) \frac{z_1^{n_1} \cdots z_r^{n_r}}{n_1! \cdots n_r!}, \end{aligned}$$

where we suppress the dependence on \mathbf{w} in the notation for better readability.

We are now ready to state our first main result, an *exponential principle*, which generalises Part (a) of the main result in [10].

Theorem 1. *Let r be a positive integer, $F : \text{Set}^r \rightarrow \text{Set}$ an r -sort species, and let $\eta : F \times F \rightarrow F \circ \amalg$ be a natural transformation. If F is decomposable and η is a composition operator for F , then the generating functions GF_F and GF_{F_η} are connected via the relation*

$$\text{GF}_F(z_1, \dots, z_r) = \exp(\text{GF}_{F_\eta}(z_1, \dots, z_r)). \quad (2.3)$$

The proof of Theorem 1 is given in Section 4. It requires several preparatory results, which are established in the next section.

In analogy to [10], there is a refinement of Theorem 1 in the spirit of [17, 21], which we explain next. Making use of the map F_η defined above, we define a sequence of mappings

$$F_\eta^{(k)} : \text{Ob}(\text{Set}^r) \rightarrow \text{Ob}(\text{Set}), \quad k \geq 0,$$

with the property that $F_\eta^{(k)}[\Omega] \subseteq F[\Omega]$ by induction on k via

$$F_\eta^{(0)}[\Omega] := \begin{cases} F[\emptyset], & \Omega = \emptyset \\ \emptyset, & \Omega \neq \emptyset \end{cases}$$

³As already remarked in the introduction, it would be easy to generalise our set-up to cover weighted multisort species in full generality, by relaxing the condition that $F[\Omega]$ needs to be finite, and requiring instead that each preimage $w_\Omega^{-1}(\lambda)$ is finite and that the ring Λ is *multiplicatively finite*, in the sense that the number of different product representations of $\lambda \in \Lambda$ is always finite.

⁴For a non-negative integer n , we write $[n]$ for the standard set $\{1, 2, \dots, n\}$ of cardinality n .

and

$$F_{\boldsymbol{\eta}}^{(k)}[\boldsymbol{\Omega}] := \bigcup_{\substack{\boldsymbol{\Omega}_1 \in \text{Ob}(\mathbf{Set}^r) \\ \boldsymbol{\Omega}_1 \subseteq \boldsymbol{\Omega}}} \eta(F_{\boldsymbol{\eta}}[\boldsymbol{\Omega}_1] \times F_{\boldsymbol{\eta}}^{(k-1)}[\boldsymbol{\Omega} - \boldsymbol{\Omega}_1]), \quad k \geq 1. \quad (2.4)$$

As the definition suggests, one should think of $F_{\boldsymbol{\eta}}^{(k)}[\boldsymbol{\Omega}]$ as the subset of objects in $F[\boldsymbol{\Omega}]$ consisting of exactly k “indecomposable” elements (components).

An immediate induction on

$$\|\boldsymbol{\Omega}\| = \|(\Omega^{(1)}, \Omega^{(2)}, \dots, \Omega^{(r)})\| := \sum_{j=1}^r |\Omega^{(j)}| \quad (2.5)$$

shows that

$$F_{\boldsymbol{\eta}}^{(k)}[\boldsymbol{\Omega}] = \emptyset, \quad k > \|\boldsymbol{\Omega}\|. \quad (2.6)$$

Again, by definition, $F_{\boldsymbol{\eta}}^{(k)}$ is just a *map* from $\text{Ob}(\mathbf{Set}^r)$ to $\text{Ob}(\mathbf{Set})$. In Lemma 16 in Section 5 we shall show that $F_{\boldsymbol{\eta}}^{(k)}$ is in fact a functor, that is, an r -sort species.

It is not difficult to see that, for any $\boldsymbol{\Omega} \in \text{Ob}(\mathbf{Set}^r)$, the sets $F_{\boldsymbol{\eta}}^{(k)}[\boldsymbol{\Omega}]$ with $k = 0, 1, 2, \dots$ cover all of $F[\boldsymbol{\Omega}]$.

Proposition 2. *For every $\boldsymbol{\Omega} \in \text{Ob}(\mathbf{Set}^r)$, we have*

$$F[\boldsymbol{\Omega}] = \bigcup_{k \geq 0} F_{\boldsymbol{\eta}}^{(k)}[\boldsymbol{\Omega}]. \quad (2.7)$$

The proof of Proposition 2 can be found in Section 6.

In [10], a second axiom, (D2), was imposed on the composition operators $\boldsymbol{\eta}$ for obtaining a refined exponential principle that takes into account the filtration given by the sets $F_{\boldsymbol{\eta}}^{(k)}[\boldsymbol{\Omega}]$ appearing on the right-hand side of (2.7) (for the case of 1-sort species): it required pairwise disjointness of the sets $F_{\boldsymbol{\eta}}^{(k)}[\boldsymbol{\Omega}]$ for $k = 0, 1, 2, \dots$. The proposition below says that, actually, this assertion is a consequence of Axiom (D1) (within the general set-up), and this is even true for multisort species.

Proposition 3. *If $F : \mathbf{Set}^r \rightarrow \mathbf{Set}$ is an r -sort species and $\boldsymbol{\eta}$ is a composition operator for F , then we have, for⁵ $k, \ell \in \mathbb{N}_0$ and $k \neq \ell$,*

$$F_{\boldsymbol{\eta}}^{(k)}[\boldsymbol{\Omega}] \cap F_{\boldsymbol{\eta}}^{(\ell)}[\boldsymbol{\Omega}] = \emptyset, \quad \boldsymbol{\Omega} \in \text{Ob}(\mathbf{Set}^r). \quad (2.8)$$

Proposition 3 is also proved in Section 6. Its proof depends crucially on the fact that “ $\boldsymbol{\eta}$ -bracketings” of F -sets and $F_{\boldsymbol{\eta}}$ -sets do not depend on the order of the terms $F[\boldsymbol{\Omega}]$ respectively $F_{\boldsymbol{\eta}}[\boldsymbol{\Omega}]$ involved, nor on the type of bracketing used; see Lemmas 11 and 15 in Sections 3 and 5, respectively. From a technical point of view, this is the decisive improvement over the results in [10], and it is the reason that the dependence of Axiom (D2) from Axiom (D1) was not observed there.

Given a Λ -weight \boldsymbol{w} on $(F, \boldsymbol{\eta})$, the above propositions allow us to refine the weighting to

$$\tilde{w}_{\boldsymbol{\Omega}}(x) := y^k w_{\boldsymbol{\Omega}}(x), \quad x \in F_{\boldsymbol{\eta}}^{(k)}[\boldsymbol{\Omega}].$$

⁵We denote by \mathbb{N}_0 the set of non-negative integers.

We then can define the refined generating function

$$\begin{aligned} \widetilde{\text{GF}}_F(z_1, \dots, z_r, y) &:= \sum_{n_1, \dots, n_r \geq 0} \sum_{x \in F([n_1], \dots, [n_r])} \widetilde{w}_{([n_1], \dots, [n_r])}(x) \frac{z_1^{n_1} \cdots z_r^{n_r}}{n_1! \cdots n_r!} \\ &= \sum_{n_1, \dots, n_r \geq 0} \sum_{k \geq 0} \sum_{x \in F_\eta^{(k)}([n_1], \dots, [n_r])} y^k w_{([n_1], \dots, [n_r])}(x) \frac{z_1^{n_1} \cdots z_r^{n_r}}{n_1! \cdots n_r!}. \end{aligned}$$

With the above notation, we have the following refinement of Theorem 1, which is our second main result.

Theorem 4. *Under the hypotheses of Theorem 1, we have*

$$\widetilde{\text{GF}}_F(z_1, \dots, z_r, y) = \exp(y \text{GF}_{F_\eta}(z_1, \dots, z_r)), \quad (2.9)$$

as well as

$$\widetilde{\text{GF}}_F(z_1, \dots, z_r, y) = (\text{GF}_F(z_1, \dots, z_r))^y. \quad (2.10)$$

The proof of Theorem 4 is given in Section 6, as a simple consequence of (the proof of) Proposition 3.

3. AUXILIARY RESULTS, I

The purpose of this section is to establish several lemmas, which will be needed in the next section in the proof of Theorem 1. At the same time, they also form the basis for the proofs of the auxiliary results in Section 5, which eventually will lead to proofs of Propositions 2 and 3, and of Theorem 4, in Section 6. In all of this section, we assume that F is a decomposable r -sort species with composition operator η .

Lemma 5. *We have $|F[\emptyset]| = 1$.*

Proof. By the injectivity of $\eta_{(\emptyset, \emptyset)} : F[\emptyset] \times F[\emptyset] \rightarrow F[\emptyset]$, the set $F[\emptyset]$ is either empty or a 1-set. Suppose that $F[\emptyset] = \emptyset$. Choose $\Omega_1 = (\Omega_1^{(1)}, \dots, \Omega_1^{(r)}) \in \text{Ob}(\mathbf{Set}^r)$ with $F[\Omega_1] \neq \emptyset$, and $\Omega_2 = (\Omega_2^{(1)}, \dots, \Omega_2^{(r)}) \in \text{Ob}(\mathbf{Set}^r)$ such that $\Omega_1 \cap \Omega_2 = \emptyset$ and $|\Omega_1^{(i)}| = |\Omega_2^{(i)}|$ for $1 \leq i \leq r$. Now consider (D1) for the partition $\Omega := \Omega_1 \amalg \Omega_2$, $\widetilde{\Omega}_i := \Omega_i$ for $i = 1, 2$. By the functoriality of F , we also have $F[\Omega_2] \neq \emptyset$ and, consequently, the left-hand side of (2.2) is non-empty, whereas the right-hand side of (2.2) would be empty in case $F[\emptyset] = \emptyset$, a contradiction. \square

Lemma 6 (COMMUTATIVITY FOR (F, η)). *For every pair $(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_r)$, we have*

$$\eta(F[\Omega_1] \times F[\Omega_2]) = \eta(F[\Omega_2] \times F[\Omega_1]). \quad (3.1)$$

Proof. Applying Axiom (D1) to the partitions

$$\Omega := \Omega_1 \amalg \Omega_2 = \Omega_2 \amalg \Omega_1,$$

we find that

$$\begin{aligned} \mathfrak{J} &:= \eta(F[\Omega_1] \times F[\Omega_2]) \cap \eta(F[\Omega_2] \times F[\Omega_1]) \\ &= \eta(\eta(F[\Omega_1 \cap \Omega_2] \times F[\Omega_1]) \times \eta(F[\Omega_2] \times F[\Omega_1 \cap \Omega_2])). \end{aligned}$$

By Lemma 5 and injectivity of the η -maps, the map

$$\eta_{(\emptyset, \Omega_1)} : F[\emptyset] \times F[\Omega_1] \rightarrow F[\Omega_1]$$

is surjective; that is,

$$\eta(F[\Omega_1 \cap \Omega_2] \times F[\Omega_1]) = F[\Omega_1].$$

Similarly, we have

$$\eta(F[\Omega_2] \times F[\Omega_1 \cap \Omega_2]) = F[\Omega_2].$$

Thus,

$$\mathfrak{J} = \eta(F[\Omega_1] \times F[\Omega_2]).$$

By an analogous application of (D1) and Lemma 5 to the partitions

$$\Omega = \Omega_2 \amalg \Omega_1 = \Omega_1 \amalg \Omega_2,$$

we find that

$$\mathfrak{J} = \eta(F[\Omega_2] \times F[\Omega_1]),$$

and the proof is complete. \square

Lemma 7 (3-ASSOCIATIVITY FOR (F, η)). *For pairwise disjoint $\Omega_1, \Omega_2, \Omega_3 \in \text{Ob}(\mathbf{Set}^r)$, we have*

$$\eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times F[\Omega_3]) = \eta(F[\Omega_1] \times \eta(F[\Omega_2] \times F[\Omega_3])). \quad (3.2)$$

Proof. We shall show that both sides of (3.2) equal the intersection

$$\mathfrak{J} := \eta(F[\Omega_1 \amalg \Omega_2] \times F[\Omega_3]) \cap \eta(F[\Omega_1] \times F[\Omega_2 \amalg \Omega_3]).$$

Applying (D1) to the partitions

$$\Omega := (\Omega_1 \amalg \Omega_2) \amalg \Omega_3 = \Omega_1 \amalg (\Omega_2 \amalg \Omega_3),$$

we find that

$$\mathfrak{J} = \eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times \eta(F[\Omega_1 \cap \Omega_3] \times F[\Omega_3]));$$

and, arguing as in the proof of Lemma 6, this equation simplifies to

$$\mathfrak{J} = \eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times F[\Omega_3]).$$

The same argument, when applied to the partitions

$$\Omega = \Omega_1 \amalg (\Omega_2 \amalg \Omega_3) = (\Omega_1 \amalg \Omega_2) \amalg \Omega_3,$$

yields

$$\mathfrak{J} = \eta(F[\Omega_1] \times \eta(F[\Omega_2] \times F[\Omega_3])),$$

whence (3.2). \square

For the sake of convenience, for pairwise disjoint elements $\Omega_1, \dots, \Omega_m \in \text{Ob}(\mathbf{Set}^r)$, let us call expressions formed by applying η -maps to $F[\Omega_1], \dots, F[\Omega_m]$ (in any order), with each set $F[\Omega_i]$ occurring exactly once, an η -bracketing of $F[\Omega_1], \dots, F[\Omega_m]$. More formally, for any $\Omega \in \text{Ob}(\mathbf{Set}^r)$, we call the set $F[\Omega]$ an η -bracketing of $F[\Omega]$; and, given an η -bracketing B_1 of $F[\Omega_{i_1}], \dots, F[\Omega_{i_k}]$ and an η -bracketing B_2 of $F[\Omega_{i_{k+1}}], \dots, F[\Omega_{i_m}]$, with $1 \leq k \leq m-1$ and $\{i_1, i_2, \dots, i_m\} = \{1, 2, \dots, m\}$, the expression $\eta(B_1 \times B_2)$ is, by definition, an η -bracketing of $F[\Omega_1], \dots, F[\Omega_m]$. For example, the left-hand side and the right-hand side of (3.2) are two possible η -bracketings of $F[\Omega_1], F[\Omega_2], F[\Omega_3]$, as are

$$\eta(\eta(F[\Omega_3] \times F[\Omega_2]) \times F[\Omega_1]) \quad \text{and} \quad \eta(F[\Omega_3] \times \eta(F[\Omega_1] \times F[\Omega_2])).$$

A simple consequence of Lemma 6 and (the proof of) Lemma 7 is the following fact.

Corollary 8. *All η -bracketings of $F[\Omega_1], F[\Omega_2], F[\Omega_3]$ are equal to*

$$\eta(F[\Omega_1 \amalg \Omega_2] \times F[\Omega_3]) \cap \eta(F[\Omega_1 \amalg \Omega_3] \times F[\Omega_2]) \cap \eta(F[\Omega_2 \amalg \Omega_3] \times F[\Omega_1]). \quad (3.3)$$

Proof. From the proof of Lemma 7, we know that

$$\eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times F[\Omega_3]) = \eta(F[\Omega_1 \amalg \Omega_2] \times F[\Omega_3]) \cap \eta(F[\Omega_1] \times F[\Omega_2 \amalg \Omega_3]).$$

If, in this equation, we interchange Ω_1 and Ω_2 and use Lemma 6 (commutativity), then we obtain

$$\eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times F[\Omega_3]) = \eta(F[\Omega_1 \amalg \Omega_2] \times F[\Omega_3]) \cap \eta(F[\Omega_2] \times F[\Omega_1 \amalg \Omega_3]).$$

Both equations together, plus another application of Lemma 6, imply our claim. \square

Lemma 9 (4-PERMUTABILITY FOR (F, η)). *If $\Omega_1, \Omega_2, \Omega_3, \Omega_4$ are pairwise disjoint elements of $\text{Ob}(\text{Set}^r)$, then all η -bracketings of $F[\Omega_1], F[\Omega_2], F[\Omega_3], F[\Omega_4]$ are equal to each other.*

Proof. The possible η -bracketings of $F[\Omega_1], F[\Omega_2], F[\Omega_3], F[\Omega_4]$ are

$$\eta(\eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times F[\Omega_3]) \times F[\Omega_4]), \quad (3.4)$$

$$\eta(\eta(F[\Omega_1] \times \eta(F[\Omega_2] \times F[\Omega_3])) \times F[\Omega_4]), \quad (3.5)$$

$$\eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times \eta(F[\Omega_3] \times F[\Omega_4])), \quad (3.6)$$

$$\eta(F[\Omega_1] \times \eta(\eta(F[\Omega_2] \times F[\Omega_3]) \times F[\Omega_4])), \quad (3.7)$$

$$\eta(F[\Omega_1] \times \eta(F[\Omega_2] \times \eta(F[\Omega_3] \times F[\Omega_4]))), \quad (3.8)$$

together with all expressions arising from the above by permuting $F[\Omega_1], F[\Omega_2], F[\Omega_3], F[\Omega_4]$. By Lemma 7 (3-associativity), the bracketing (3.4) equals the bracketing (3.5), and the bracketing (3.7) equals the bracketing (3.8). It suffices therefore to prove the equality of (3.4), (3.6), (3.8), and all expressions arising from these three by permuting $F[\Omega_1], F[\Omega_2], F[\Omega_3], F[\Omega_4]$.

Applying (D1) to the partitions

$$\Omega := (\Omega_1 \amalg \Omega_2) \amalg (\Omega_3 \amalg \Omega_4) = (\Omega_1 \amalg \Omega_3) \amalg (\Omega_2 \amalg \Omega_4),$$

we find that

$$\begin{aligned} \eta(F[\Omega_1 \amalg \Omega_2] \times F[\Omega_3 \amalg \Omega_4]) \cap \eta(F[\Omega_1 \amalg \Omega_3] \times F[\Omega_2 \amalg \Omega_4]) \\ = \eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times \eta(F[\Omega_3] \times F[\Omega_4])). \end{aligned}$$

Using this equation, as well as the one which arises by interchanging Ω_1 and Ω_2 and applying Lemma 6 (commutativity) on the resulting right-hand side, we obtain

$$\begin{aligned} \eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times \eta(F[\Omega_3] \times F[\Omega_4])) \\ = \eta(F[\Omega_1 \amalg \Omega_2] \times F[\Omega_3 \amalg \Omega_4]) \cap \eta(F[\Omega_1 \amalg \Omega_3] \times F[\Omega_2 \amalg \Omega_4]) \\ \cap \eta(F[\Omega_2 \amalg \Omega_3] \times F[\Omega_1 \amalg \Omega_4]). \quad (3.9) \end{aligned}$$

On the other hand, by Corollary 8, the expression

$$\eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times F[\Omega_3])$$

equals the expression (3.3). Therefore, ‘‘bracketing’’ these two expressions by $\eta(\cdot \times F[\Omega_4])$, using the injectivity of η , and applying Lemma 7 (3-associativity) to the expression resulting from (3.3), we arrive at

$$\begin{aligned} \eta(\eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times F[\Omega_3]) \times F[\Omega_4]) \\ = \eta(F[\Omega_1 \amalg \Omega_2] \times \eta(F[\Omega_3] \times F[\Omega_4])) \cap \eta(F[\Omega_1 \amalg \Omega_3] \times \eta(F[\Omega_2] \times F[\Omega_4])) \\ \cap \eta(F[\Omega_2 \amalg \Omega_3] \times \eta(F[\Omega_1] \times F[\Omega_4])). \quad (3.10) \end{aligned}$$

Since $\eta(F[\Omega_3] \times F[\Omega_4]) \subseteq F[\Omega_3 \amalg \Omega_4]$, and similar inclusions hold for other combinations of the Ω_i 's, we have

$$\begin{aligned} \eta(F[\Omega_1 \amalg \Omega_2] \times \eta(F[\Omega_3] \times F[\Omega_4])) &\subseteq \eta(F[\Omega_1 \amalg \Omega_2] \times F[\Omega_3 \amalg \Omega_4]), \\ \eta(F[\Omega_1 \amalg \Omega_3] \times \eta(F[\Omega_2] \times F[\Omega_4])) &\subseteq \eta(F[\Omega_1 \amalg \Omega_3] \times F[\Omega_2 \amalg \Omega_4]), \\ \eta(F[\Omega_2 \amalg \Omega_3] \times \eta(F[\Omega_1] \times F[\Omega_4])) &\subseteq \eta(F[\Omega_2 \amalg \Omega_3] \times F[\Omega_1 \amalg \Omega_4]). \end{aligned}$$

Altogether, these inclusions imply that the right-hand side of (3.10) is contained in the right-hand side of (3.9). We infer that

$$\eta(\eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times F[\Omega_3]) \times F[\Omega_4]) \subseteq \eta(\eta(F[\Omega_1] \times F[\Omega_2]) \times \eta(F[\Omega_3] \times F[\Omega_4])). \quad (3.11)$$

However, by injectivity of the η -maps, both sides of (3.11) have cardinality

$$|F[\Omega_1]| \cdot |F[\Omega_2]| \cdot |F[\Omega_3]| \cdot |F[\Omega_4]|.$$

Hence, they must be equal, which proves the equality of (3.4) and (3.6).

An analogous argument proves equality of (3.6) and (3.8).

Since, by Lemma 6 (commutativity), the right-hand side of (3.9) is invariant under permutation of $\Omega_1, \Omega_2, \Omega_3, \Omega_4$, the proof is complete. \square

The (proof of the) above lemma, combined with Lemma 6 and Corollary 8, leads to the following observation.

Corollary 10. *All η -bracketings of $F[\Omega_1], F[\Omega_2], F[\Omega_3], F[\Omega_4]$ are equal to*

$$\begin{aligned} &\eta(F[\Omega_1 \amalg \Omega_2] \times F[\Omega_3 \amalg \Omega_4]) \cap \eta(F[\Omega_1 \amalg \Omega_3] \times F[\Omega_2 \amalg \Omega_4]) \\ &\quad \cap \eta(F[\Omega_2 \amalg \Omega_3] \times F[\Omega_1 \amalg \Omega_4]) \\ &\quad \cap \eta(F[\Omega_1] \times F[\Omega_2 \amalg \Omega_3 \amalg \Omega_4]) \cap \eta(F[\Omega_2] \times F[\Omega_1 \amalg \Omega_3 \amalg \Omega_4]) \\ &\quad \cap \eta(F[\Omega_3] \times F[\Omega_1 \amalg \Omega_2 \amalg \Omega_4]) \cap \eta(F[\Omega_4] \times F[\Omega_1 \amalg \Omega_2 \amalg \Omega_3]). \end{aligned} \quad (3.12)$$

Proof. By Corollary 8, we have

$$\begin{aligned} &\eta(F[\Omega_1 \amalg \Omega_2] \times \eta(F[\Omega_3] \times F[\Omega_4])) \\ &= \eta(F[\Omega_1 \amalg \Omega_2 \amalg \Omega_3] \times F[\Omega_4]) \cap \eta(F[\Omega_1 \amalg \Omega_2 \amalg \Omega_4] \times F[\Omega_3]) \\ &\quad \cap \eta(F[\Omega_3 \amalg \Omega_4] \times F[\Omega_1 \amalg \Omega_2]). \end{aligned}$$

Analogous identities hold for the other two terms on the right-hand side of (3.10). If we combine this with Lemma 6 (commutativity) and Lemma 9 (4-permutability), then the claim follows immediately. \square

Before we state the general permutability result, let us introduce the following shorthand notation, which will be used in its proof. Given a subset I of $[m]$, where $I = \{i_1, i_2, \dots, i_k\}$ with $i_1 < i_2 < \dots < i_k$, we let Ω_I stand for $\Omega_{i_1} \amalg \dots \amalg \Omega_{i_k}$.

Lemma 11 (*m*-PERMUTABILITY FOR (F, η)). *If $\Omega_1, \dots, \Omega_m$ are pairwise disjoint elements of $\text{Ob}(\text{Set}^r)$, then all η -bracketings of $F[\Omega_1], \dots, F[\Omega_m]$ are equal to each other.*

Proof. We shall prove by induction on m that, for each $m \geq 2$, all $\boldsymbol{\eta}$ -bracketings of $F[\boldsymbol{\Omega}_1], \dots, F[\boldsymbol{\Omega}_m]$ equal

$$\bigcap_{\substack{I, J \subseteq [m] \\ I \cup J = [m], I \cap J = \emptyset \\ I \neq \emptyset \neq J}} \eta(F[\boldsymbol{\Omega}_I] \times F[\boldsymbol{\Omega}_J]). \quad (3.13)$$

For $m = 2$, the assertion follows from Lemma 6 (commutativity). For $m = 3$, the assertion is equivalent to Corollary 8 (again, modulo Lemma 6), and for $m = 4$, the assertion is equivalent to Corollary 10.

Now let $m \geq 5$, and let us suppose that the assertion is true up to $m - 1$. Consider an $\boldsymbol{\eta}$ -bracketing of $\boldsymbol{\Omega}_1, \dots, \boldsymbol{\Omega}_m$. There are three possibilities. Either this bracketing has the form

$$\eta(\eta(E_1) \times \eta(E_2)), \quad (3.14)$$

where E_1 and E_2 are expressions involving $\boldsymbol{\eta}$ -maps and distinct $\boldsymbol{\Omega}_i$'s, each of them involving at least two $\boldsymbol{\Omega}_i$'s, or the form

$$\eta(F[\boldsymbol{\Omega}_i] \times \eta(E_3)), \quad (3.15)$$

where E_3 involves $\boldsymbol{\eta}$ -maps and $\boldsymbol{\Omega}_1, \dots, \boldsymbol{\Omega}_{i-1}, \boldsymbol{\Omega}_{i+1}, \dots, \boldsymbol{\Omega}_m$, for some i , or the form

$$\eta(\eta(E_4) \times F[\boldsymbol{\Omega}_i]), \quad (3.16)$$

where E_4 involves $\boldsymbol{\eta}$ -maps and $\boldsymbol{\Omega}_1, \dots, \boldsymbol{\Omega}_{i-1}, \boldsymbol{\Omega}_{i+1}, \dots, \boldsymbol{\Omega}_m$, for some i .

We start by considering (3.14). Let us assume that E_1 involves all $\boldsymbol{\Omega}_r$'s for $r \in R$, and E_2 involves all $\boldsymbol{\Omega}_s$'s for $s \in S$, with $R \cup S = [m]$, $R \cap S = \emptyset$, $R \neq \emptyset \neq S$. By the inductive hypothesis, we know that

$$\eta(E_1) = \bigcap_{\substack{I_1, J_1 \subseteq R \\ I_1 \cup J_1 = R, I_1 \cap J_1 = \emptyset \\ I_1 \neq \emptyset \neq J_1}} \eta(F[\boldsymbol{\Omega}_{I_1}] \times F[\boldsymbol{\Omega}_{J_1}]) \quad (3.17)$$

and

$$\eta(E_2) = \bigcap_{\substack{I_2, J_2 \subseteq S \\ I_2 \cup J_2 = S, I_2 \cap J_2 = \emptyset \\ I_2 \neq \emptyset \neq J_2}} \eta(F[\boldsymbol{\Omega}_{I_2}] \times F[\boldsymbol{\Omega}_{J_2}]). \quad (3.18)$$

If we substitute (3.17) and (3.18) in (3.14) and use injectivity of the $\boldsymbol{\eta}$ -maps, then we obtain

$$\bigcap_{\substack{I_1, J_1 \subseteq R \\ I_1 \cup J_1 = R, I_1 \cap J_1 = \emptyset \\ I_1 \neq \emptyset \neq J_1}} \bigcap_{\substack{I_2, J_2 \subseteq S \\ I_2 \cup J_2 = S, I_2 \cap J_2 = \emptyset \\ I_2 \neq \emptyset \neq J_2}} \eta(\eta(F[\boldsymbol{\Omega}_{I_1}] \times F[\boldsymbol{\Omega}_{J_1}]) \times \eta(F[\boldsymbol{\Omega}_{I_2}] \times F[\boldsymbol{\Omega}_{J_2}])).$$

We may now apply Corollary 10 to each of the terms on the right-hand side of this equation. It is not difficult to see that, together with Lemma 6 (commutativity), we obtain (3.13).

Next we consider (3.15). By the inductive hypothesis, we know that

$$\eta(E_3) = \bigcap_{\substack{I, J \subseteq [m] - \{i\} \\ I \cup J = [m] - \{i\}, I \cap J = \emptyset \\ I \neq \emptyset \neq J}} \eta(F[\boldsymbol{\Omega}_I] \times F[\boldsymbol{\Omega}_J]).$$

If we substitute this in (3.15) and use injectivity of the η -maps, then we obtain

$$\bigcap_{\substack{I, J \subseteq [m] - \{i\} \\ I \cup J = [m] - \{i\}, I \cap J = \emptyset \\ I \neq \emptyset \neq J}} \eta(F[\Omega_i] \times \eta(F[\Omega_I] \times F[\Omega_J])).$$

We may now apply Corollary 8 to each of the terms on the right-hand side of this equation. It is not difficult to see that, together with Lemma 6 (commutativity), we obtain (3.13).

The argument for (3.16) is analogous. This completes the proof of the lemma. \square

For $\Omega = (\Omega^{(1)}, \dots, \Omega^{(r)}) \in \text{Ob}(\mathbf{Set}^r)$ and an integer $\rho \in [r]$, we write $(\omega, \rho) \in \Omega$ to mean $\omega \in \Omega^{(\rho)}$. This is the concept of *base point* needed in the present context.

Lemma 12. *For non-empty $\Omega \in \text{Ob}(\mathbf{Set}^r)$ and every choice of base point $(\omega, \rho) \in \Omega$, we have*

$$F[\Omega] = \bigcup_{\substack{\Omega_1 \in \text{Ob}(\mathbf{Set}^r) \\ (\omega, \rho) \in \Omega_1 \subseteq \Omega}} \eta(F_{\eta}[\Omega_1] \times F[\Omega - \Omega_1]). \quad (3.19)$$

Proof. Let $x \in F[\Omega]$ be an arbitrary element, and consider the totality of all $\Omega_1 \in \text{Ob}(\mathbf{Set}^r)$ such that $(\omega, \rho) \in \Omega_1 \subseteq \Omega$ and $x \in \eta(F_{\eta}[\Omega_1] \times F[\Omega - \Omega_1])$. Such Ω_1 's do exist; for instance $\Omega_1 = \Omega$ has these properties, since the map

$$\eta_{(\Omega, \emptyset)} : F[\Omega] \times F[\emptyset] \hookrightarrow F[\Omega]$$

is surjective. Among these Ω_1 's we choose one of minimal norm $\|\Omega_1\|$ (recall the definition in (2.5)), say $\Omega_1(x)$. Now suppose that $x \notin \eta(F_{\eta}[\Omega_1(x)] \times F[\Omega - \Omega_1(x)])$. Then, by the definition of F_{η} , the injectivity of η , and the choice of $\Omega_1(x)$, we must have

$$\begin{aligned} x &\in \eta\left((F[\Omega_1(x)] - F_{\eta}[\Omega_1(x)]) \times F[\Omega - \Omega_1(x)]\right) \\ &= \bigcup_{\substack{(I, J) \in \text{Ob}(\mathfrak{D}_r) \\ I \amalg J = \Omega_1(x) \\ I \neq \emptyset \neq J}} \eta\left(\eta(F[I] \times F[J]) \times F[\Omega - \Omega_1(x)]\right). \end{aligned}$$

Consequently, there exists $(I_1, J_1) \in \text{Ob}(\mathfrak{D}_r)$ such that $I_1 \amalg J_1 = \Omega_1(x)$, $I_1 \neq \emptyset \neq J_1$, and

$$x \in \eta\left(\eta(F[I_1] \times F[J_1]) \times F[\Omega - \Omega_1(x)]\right).$$

Using Corollary 8 (3-permutability), we see that the latter set is contained in both $\eta(F[I_1] \times F[\Omega - I_1])$ and $\eta(F[J_1] \times F[\Omega - J_1])$. The base point (ω, ρ) is contained in I_1 or J_1 ; to fix ideas, say $(\omega, \rho) \in I_1$. Hence, we arrive at the assertion that

$$x \in \eta(F[I_1] \times F[\Omega - I_1]), (\omega, \rho) \in I_1 \subseteq \Omega, \|I_1\| < \|\Omega_1(x)\|,$$

contradicting the choice of $\Omega_1(x)$. We conclude that x is indeed contained in

$$\eta(F_{\eta}[\Omega_1(x)] \times F[\Omega - \Omega_1(x)]),$$

and (3.19) is proven. \square

Lemma 13. *The right-hand side of (3.19) is a disjoint union.*

Proof. In the context of Lemma 12, let $\Omega_1, \Omega_2 \in \text{Ob}(\mathbf{Set}^r)$ be such that $(\omega, \rho) \in \Omega_i \subseteq \Omega$ and $\Omega_1 \neq \Omega_2$, say $\Omega_1 \not\subseteq \Omega_2$. It is enough to show that

$$\mathfrak{J} := \eta(F[\Omega_1] \times F[\Omega - \Omega_1]) \cap \eta(F[\Omega_2] \times F[\Omega - \Omega_2])$$

has an empty intersection with $\eta(F_\eta[\Omega_1] \times F[\Omega - \Omega_1])$. But, by (D1), we have

$$\mathfrak{J} \subseteq \eta\left(\eta(F[\Omega_1 \cap \Omega_2] \times F[\Omega_1 - \Omega_2]) \times F[\Omega - \Omega_1]\right),$$

and, by definition of F_η and the fact that $(\omega, \rho) \in \Omega_1 \cap \Omega_2 \neq \emptyset \neq \Omega_1 - \Omega_2$, we have

$$F_\eta[\Omega_1] \cap \eta(F[\Omega_1 \cap \Omega_2] \times F[\Omega_1 - \Omega_2]) = \emptyset.$$

Consequently, by the injectivity of η , we must indeed have

$$\begin{aligned} & \eta(F_\eta[\Omega_1] \times F[\Omega - \Omega_1]) \cap \mathfrak{J} \\ & \subseteq \eta(F_\eta[\Omega_1] \times F[\Omega - \Omega_1]) \cap \eta\left(\eta(F[\Omega_1 \cap \Omega_2] \times F[\Omega_1 - \Omega_2]) \times F[\Omega - \Omega_1]\right) = \emptyset, \end{aligned}$$

as required. \square

Lemma 14 (FUNCTORIALITY OF F_η). *Let $\Omega, \tilde{\Omega} \in \text{Ob}(\mathbf{Set}^r)$, and let $f : \Omega \rightarrow \tilde{\Omega}$ be a morphism. Then*

$$F[f](F_\eta[\Omega]) = F_\eta[\tilde{\Omega}];$$

that is, setting $F_\eta[f] := F[f]|_{F_\eta[\Omega]}$, we get a functor $F_\eta : \mathbf{Set}^r \rightarrow \mathbf{Set}$.

Proof. The assertion is obvious if $\Omega = \emptyset$, so we may suppose that $\Omega \neq \emptyset$. Then, using the naturality of η (that is, the diagram (2.1)), we have

$$\begin{aligned} F[f](F_\eta[\Omega]) &= F[f] \left(F[\Omega] - \bigcup_{\substack{(\mathbf{I}_1, \mathbf{J}_1) \in \text{Ob}(\mathfrak{D}_r) \\ \mathbf{I}_1 \amalg \mathbf{J}_1 = \Omega \\ \mathbf{I}_1 \neq \emptyset \neq \mathbf{J}_1}} \eta(F[\mathbf{I}_1] \times F[\mathbf{J}_1]) \right) \\ &= F[\tilde{\Omega}] - \bigcup_{\substack{(\mathbf{I}_1, \mathbf{J}_1) \in \text{Ob}(\mathfrak{D}_r) \\ \mathbf{I}_1 \amalg \mathbf{J}_1 = \Omega \\ \mathbf{I}_1 \neq \emptyset \neq \mathbf{J}_1}} (F \circ \amalg) [(f|_{\mathbf{I}_1}, f|_{\mathbf{J}_1})] (\eta(F[\mathbf{I}_1] \times F[\mathbf{J}_1])) \\ &= F[\tilde{\Omega}] - \bigcup_{\substack{(\mathbf{I}_1, \mathbf{J}_1) \in \text{Ob}(\mathfrak{D}_r) \\ \mathbf{I}_1 \amalg \mathbf{J}_1 = \Omega \\ \mathbf{I}_1 \neq \emptyset \neq \mathbf{J}_1}} \eta(F[f|_{\mathbf{I}_1}](F[\mathbf{I}_1]) \times F[f|_{\mathbf{J}_1}](F[\mathbf{J}_1])) \\ &= F[\tilde{\Omega}] - \bigcup_{\substack{(\mathbf{I}_2, \mathbf{J}_2) \in \text{Ob}(\mathfrak{D}_r) \\ \mathbf{I}_2 \amalg \mathbf{J}_2 = \tilde{\Omega} \\ \mathbf{I}_2 \neq \emptyset \neq \mathbf{J}_2}} \eta(F[\mathbf{I}_2] \times F[\mathbf{J}_2]) \\ &= F[\tilde{\Omega}]. \end{aligned}$$

\square

4. PROOF OF THEOREM 1

For convenience, let us “extend” the Λ -weight w to subsets of $F[\Omega]$, for all $\Omega \in \text{Ob}(\mathbf{Set}^r)$. To be precise, for $A \subseteq F[\Omega]$, we define

$$w_{\Omega}(A) := \sum_{x \in A} w_{\Omega}(x).$$

Given disjoint $\Omega_1, \Omega_2 \in \text{Ob}(\mathbf{Set}^r)$ and subsets $A_1 \subseteq F_{\eta}[\Omega_1]$ and $A_2 \subseteq F[\Omega_2]$, Axiom (W2) says that

$$w_{\Omega_1 \amalg \Omega_2}(\eta(A_1 \times A_2)) = w_{\Omega_1}(A_1) \cdot w_{\Omega_2}(A_2). \quad (4.1)$$

In the sequel, we shall suppress the indices of weights w for better readability, the indices always being clear from the context.

As a direct consequence of Lemmas 12 and 13, of the injectivity of the η -maps, and of (4.1), we have that, for $n_1, n_2, \dots, n_r \in \mathbb{N}_0$ and $n_{\rho} > 0$,

$$\begin{aligned} & w(F([n_1], \dots, [n_r])) \\ &= \sum_{\substack{\Omega_1^{(i)} \subseteq [n_i] \ (1 \leq i \leq r) \\ 1 \in \Omega_1^{(\rho)}}} w(F_{\eta}[(\Omega_1^{(1)}, \dots, \Omega_1^{(r)})]) \cdot w(F([n_1 - \Omega_1^{(1)}, \dots, [n_r - \Omega_1^{(r)}])). \end{aligned} \quad (4.2)$$

Using the functoriality of F and F_{η} , together with Axiom (W1), each Ω_1 with

$$\Omega_1 = (\Omega_1^{(1)}, \dots, \Omega_1^{(r)}) \subseteq ([n_1], \dots, [n_r]),$$

$(1, \rho) \in \Omega_1$, and cardinalities $|\Omega_1^{(i)}| = \mu_i$ ($1 \leq i \leq r$), is seen to contribute

$$w(F_{\eta}([\mu_1], \dots, [\mu_r])) \cdot w(F([n_1 - \mu_1], \dots, [n_r - \mu_r])) \quad (4.3)$$

to the right-hand side of (4.2). We observe that (4.3) does not depend upon Ω_1 itself, but only on the cardinalities μ_1, \dots, μ_r of the components $\Omega_1^{(1)}, \dots, \Omega_1^{(r)}$. Therefore, the

$$\frac{\mu_{\rho}}{n_{\rho}} \prod_{1 \leq i \leq r} \binom{n_i}{\mu_i}$$

elements $\Omega_1 \in \text{Ob}(\mathbf{Set}^r)$ with $\Omega_1 \subseteq ([n_1], \dots, [n_r])$, $(1, \rho) \in \Omega_1$, and such that $|\Omega_1^{(i)}| = \mu_i$ for $1 \leq i \leq r$, contribute

$$\frac{\mu_{\rho}}{n_{\rho}} \left(\prod_{1 \leq i \leq r} \binom{n_i}{\mu_i} \right) w(F_{\eta}([\mu_1], \dots, [\mu_r])) \cdot w(F([n_1 - \mu_1], \dots, [n_r - \mu_r]))$$

to the right-hand side of (4.2), and we obtain

$$\begin{aligned} w(F([n_1], \dots, [n_r])) &= \sum_{0 \leq \mu_i \leq n_i \ (1 \leq i \leq r)} \frac{\mu_{\rho}}{n_{\rho}} \left(\prod_{1 \leq i \leq r} \binom{n_i}{\mu_i} \right) w(F_{\eta}([\mu_1], \dots, [\mu_r])) \\ &\quad \cdot w(F([n_1 - \mu_1], \dots, [n_r - \mu_r])), \end{aligned} \quad n_{\rho} > 0,$$

or, equivalently,

$$n_\rho \cdot w(F([n_1], \dots, [n_r])) = \sum_{0 \leq \mu_i \leq n_i \ (1 \leq i \leq r)} \mu_\rho \left(\prod_{1 \leq i \leq r} \binom{n_i}{\mu_i} \right) w(F_\eta([[\mu_1], \dots, [\mu_r]])) \cdot w(F([n_1 - \mu_1], \dots, [n_r - \mu_r])), \quad (4.4)$$

as long as $n_\rho > 0$. However, Equation (4.4) holds as well for $n_\rho = 0$, with both sides vanishing, so that we are allowed to drop the restriction $n_\rho > 0$.

Fix $\rho \in [r]$, multiply both sides of (4.4) by

$$z_1^{n_1} \cdots z_{\rho-1}^{n_{\rho-1}} z_\rho^{n_\rho-1} z_{\rho+1}^{n_{\rho+1}} \cdots z_r^{n_r} / (n_1! \cdots n_r!),$$

and sum over all tuples $(n_1, \dots, n_r) \in \mathbb{N}_0^r$, to get

$$\begin{aligned} & \sum_{n_1, \dots, n_r \geq 0} w(F([n_1], \dots, [n_r])) \frac{z_1^{n_1} \cdots z_{\rho-1}^{n_{\rho-1}} z_\rho^{n_\rho-1} z_{\rho+1}^{n_{\rho+1}} \cdots z_r^{n_r}}{n_1! \cdots n_{\rho-1}! (n_\rho - 1)! n_{\rho+1}! \cdots n_r!} \\ &= \sum_{n_1, \dots, n_r \geq 0} \sum_{0 \leq \mu_i \leq n_i \ (1 \leq i \leq r)} \frac{w(F_\eta([[\mu_1], \dots, [\mu_r]]))}{\mu_1! \cdots \mu_{\rho-1}! (\mu_\rho - 1)! \mu_{\rho+1}! \cdots \mu_r!} \\ & \quad \cdot \frac{w(F([n_1 - \mu_1], \dots, [n_r - \mu_r]))}{(n_1 - \mu_1)! \cdots (n_r - \mu_r)!} z_1^{n_1} \cdots z_{\rho-1}^{n_{\rho-1}} z_\rho^{n_\rho-1} z_{\rho+1}^{n_{\rho+1}} \cdots z_r^{n_r}, \quad (4.5) \end{aligned}$$

where $1/(-1)!$ has to be interpreted as 0. The left-hand side equals

$$\frac{\partial \text{GF}_F}{\partial z_\rho},$$

while the right-hand side is identified as

$$\frac{\partial \text{GF}_{F_\eta}}{\partial z_\rho} \text{GF}_F;$$

whence the equations

$$\frac{\partial \text{GF}_F}{\partial z_\rho} = \frac{\partial \text{GF}_{F_\eta}}{\partial z_\rho} \text{GF}_F, \quad 1 \leq \rho \leq r. \quad (4.6)$$

Set

$$Q(z_1, \dots, z_r) := \text{GF}_F(z_1, \dots, z_r) \exp(-\text{GF}_{F_\eta}(z_1, \dots, z_r)).$$

Then, in view of Equations (4.6), the series Q satisfies

$$\frac{\partial Q}{\partial z_\rho} = 0, \quad 1 \leq \rho \leq r.$$

These last equations force Q to be independent of z_1, z_2, \dots, z_r . However, since $\text{GF}_{F_\eta}(0, \dots, 0) = 0$ by definition of F_η , and since $\text{GF}_F(0, \dots, 0) = 1$ by Axiom (W0) and Lemma 5, direct inspection shows that

$$Q(0, 0, \dots, 0) = 1,$$

and (2.3) follows. \square

5. AUXILIARY RESULTS, II

In this section, we complement the results of Section 3 by establishing several further results which will be needed in the proofs of Proposition 3 and Theorem 4, to be given in the next section. The first lemma provides the analogue of Lemma 11 for F_η , namely that arbitrary permutability holds also for η -bracketings of F_η -images (see the subsequent paragraph for the precise definition). All the remaining lemmas concern the maps $F_\eta^{(k)}$. In all of this section, we assume that F is a decomposable r -sort species with composition operator η .

In complete analogy with the corresponding definition in Section 3, we define the concept of an η -bracketing of $F_\eta[\Omega_1], \dots, F_\eta[\Omega_m]$ for pairwise disjoint elements $\Omega_1, \dots, \Omega_m \in \text{Ob}(\mathbf{Set}^r)$: simply replace F by F_η everywhere in the definition just after the proof of Lemma 7.

Lemma 15 (*m -PERMUTABILITY FOR (F_η, η)*). *If $\Omega_1, \dots, \Omega_m$ are pairwise disjoint elements of $\text{Ob}(\mathbf{Set}^r)$, then all η -bracketings of $F_\eta[\Omega_1], \dots, F_\eta[\Omega_m]$ are equal to each other.*

Proof. Since $F_\eta[\emptyset] = \emptyset$ by definition of F_η , our claim holds if at least one of $\Omega_1, \dots, \Omega_m$ equals \emptyset . Hence, we may assume that all of $\Omega_1, \dots, \Omega_m$ are non-empty.

Next we note that, for sets $M_1, \dots, M_m, A_1, \dots, A_m$, we have

$$\begin{aligned} & (M_1 - A_1) \times (M_2 - A_2) \times \cdots \times (M_m - A_m) \\ &= M_1 \times M_2 \times \cdots \times M_m - \left(\bigcup_{k=1}^m M_1 \times \cdots \times M_{k-1} \times A_k \times M_{k+1} \times \cdots \times M_m \right). \end{aligned} \quad (5.1)$$

Now assume that we are given two η -bracketings of $F_\eta[\Omega_1], \dots, F_\eta[\Omega_m]$, say

$$B_\eta(F_\eta[\Omega_1], \dots, F_\eta[\Omega_m]) \quad \text{and} \quad \bar{B}_\eta(F_\eta[\Omega_1], \dots, F_\eta[\Omega_m]).$$

Substituting the definition of F_η into $B_\eta(F_\eta[\Omega_1], \dots, F_\eta[\Omega_m])$, and applying Identity (5.1) plus injectivity of η -maps, we find that

$$\begin{aligned} & B_\eta(F_\eta[\Omega_1], \dots, F_\eta[\Omega_m]) = B_\eta(F[\Omega_1], \dots, F[\Omega_m]) \\ & - \bigcup_{k=1}^m B_\eta \left(F[\Omega_1], \dots, F[\Omega_{k-1}], \bigcup_{\substack{(\mathbf{I}, \mathbf{J}) \in \text{Ob}(\mathfrak{D}_r) \\ \mathbf{I} \amalg \mathbf{J} = \Omega_k \\ \mathbf{I} \neq \emptyset \neq \mathbf{J}}} \eta(F[\mathbf{I}] \times F[\mathbf{J}]), F[\Omega_{k+1}], \dots, F[\Omega_m] \right) \\ &= B_\eta(F[\Omega_1], \dots, F[\Omega_m]) \\ & - \bigcup_{k=1}^m \bigcup_{\substack{(\mathbf{I}, \mathbf{J}) \in \text{Ob}(\mathfrak{D}_r) \\ \mathbf{I} \amalg \mathbf{J} = \Omega_k \\ \mathbf{I} \neq \emptyset \neq \mathbf{J}}} B_\eta(F[\Omega_1], \dots, F[\Omega_{k-1}], \eta(F[\mathbf{I}] \times F[\mathbf{J}]), F[\Omega_{k+1}], \dots, F[\Omega_m]). \end{aligned}$$

The same argument shows that $\bar{B}_\eta(F_\eta[\Omega_1], \dots, F_\eta[\Omega_m])$ equals the last expression where every occurrence of B_η is replaced by \bar{B}_η . By Lemma 11 (m -permutability for (F, η)), we have

$$B_\eta(F[\Omega_1], \dots, F[\Omega_m]) = \bar{B}_\eta(F[\Omega_1], \dots, F[\Omega_m])$$

and

$$\begin{aligned} B_{\eta}(F[\Omega_1], \dots, F[\Omega_{k-1}], \eta(F[\mathbf{I}] \times F[\mathbf{J}]), F[\Omega_{k+1}], \dots, F[\Omega_m]) \\ = \bar{B}_{\eta}(F[\Omega_1], \dots, F[\Omega_{k-1}], \eta(F[\mathbf{I}] \times F[\mathbf{J}]), F[\Omega_{k+1}], \dots, F[\Omega_m]), \end{aligned}$$

hence

$$B_{\eta}(F_{\eta}[\Omega_1], \dots, F_{\eta}[\Omega_m]) = \bar{B}_{\eta}(F_{\eta}[\Omega_1], \dots, F_{\eta}[\Omega_m]),$$

which establishes our claim. \square

Lemma 16 (FUNCTORIALITY OF $F_{\eta}^{(k)}$). *For each morphism $f : \Omega \rightarrow \tilde{\Omega}$ in \mathbf{Set}^r and every integer $k \geq 0$, we have*

$$F[f](F_{\eta}^{(k)}[\Omega]) = F_{\eta}^{(k)}[\tilde{\Omega}].$$

Proof. We use induction on k , our claim being obvious for $k = 0$. Suppose that the assertion holds for $0 \leq k < K$ with some $K \geq 1$. Then, using the definition of $F_{\eta}^{(k)}$, the functoriality of F_{η} already demonstrated in Lemma 14, the inductive hypothesis, as well as the naturality of η , we find that

$$\begin{aligned} F[f](F_{\eta}^{(K)}[\Omega]) &= F[f] \left(\bigcup_{\substack{\Omega' \in \text{Ob}(\mathbf{Set}^r) \\ \Omega' \subseteq \Omega}} \eta(F_{\eta}[\Omega'] \times F_{\eta}^{(K-1)}[\Omega - \Omega']) \right) \\ &= \bigcup_{\substack{\Omega' \in \text{Ob}(\mathbf{Set}^r) \\ \Omega' \subseteq \Omega}} (F \circ \Pi) [(\mathbf{f}|_{\Omega'}, \mathbf{f}|_{\Omega - \Omega'}) (\eta(F_{\eta}[\Omega'] \times F_{\eta}^{(K-1)}[\Omega - \Omega']))] \\ &= \bigcup_{\substack{\Omega' \in \text{Ob}(\mathbf{Set}^r) \\ \Omega' \subseteq \Omega}} \eta(F[\mathbf{f}|_{\Omega'}](F_{\eta}[\Omega']) \times F[\mathbf{f}|_{\Omega - \Omega'}](F_{\eta}^{(K-1)}[\Omega - \Omega'])) \\ &= \bigcup_{\substack{\tilde{\Omega}' \in \text{Ob}(\mathbf{Set}^r) \\ \tilde{\Omega}' \subseteq \tilde{\Omega}}} \eta(F_{\eta}[\tilde{\Omega}'] \times F_{\eta}^{(K-1)}[\tilde{\Omega} - \tilde{\Omega}']) \\ &= F_{\eta}^{(K)}[\tilde{\Omega}]. \end{aligned}$$

\square

Lemma 17. *The functors $F_{\eta}^{(1)}$ and F_{η} coincide.*

Proof. It suffices to show that $F_{\eta}^{(1)}[\Omega] = F_{\eta}[\Omega]$ for every $\Omega \in \text{Ob}(\mathbf{Set}^r)$. By (2.6), this holds if $\Omega = \emptyset$, so assume that $\Omega \neq \emptyset$. Then, using the definition of $F_{\eta}^{(k)}$, the injectivity

of η -maps, Lemma 7 (3-associativity), and Lemma 5, we have

$$\begin{aligned}
F_{\eta}^{(1)}[\Omega] &= \bigcup_{\substack{\Omega_1 \in \text{Ob}(\mathbf{Set}^r) \\ \Omega_1 \subseteq \Omega}} \eta(F_{\eta}[\Omega_1] \times F_{\eta}^{(0)}[\Omega - \Omega_1]) \\
&= \eta(F_{\eta}[\Omega] \times F[\emptyset]) \\
&= \eta\left(\left(F[\Omega] - \bigcup_{\substack{(\mathbf{I}, \mathbf{J}) \in \text{Ob}(\mathfrak{D}_r) \\ \mathbf{I} \amalg \mathbf{J} = \Omega \\ \mathbf{I} \neq \emptyset \neq \mathbf{J}}} \eta(F[\mathbf{I}] \times F[\mathbf{J}])\right) \times F[\emptyset]\right) \\
&= \eta(F[\Omega] \times F[\emptyset]) - \bigcup_{\substack{(\mathbf{I}, \mathbf{J}) \in \text{Ob}(\mathfrak{D}_r) \\ \mathbf{I} \amalg \mathbf{J} = \Omega \\ \mathbf{I} \neq \emptyset \neq \mathbf{J}}} \eta\left(\eta(F[\mathbf{I}] \times F[\mathbf{J}]) \times F[\emptyset]\right) \\
&= F[\Omega] - \bigcup_{\substack{(\mathbf{I}, \mathbf{J}) \in \text{Ob}(\mathfrak{D}_r) \\ \mathbf{I} \amalg \mathbf{J} = \Omega \\ \mathbf{I} \neq \emptyset \neq \mathbf{J}}} \eta(F[\mathbf{I}] \times F[\mathbf{J}]) \\
&= F_{\eta}[\Omega],
\end{aligned}$$

proving our claim. \square

In the next lemma, we require again the concept of a base point, which was introduced just before Lemma 12.

Lemma 18. *For every non-empty $\Omega \in \text{Ob}(\mathbf{Set}^r)$, each choice of base point $(\omega, \rho) \in \Omega$, and every integer $k \geq 1$, we have*

$$F_{\eta}^{(k)}[\Omega] = \coprod_{\substack{\Omega_1 \in \text{Ob}(\mathbf{Set}^r) \\ (\omega, \rho) \in \Omega_1 \subseteq \Omega}} \eta(F_{\eta}[\Omega_1] \times F_{\eta}^{(k-1)}[\Omega - \Omega_1]). \quad (5.2)$$

Proof. The fact that the terms on the right-hand side of (5.2) are pairwise disjoint follows from Lemma 13, since a term $\eta(F_{\eta}[\Omega_1] \times F_{\eta}^{(k-1)}[\Omega - \Omega_1])$ is contained in the larger set $\eta(F_{\eta}[\Omega_1] \times F[\Omega - \Omega_1])$.

An immediate induction using the definition of $F_{\eta}^{(m)}$ shows that, for all $m \geq 2$, we have

$$F_{\eta}^{(m)}[\Omega] = \bigcup_{\substack{\Omega_1, \dots, \Omega_m \in \text{Ob}(\mathbf{Set}^r) \\ \Omega_1 \amalg \dots \amalg \Omega_m = \Omega}} \eta(F_{\eta}[\Omega_1] \times \eta(F_{\eta}[\Omega_2] \times \dots \times \eta(F_{\eta}[\Omega_{m-1}] \times F_{\eta}[\Omega_m]) \dots)). \quad (5.3)$$

We first consider the case where $k = 1$. Here, by definition of $F_{\eta}^{(0)}[\emptyset]$, the only contribution to the union on the right-hand side of (5.2) arises for $\Omega_1 = \Omega$. In that situation, we have

$$\eta(F_{\eta}[\Omega] \times F_{\eta}^{(0)}[\emptyset]) = \eta(F_{\eta}[\Omega] \times F[\emptyset]).$$

However, this is also the only contribution on the right-hand side of the definition of $F_{\eta}^{(1)}$ given in (2.4), thus proving (5.2) for $k = 1$.

Now we consider the case where $k \geq 2$. If $k \geq 3$, then, given $\Omega \in \text{Ob}(\mathbf{Set}^r)$, we substitute the right-hand side of (5.3) with $m = k - 1$ in (5.2). As a result, we obtain

$$\bigcup_{\substack{\Omega_1, \Omega_2, \dots, \Omega_k \in \text{Ob}(\mathbf{Set}^r) \\ \Omega_1 \amalg \Omega_2 \amalg \dots \amalg \Omega_k = \Omega \\ (\omega, \rho) \in \Omega_1}} \eta(F_\eta[\Omega_1] \times \eta(F_\eta[\Omega_2] \times \dots \times \eta(F_\eta[\Omega_{k-1}] \times F_\eta[\Omega_k]) \dots)) \quad (5.4)$$

for the right-hand side of (5.2). We note that Expression (5.4) also agrees with the right-hand side of (5.2) for $k = 2$ (taking into account the fact that we already know that the union on the right-hand side of (5.2) is a disjoint union).

Expression (5.4) is almost (5.3) with $m = k$, except that Ω_1 is distinguished by having to contain the given base point (ω, ρ) . However, by Lemma 15 (m -permutability for (F_η, η)), the ordering of $\Omega_1, \Omega_2, \dots, \Omega_k$ in the η -bracketing in the union on the right-hand side of (5.4) is of no relevance. Thus, the restriction that $(\omega, \rho) \in \Omega_1$ can be dropped. This shows that the right-hand side of (5.2) equals $F_\eta^{(k)}[\Omega]$, as claimed. \square

6. PROOFS OF PROPOSITIONS 2 AND 3, AND OF THEOREM 4

We begin this section with the proof of Proposition 2. With Lemma 18 in hand, we are finally in the position to also establish Proposition 3. Theorem 4 is then a simple consequence of an identity on which the proof of Proposition 3 rests (see (6.1) below). Although the property expressed in this proposition is of a structural nature, our proof relies in fact on a counting argument. It would be desirable to find an alternative approach more in keeping with the actual nature of Proposition 3.

Proof of Proposition 2. We use induction on $\|\Omega\|$, where $\|\cdot\|$ has been defined in (2.5). By (2.6) and the definition of $F_\eta^{(0)}$, the statement holds if $\|\Omega\| = 0$, that is, if $\Omega = \emptyset$. Let Ω be such that $\|\Omega\| = N$ for some integer $N > 0$, and suppose that (2.7) holds for all $\Omega' \in \text{Ob}(\mathbf{Set}^r)$ of norm strictly less than N . Then we have $\Omega \neq \emptyset$, and therefore

$$\begin{aligned} \bigcup_{k \geq 0} F_\eta^{(k)}[\Omega] &= \bigcup_{k \geq 1} F_\eta^{(k)}[\Omega] \\ &= \bigcup_{k \geq 1} \bigcup_{\substack{\Omega_1 \in \text{Ob}(\mathbf{Set}^r) \\ \Omega_1 \subseteq \Omega}} \eta(F_\eta[\Omega_1] \times F_\eta^{(k-1)}[\Omega - \Omega_1]) \\ &= \bigcup_{\substack{\Omega_1 \in \text{Ob}(\mathbf{Set}^r) \\ \Omega_1 \subseteq \Omega}} \eta\left(F_\eta[\Omega_1] \times \left(\bigcup_{k \geq 1} F_\eta^{(k-1)}[\Omega - \Omega_1]\right)\right) \\ &= \bigcup_{\substack{\Omega_1 \in \text{Ob}(\mathbf{Set}^r) \\ \emptyset \neq \Omega_1 \subseteq \Omega}} \eta\left(F_\eta[\Omega_1] \times \left(\bigcup_{k \geq 0} F_\eta^{(k)}[\Omega - \Omega_1]\right)\right) \\ &= \bigcup_{\substack{\Omega_1 \in \text{Ob}(\mathbf{Set}^r) \\ \Omega_1 \subseteq \Omega}} \eta(F_\eta[\Omega_1] \times F[\Omega - \Omega_1]) \\ &= F[\Omega]. \end{aligned}$$

Here, we have used Lemma 12 for the last equality, and the inductive hypothesis in the second but last step (here it is important that $\Omega_1 \neq \emptyset$ in order to guarantee that $\|\Omega - \Omega_1\| < \|\Omega\|$). \square

Proof of Proposition 3 and of Theorem 4. For $k \geq 0$, let us define the generating function for $F_{\boldsymbol{\eta}}^{(k)}$ by

$$\text{GF}_{F_{\boldsymbol{\eta}}^{(k)}}(z_1, \dots, z_r) := \sum_{n_1, \dots, n_r \geq 0} \sum_{x \in F_{\boldsymbol{\eta}}^{(k)}([n_1], \dots, [n_r])} w_{([n_1], \dots, [n_r])}(x) \frac{z_1^{n_1} \cdots z_r^{n_r}}{n_1! \cdots n_r!}.$$

Again, in the sequel, we shall suppress the indices to weights w for better readability, the indices always being clear from the context.

The first step consists in showing that

$$\text{GF}_{F_{\boldsymbol{\eta}}^{(k)}}(z_1, \dots, z_r) = \frac{1}{k!} \left(\text{GF}_{F_{\boldsymbol{\eta}}}(z_1, \dots, z_r) \right)^k. \quad (6.1)$$

By definition of $F_{\boldsymbol{\eta}}^{(0)}$, the left-hand side of (6.1) equals 1, so that (6.1) holds for $k = 0$. Therefore, we may in the sequel assume that $k \geq 1$.

We now proceed in a manner similar to the proof of Theorem 1 given in Section 4. Here, however, we use Lemma 18 instead of Lemmas 12 and 13, and we also need the functoriality of $F_{\boldsymbol{\eta}}^{(m)}$ for $m = 0, 1, 2, \dots$ established in Lemma 16. In this way, we obtain from (5.2) the identity

$$\begin{aligned} n_{\rho} \cdot w(F_{\boldsymbol{\eta}}^{(k)}([n_1], \dots, [n_r])) &= \sum_{0 \leq \mu_i \leq n_i \ (1 \leq i \leq r)} \mu_{\rho} \left(\prod_{1 \leq i \leq r} \binom{n_i}{\mu_i} \right) w(F_{\boldsymbol{\eta}}([[\mu_1], \dots, [\mu_r]])) \\ &\quad \cdot w(F_{\boldsymbol{\eta}}^{(k-1)}([n_1 - \mu_1], \dots, [n_r - \mu_r])). \end{aligned} \quad (6.2)$$

Fixing $\rho \in [r]$, multiplying both sides of (6.2) by

$$z_1^{n_1} \cdots z_{\rho-1}^{n_{\rho-1}} z_{\rho}^{n_{\rho}-1} z_{\rho+1}^{n_{\rho+1}} \cdots z_r^{n_r} / (n_1! \cdots n_r!),$$

and summing over all tuples $(n_1, \dots, n_r) \in \mathbb{N}_0^r$, gives

$$\begin{aligned} &\sum_{n_1, \dots, n_r \geq 0} w(F_{\boldsymbol{\eta}}^{(k)}([n_1], \dots, [n_r])) \frac{z_1^{n_1} \cdots z_{\rho-1}^{n_{\rho-1}} z_{\rho}^{n_{\rho}-1} z_{\rho+1}^{n_{\rho+1}} \cdots z_r^{n_r}}{n_1! \cdots n_{\rho-1}! (n_{\rho} - 1)! n_{\rho+1}! \cdots n_r!} \\ &= \sum_{n_1, \dots, n_r \geq 0} \sum_{0 \leq \mu_i \leq n_i \ (1 \leq i \leq r)} \frac{w(F_{\boldsymbol{\eta}}([[\mu_1], \dots, [\mu_r]]))}{\mu_1! \cdots \mu_{\rho-1}! (\mu_{\rho} - 1)! \mu_{\rho+1}! \cdots \mu_r!} \\ &\quad \cdot \frac{w(F_{\boldsymbol{\eta}}^{(k-1)}([n_1 - \mu_1], \dots, [n_r - \mu_r]))}{(n_1 - \mu_1)! \cdots (n_r - \mu_r)!} z_1^{n_1} \cdots z_{\rho-1}^{n_{\rho-1}} z_{\rho}^{n_{\rho}-1} z_{\rho+1}^{n_{\rho+1}} \cdots z_r^{n_r}, \end{aligned} \quad (6.3)$$

where, again, $1/(-1)!$ has to be interpreted as 0. The left-hand side equals

$$\frac{\partial \text{GF}_{F_{\boldsymbol{\eta}}^{(k)}}}{\partial z_{\rho}},$$

while the right-hand side is identified as

$$\frac{\partial \text{GF}_{F_{\boldsymbol{\eta}}}}{\partial z_{\rho}} \text{GF}_{F_{\boldsymbol{\eta}}^{(k-1)}},$$

whence the equations

$$\frac{\partial \text{GF}_{F_\eta^{(k)}}}{\partial z_\rho} = \frac{\partial \text{GF}_{F_\eta}}{\partial z_\rho} \text{GF}_{F_\eta^{(k-1)}}, \quad 1 \leq \rho \leq r. \quad (6.4)$$

Assuming inductively that

$$\text{GF}_{F_\eta^{(k-1)}} = \frac{1}{(k-1)!} (\text{GF}_{F_\eta})^{k-1},$$

we infer from (6.4) that

$$\text{GF}_{F_\eta^{(k)}} = \frac{1}{k!} (\text{GF}_{F_\eta})^k + C,$$

where C is independent of z_1, z_2, \dots, z_r . Making use of the facts that $\text{GF}_{F_\eta^{(k)}}(0, \dots, 0) = 0$ (since $k \geq 1$) and that $\text{GF}_{F_\eta}(0, \dots, 0) = 0$, we see that $C = 0$, which proves (6.1).

On the other hand, by Theorem 1, we know that

$$\text{GF}_F(z_1, z_2, \dots, z_r) = \exp(\text{GF}_{F_\eta}(z_1, z_2, \dots, z_r)),$$

or, equivalently,

$$\text{GF}_F(z_1, z_2, \dots, z_r) = \sum_{k \geq 0} \frac{1}{k!} (\text{GF}_{F_\eta}(z_1, z_2, \dots, z_r))^k. \quad (6.5)$$

If there were a non-empty intersection between $F_\eta^{(k_1)}[\Omega]$ and $F_\eta^{(k_2)}[\Omega]$, for some k_1, k_2 with $k_1 < k_2$ and some $\Omega \in \text{Ob}(\mathbf{Set}^r)$, then Proposition 2 would contradict (6.5) and (6.1). This proves the assertion of Proposition 3.

The proof of Theorem 4 is now easily completed. By definition of $\widetilde{\text{GF}}_F(z_1, \dots, z_r, y)$, we have

$$\widetilde{\text{GF}}_F(z_1, \dots, z_r, y) = \sum_{k \geq 0} y^k \text{GF}_{F_\eta^{(k)}}(z_1, \dots, z_r).$$

If we now substitute (6.1), then we immediately obtain (2.9). Identity (2.10) results from using Theorem 1 to express $\text{GF}_{F_\eta}(z_1, \dots, z_r)$ in terms of $\text{GF}_F(z_1, \dots, z_r)$ and substituting the result in (2.9). \square

7. ILLUSTRATIONS, I: THREE EXAMPLES

We give here three illustrations for the application of our theory. In the first and second example below, bipartite graphs are considered. Example 1 is, in some sense, “standard,” since it addresses the case where the composition operator η consists in “putting objects together,” so that the combinatorial objects in our (in this case, 2-sort) species are sets of indecomposable objects, a situation which is well covered by classical species theory. In Example 2, however, the composition operator η is different, “non-standard,” so that classical species theory does not apply, but our (extension of species) theory does. On the other hand, we shall see in Section 9 that this composition operator is pointwise associative and commutative (for the precise definition see Section 9), and that this family of composition operators is closely related to the classical operation of “putting objects together.” (See Theorem 22 for the precise statement.) Our last example in this section, Example 3, presents an example of a composition operator that is neither pointwise associative nor pointwise commutative. A particular aspect demonstrated by Examples 1 and 2 that we want to highlight is that composition operators need not be unique.

Example 1 (BIPARTITE GRAPHS I). Let the 2-sort species $F : \mathbf{Set}^2 \rightarrow \mathbf{Set}$ be defined by

$$F[\Omega] = F[(\Omega^{(1)}, \Omega^{(2)})] := 2^{\Omega^{(1)} \times \Omega^{(2)}}, \quad \Omega = (\Omega^{(1)}, \Omega^{(2)}) \in \text{Ob}(\mathbf{Set}^2).$$

Thus, $F[(\Omega^{(1)}, \Omega^{(2)})]$ can be considered as set of all bipartite graphs, where the set of “white” vertices is $\Omega^{(1)}$ and the set of “black” vertices is $\Omega^{(2)}$. For $(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_2)$, $b_1 \in F[\Omega_1]$ and $b_2 \in F[\Omega_2]$, put

$$\eta_{(\Omega_1, \Omega_2)}((b_1, b_2)) := b_1 \amalg b_2.$$

This means that $\eta_{(\Omega_1, \Omega_2)}$ merely forms the disjoint union of the bipartite graphs b_1 and b_2 . Then it is not difficult to see that η is a natural transformation satisfying (D1). Moreover, $F_\eta[\Omega]$ consists of the *connected* bipartite graphs with bipartition $\Omega = (\Omega^{(1)}, \Omega^{(2)})$.

For a weight, we choose $\Lambda = \mathbb{Z}[t]$ and

$$w_\Omega(b) := t^{|b|}, \quad b \in F[\Omega].$$

Again, it is not difficult to see that \mathbf{w} satisfies Axioms (W0)–(W2); that is, \mathbf{w} is a Λ -weight on (F, η) .

Theorem 1 then says that

$$\text{GF}_F(z_1, z_2) = \exp(\text{GF}_{F_\eta}(z_1, z_2)), \quad (7.1)$$

where

$$\text{GF}_F(z_1, z_2) = \sum_{n_1, n_2 \geq 0} \sum_{b \in F[(\{n_1\}, \{n_2\})]} t^{|b|} \frac{z_1^{n_1} z_2^{n_2}}{n_1! n_2!}$$

and

$$\text{GF}_{F_\eta}(z_1, z_2) = \sum_{n_1, n_2 \geq 0} \sum_{b \in F_\eta[(\{n_1\}, \{n_2\})]} t^{|b|} \frac{z_1^{n_1} z_2^{n_2}}{n_1! n_2!}.$$

However, by straightforward counting, one sees that

$$\text{GF}_F(z_1, z_2) = \sum_{n_1, n_2 \geq 0} (1+t)^{n_1 n_2} \frac{z_1^{n_1} z_2^{n_2}}{n_1! n_2!}.$$

From (7.1), it then follows that the generating function for connected bipartite graphs is given by

$$\text{GF}_{F_\eta}(z_1, z_2) = \log \left(\sum_{n_1, n_2 \geq 0} (1+t)^{n_1 n_2} \frac{z_1^{n_1} z_2^{n_2}}{n_1! n_2!} \right),$$

while (2.10) implies that

$$\begin{aligned} \widetilde{\text{GF}}_F(z_1, z_2) &:= \sum_{n_1, n_2 \geq 0} \sum_{b \in F[(\{n_1\}, \{n_2\})]} t^{|b|} y^{\#(\text{connected components of } b)} \frac{z_1^{n_1} z_2^{n_2}}{n_1! n_2!} \\ &= \left(\sum_{n_1, n_2 \geq 0} (1+t)^{n_1 n_2} \frac{z_1^{n_1} z_2^{n_2}}{n_1! n_2!} \right)^y. \end{aligned} \quad (7.2)$$

This example can be considered as a 2-dimensional analogue of the example in [10, Sec. 3] (with the first of the two composition operators considered there). The knowledgeable reader will recognise (7.2) as the exponential generating function for the Tutte polynomials of complete bipartite graphs (cf. e.g. [21, Eq. (3.10)]).

Example 2 (BIPARTITE GRAPHS II). Let $F : \mathbf{Set}^2 \rightarrow \mathbf{Set}$ be as in Example 1. Here, for $(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_2)$, where $\Omega_i = (\Omega_i^{(1)}, \Omega_i^{(2)})$, $i = 1, 2$, for $b_1 \in F[\Omega_1]$ and $b_2 \in F[\Omega_2]$, we put

$$\eta'_{(\Omega_1, \Omega_2)}((b_1, b_2)) := b_1 \amalg b_2 \amalg \left(\Omega_1^{(1)} \times \Omega_2^{(2)} \right) \amalg \left(\Omega_1^{(2)} \times \Omega_2^{(1)} \right).$$

The graph $\eta'_{(\Omega_1, \Omega_2)}((b_1, b_2))$ can be considered as a kind of bipartite completion of the disjoint union of b_1 and b_2 . Again, it is not difficult to see that η' is a natural transformation satisfying (D1). Moreover, $F_{\eta'}(\Omega)$ consists of the *complements* of connected bipartite graphs with bipartition $\Omega = (\Omega^{(1)}, \Omega^{(2)})$, where the complement b^c of a bipartite graph $b \in F[\Omega]$ is defined as $b^c := (\Omega^{(1)} \times \Omega^{(2)}) - b$.

If we now were to choose the weight of Example 1, then Axiom (W2) would be violated. Instead, with $\Lambda = \mathbb{Z}[t]$, we set

$$w'_\Omega(b) := t^{|\Omega^{(1)}| \cdot |\Omega^{(2)}| - |b|}, \quad b \in F[\Omega] = F[(\Omega^{(1)}, \Omega^{(2)})].$$

Then it is not difficult to see that w' does satisfy Axioms (W0)–(W2); that is, w' is a Λ -weight on (F, η') .

Theorem 1 then says that

$$\text{GF}_F(z_1, z_2) = \exp \left(\text{GF}_{F_{\eta'}}(z_1, z_2) \right), \quad (7.3)$$

where

$$\text{GF}_F(z_1, z_2) = \sum_{n_1, n_2 \geq 0} \sum_{b \in F[(n_1), (n_2)]} t^{n_1 n_2 - |b|} \frac{z_1^{n_1} z_2^{n_2}}{n_1! n_2!}$$

and

$$\text{GF}_{F_{\eta'}}(z_1, z_2) = \sum_{n_1, n_2 \geq 0} \sum_{b \in F_{\eta'}[(n_1), (n_2)]} t^{n_1 n_2 - |b|} \frac{z_1^{n_1} z_2^{n_2}}{n_1! n_2!}.$$

Again, by straightforward counting, one sees that

$$\text{GF}_F(z_1, z_2) = \sum_{n_1, n_2 \geq 0} (1+t)^{n_1 n_2} \frac{z_1^{n_1} z_2^{n_2}}{n_1! n_2!},$$

and we obtain the formulae

$$\text{GF}_{F_{\eta'}}(z_1, z_2) = \log \left(\sum_{n_1, n_2 \geq 0} (1+t)^{n_1 n_2} \frac{z_1^{n_1} z_2^{n_2}}{n_1! n_2!} \right)$$

and

$$\begin{aligned} \widetilde{\text{GF}}_F(z_1, z_2) &:= \sum_{n_1, n_2 \geq 0} \sum_{b \in F[(n_1), (n_2)]} t^{n_1 n_2 - |b|} y^{\#(\text{connected components of } b^c)} \frac{z_1^{n_1} z_2^{n_2}}{n_1! n_2!} \\ &= \left(\sum_{n_1, n_2 \geq 0} (1+t)^{n_1 n_2} \frac{z_1^{n_1} z_2^{n_2}}{n_1! n_2!} \right)^y. \end{aligned} \quad (7.4)$$

This example can be viewed as a 2-dimensional analogue of the example in [10, Sec. 3] (with the second of the two composition operators considered there).

The alert reader will have noticed that the η' -maps could have been alternatively defined by

$$\eta'_{(\Omega_1, \Omega_2)}((b_1, b_2)) := (b_1^c \amalg b_2^c)^c, \quad (7.5)$$

where the complements have to be taken in the appropriate complete bipartite graphs. This construction will be generalised in Section 9.

Example 3 (BINARY FUNCTIONS). Let the (1-sort) species $F : \mathbf{Set} \rightarrow \mathbf{Set}$ be defined by

$$F[\Omega] := \{0, 1\}^\Omega, \quad \Omega \in \text{Ob}(\mathbf{Set}).$$

For $(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_1)$, $f_1 \in F[\Omega_1]$, and $f_2 \in F[\Omega_2]$, put

$$(\eta_{(\Omega_1, \Omega_2)}((f_1, f_2)))(\omega) := \begin{cases} f_1(\omega), & \text{if } \omega \in \Omega_1, \\ 1 - f_2(\omega), & \text{if } \omega \in \Omega_2. \end{cases}$$

Then it is easy to see that η is a natural transformation satisfying (D1). Moreover,

$$F_\eta[\Omega] = \begin{cases} \{0_\Omega, 1_\Omega\}, & \text{if } |\Omega| = 1, \\ \{\}, & \text{otherwise,} \end{cases}$$

where 0_Ω and 1_Ω are the constant functions on Ω taking the value 0 and 1, respectively. We note that, in contrast to Examples 1 and 2, the η -maps of the present example are *pointwise non-associative and non-commutative* (cf. Section 9); to be precise, in general we have

$$\eta_{(\Omega_1 \amalg \Omega_2, \Omega_3)}\left(\left(\eta_{(\Omega_1, \Omega_2)}((f_1, f_2)), f_3\right)\right) \neq \eta_{(\Omega_1, \Omega_2 \amalg \Omega_3)}\left(\left(f_1, \eta_{(\Omega_2, \Omega_3)}((f_2, f_3))\right)\right)$$

and

$$\eta_{(\Omega_1, \Omega_2)}((f_1, f_2)) \neq \eta_{(\Omega_2, \Omega_1)}((f_2, f_1)).$$

For the sake of completeness, we remark that, choosing the trivial weighting

$$w_\Omega(f) := 1, \quad f \in F[\Omega],$$

Theorem 1 yields the trivial identity

$$\text{GF}_F(z) = \sum_{n \geq 0} 2^n \frac{z^n}{n!} = \exp(\text{GF}_{F_\eta}(z)) = \exp(2z).$$

The construction of this example can also be generalised to produce many more (multi-sort) species with pointwise non-associative and non-commutative composition operator, see Theorem 23 in Section 9.

8. ILLUSTRATIONS, II: MAGIC SQUARES

The purpose of this section is to illustrate the increased flexibility of our present multivariate setting. We show that a number of generating function identities for combinatorial matrices found scattered throughout the literature can be uniformly explained, and generalised, in the context of our theory.

By a *combinatorial matrix* on $\Omega = (\Omega^{(1)}, \Omega^{(2)}) \in \text{Ob}(\mathbf{Set}^2)$ we shall mean any map

$$m : \Omega^{(1)} \times \Omega^{(2)} \rightarrow \mathbb{N}_0.$$

The pair of sets Ω is called the *support* of m . Let m_1, m_2 be two combinatorial matrices with supports $\Omega_1 = (\Omega_1^{(1)}, \Omega_1^{(2)})$ and $\Omega_2 = (\Omega_2^{(1)}, \Omega_2^{(2)})$, respectively, and suppose that $\Omega_1 \cap \Omega_2 = \emptyset$. Then we define their *direct sum* $m = m_1 \oplus m_2$ to be the combinatorial matrix with support $\Omega := \Omega_1 \amalg \Omega_2$ given by

$$m(\omega_1, \omega_2) := \begin{cases} m_1(\omega_1, \omega_2), & (\omega_1, \omega_2) \in \Omega_1^{(1)} \times \Omega_1^{(2)}, \\ m_2(\omega_1, \omega_2), & (\omega_1, \omega_2) \in \Omega_2^{(1)} \times \Omega_2^{(2)}, \\ 0, & \text{otherwise.} \end{cases}$$

A combinatorial matrix m on Ω is termed *s-magic*,⁶ s a positive integer, if

$$\sum_{\omega_2 \in \Omega^{(2)}} m(\omega_1, \omega_2) = s, \quad \omega_1 \in \Omega^{(1)},$$

and

$$\sum_{\omega_1 \in \Omega^{(1)}} m(\omega_1, \omega_2) = s, \quad \omega_2 \in \Omega^{(2)}.$$

Computing the sum of entries, we find that an *s-magic* matrix is necessarily square, $|\Omega^{(1)}| = |\Omega^{(2)}|$. The enumeration of *s-magic* squares has a long history, going back to MacMahon [18, §404–419]. A good account of the enumerative theory of magic squares can be found in [26, Sec. 4.6], with many pointers to further literature. For more recent work, see for instance [4, 9].

For $s \in \mathbb{N}$ and $\Omega \in \text{Ob}(\mathbf{Set}^2)$, denote by $F_s(\Omega)$ the set of all *s-magic* matrices on Ω , and by $\bar{F}_s(\Omega)$ the set of those *s-magic* matrices on Ω which do not contain s as an entry. We thus have mappings

$$F_s, \bar{F}_s : \text{Ob}(\mathbf{Set}^2) \rightarrow \text{Ob}(\mathbf{Set}),$$

which we turn into functors $F_s, \bar{F}_s : \mathbf{Set}^2 \rightarrow \mathbf{Set}$ by assigning to a morphism

$$\mathbf{f} = (f_1, f_2) : \Omega \rightarrow \tilde{\Omega}$$

in \mathbf{Set}^2 the map (denoted $F_s[\mathbf{f}]$ respectively $\bar{F}_s[\mathbf{f}]$) sending a combinatorial matrix m in the respective domain to $m \circ (f_1^{-1} \times f_2^{-1})$. Moreover, given s and a finite set Ω , let $F_s^*(\Omega)$ be the set of *symmetric s-magic* matrices on $\Omega = (\Omega, \Omega)$; that is, combinatorial matrices satisfying

$$m(\omega_1, \omega_2) = m(\omega_2, \omega_1), \quad (\omega_1, \omega_2) \in \Omega^2;$$

and denote by $\bar{F}_s^*(\Omega)$ the subset of $F_s^*(\Omega)$ consisting of those matrices with no entry equal to s . Just as above, the maps

$$F_s^*, \bar{F}_s^* : \text{Ob}(\mathbf{Set}) \rightarrow \text{Ob}(\mathbf{Set})$$

become functors $F_s^*, \bar{F}_s^* : \mathbf{Set} \rightarrow \mathbf{Set}$ by assigning to a morphism $f : \Omega \rightarrow \tilde{\Omega}$ in \mathbf{Set} the map sending a combinatorial matrix m in the respective domain to $m \circ (f^{-1} \times f^{-1})$.

Next, given $s \in \mathbb{N}$, a pair $(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_2)$, and a pair $(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_1)$, the direct sum construction provides us with injective maps

$$\begin{aligned} (\eta_s)_{(\Omega_1, \Omega_2)} &: F_s(\Omega_1) \times F_s(\Omega_2) \rightarrow F_s(\Omega_1 \amalg \Omega_2), \\ (\bar{\eta}_s)_{(\Omega_1, \Omega_2)} &: \bar{F}_s(\Omega_1) \times \bar{F}_s(\Omega_2) \rightarrow \bar{F}_s(\Omega_1 \amalg \Omega_2), \\ (\eta_s^*)_{(\Omega_1, \Omega_2)} &: F_s^*(\Omega_1) \times F_s^*(\Omega_2) \rightarrow F_s^*(\Omega_1 \amalg \Omega_2), \\ (\bar{\eta}_s^*)_{(\Omega_1, \Omega_2)} &: \bar{F}_s^*(\Omega_1) \times \bar{F}_s^*(\Omega_2) \rightarrow \bar{F}_s^*(\Omega_1 \amalg \Omega_2). \end{aligned}$$

A certain amount of checking is required in order to convince oneself that these definitions fit into the framework of Theorems 1 and 4. The next lemma states the corresponding result. We leave its proof, which essentially amounts to a routine verification, to the reader.

Lemma 19. (i) *For each $s \in \mathbb{N}$, the collection of maps*

$$\boldsymbol{\eta}_s = \left((\eta_s)_{(\Omega_1, \Omega_2)} \right)_{(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_2)}$$

⁶Strictly speaking, the correct term here would be “*s*-semi-magic,” since we do not require diagonals to sum up to s as well, see e.g. [4]. However, we prefer the term “*s*-magic” for the sake of brevity.

is a natural transformation from the functor $F_s \times F_s$ to the functor $F_s \circ \Pi$. Analogous statements hold for the functors $\bar{F}_s, F_s^*, \bar{F}_s^*$, and the families of maps

$$\begin{aligned}\bar{\eta}_s &= \left((\bar{\eta}_s)_{(\Omega_1, \Omega_2)} \right)_{(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_2)}, \\ \eta_s^* &= \left((\eta_s^*)_{(\Omega_1, \Omega_2)} \right)_{(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_1)}, \\ \bar{\eta}_s^* &= \left((\bar{\eta}_s^*)_{(\Omega_1, \Omega_2)} \right)_{(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_1)}.\end{aligned}$$

(ii) For each s , the pair (F_s, η_s) satisfies Axiom (D1), an analogous statement holding for each of the other pairs $(\bar{F}_s, \bar{\eta}_s)$, (F_s^*, η_s^*) , and $(\bar{F}_s^*, \bar{\eta}_s^*)$.

It follows from Lemma 19 and Theorem 4, that Equations (2.9) and (2.10) hold for each of the pairs (F_s, η_s) , $(\bar{F}_s, \bar{\eta}_s)$, (F_s^*, η_s^*) , and $(\bar{F}_s^*, \bar{\eta}_s^*)$; in particular, we find that

$$\widetilde{\text{GF}}_{F_s}(z_1, z_2, y) = \exp(y \text{GF}_{(F_s)\eta_s}(z_1, z_2)), \quad (8.1)$$

$$\widetilde{\text{GF}}_{\bar{F}_s}(z_1, z_2, y) = \exp(y \text{GF}_{(\bar{F}_s)\bar{\eta}_s}(z_1, z_2)), \quad (8.2)$$

$$\widetilde{\text{GF}}_{F_s^*}(z, y) = \exp(y \text{GF}_{(F_s^*)\eta_s^*}(z)), \quad (8.3)$$

$$\widetilde{\text{GF}}_{\bar{F}_s^*}(z, y) = \exp(y \text{GF}_{(\bar{F}_s^*)\bar{\eta}_s^*}(z)). \quad (8.4)$$

Note that in these identities the variable y keeps track of the number of indecomposable matrices into which the matrices which are counted by the respective generating functions on the left-hand sides can be decomposed. Clearly, the generating functions occurring in (8.1) and (8.2) can be viewed as formal power series in $z_1 z_2$ and y ; that is, $z_1 z_2$ could be replaced by a single variable. However, we prefer to keep z_1 and z_2 separate, since this is more in line with our general theory.

We note certain dependencies among the series $\widetilde{\text{GF}}_{F_s}, \widetilde{\text{GF}}_{\bar{F}_s}, \widetilde{\text{GF}}_{F_s^*}, \widetilde{\text{GF}}_{\bar{F}_s^*}$; for instance, we observe that an indecomposable s -magic matrix on $([n_1], [n_2])$ cannot contain an entry equal to s , unless $n_1 = n_2 = 1$. It follows that

$$|(F_s)_{\eta_s}([n_1], [n_2])| = \begin{cases} 1 + |(\bar{F}_s)_{\bar{\eta}_s}([1], [1])|, & n_1 = n_2 = 1, \\ |(\bar{F}_s)_{\bar{\eta}_s}([n_1], [n_2])|, & \text{otherwise,} \end{cases}$$

and hence, by Equations (8.1) and (8.2),

$$\widetilde{\text{GF}}_{\bar{F}_s}(z_1, z_2, y) = e^{-z_1 z_2 y} \widetilde{\text{GF}}_{F_s}(z_1, z_2, y). \quad (8.5)$$

Similarly, we have

$$\widetilde{\text{GF}}_{\bar{F}_s^*}(z, y) = e^{-y(z+z^2/2)} \widetilde{\text{GF}}_{F_s^*}(z, y). \quad (8.6)$$

Indeed, for $n = 1, 2$, there exist indecomposable symmetric s -magic matrices on $([n], [n])$ containing an entry s :

$$(s) \text{ and } \begin{pmatrix} 0 & s \\ s & 0 \end{pmatrix}.$$

Now let m be a symmetric s -magic matrix on $([n], [n])$ with $n \geq 3$, and suppose that m contains an entry equal to s in position (i, j) . Then, if $i = j$, we have $m = (s) \oplus m'$, where m' has support $([n] - \{i\}, [n] - \{i\})$. If, on the other hand, $i \neq j$, then, by symmetry, m also contains s in position (j, i) , and we find that m splits as

$$m = \begin{pmatrix} 0 & s \\ s & 0 \end{pmatrix} \oplus m',$$

where m' has support $([n] - \{i, j\}, [n] - \{i, j\})$, and is non-empty since $n \geq 3$. Thus, in both cases, m is in fact decomposable. Hence,

$$|(F_s^*)_{\eta_s^*}([n])| = \begin{cases} 1 + |(\bar{F}_s^*)_{\bar{\eta}_s^*}([n])|, & n = 1, 2, \\ |(\bar{F}_s^*)_{\bar{\eta}_s^*}([n])|, & \text{otherwise,} \end{cases}$$

and (8.6) follows from Equations (8.3) and (8.4).

The enumeration can be done exactly if $s = 2$. For, according to Birkhoff's Theorem (cf. [6] or [2, Corollary 8.40]), a 2-magic matrix m is the sum of two permutation matrices, say p_1 and p_2 . If m is indecomposable, then the pair $\{p_1, p_2\}$ is uniquely determined. Premultiplying by p_1^{-1} , we obtain a situation where p_1 is the identity; indecomposability forces p_2 to be the permutation matrix corresponding to a cyclic permutation. So there are $n!(n-1)!$ choices for (p_1, p_2) , and half this many choices for m (assuming, as we may, that $n > 1$). Note that this formula gives half the correct number for $n = 1$. So we have

$$|(F_2)_{\eta_2}([n_1], [n_2])| = \begin{cases} 1, & n_1 = n_2 = n = 1, \\ \frac{n!(n-1)!}{2}, & n_1 = n_2 = n > 1, \\ 0, & n_1 \neq n_2, \end{cases}$$

that is,

$$\text{GF}_{(F_2)_{\eta_2}}(z_1, z_2) = \frac{1}{2}z_1z_2 - \frac{1}{2}\log(1 - z_1z_2),$$

and therefore

$$\widetilde{\text{GF}}_{F_2}(z_1, z_2, y) = (1 - z_1z_2)^{-y/2} e^{z_1z_2y/2} \quad (8.7)$$

by Equation (8.1). Also,

$$\widetilde{\text{GF}}_{\bar{F}_2}(z_1, z_2, y) = (1 - z_1z_2)^{-y/2} e^{-z_1z_2y/2}, \quad (8.8)$$

making use of Equation (8.5) and the last result. Special cases of Identities (8.7) and (8.8) appear in [3, Sections 8.1 and 8.3] (see also [25, Eqs. (23) and (24) in Example 6.11]). For $s > 2$, enumeration is more difficult; see Stanley's paper [24] and Comtet [8, pp. 124–125] for comments in this direction; also Goulden and Jackson [14, Sections 3.4 and 3.5] for some variations on this counting problem.⁷

For symmetric matrices, it is again possible to count the indecomposables with $s = 2$. For $n > 2$, such a matrix can be represented as a graph in which every vertex has degree 2; loops are permitted, but contribute only one to the degree of a vertex. Indecomposability of the matrix is reflected in connectedness of the graph. So the graphs we must consider are paths (with a loop at each end) and cycles; and, for $n > 2$, their number is $\frac{1}{2}(n! + (n-1)!)$. Including the cases where $n \leq 2$, we obtain

$$\text{GF}_{(F_2^*)_{\eta_2^*}}(z) = \frac{z^2}{4} + \frac{z}{2(1-z)} - \frac{1}{2}\log(1-z),$$

and, hence

$$\widetilde{\text{GF}}_{F_2^*}(z, y) = (1-z)^{-y/2} \exp\left(\frac{yz^2}{4} + \frac{yz}{2(1-z)}\right), \quad (8.9)$$

as well as

$$\widetilde{\text{GF}}_{\bar{F}_2^*}(z, y) = (1-z)^{-y/2} \exp\left(-\frac{yz^2}{4} - yz + \frac{yz}{2(1-z)}\right). \quad (8.10)$$

Identity (8.9) generalises [15, Eq. (6.3)], whereas (8.10) generalises [15, Eq. (6.4)].

⁷Note however, that the formula given in [8] for $s = 3$ is erroneous.

9. CHARACTERISATION OF POINTWISE ASSOCIATIVE AND COMMUTATIVE
COMPOSITION OPERATORS

In this final section, we address the question how far the composition operator η of our theory can be from the standard operation for the species of sets of combinatorial structures given by forming the disjoint union, of which Example 1 in Section 7 is a prototypical example. While we do not have an answer in general, we are able to completely resolve the case when η is pointwise associative and commutative. Here, given an r -sort species $F : \mathbf{Set}^r \rightarrow \mathbf{Set}$ with composition operator η , we say that η is *pointwise associative* if, for all pairwise disjoint $\Omega_1, \Omega_2, \Omega_3 \in \text{Ob}(\mathbf{Set}^r)$, and all elements $x_1 \in F[\Omega_1]$, $x_2 \in F[\Omega_2]$, $x_3 \in F[\Omega_3]$, we have

$$\eta_{(\Omega_1 \amalg \Omega_2, \Omega_3)} \left(\left(\eta_{(\Omega_1, \Omega_2)}((x_1, x_2)), x_3 \right) \right) = \eta_{(\Omega_1, \Omega_2 \amalg \Omega_3)} \left(\left(x_1, \eta_{(\Omega_2, \Omega_3)}((x_2, x_3)) \right) \right),$$

and we say that η is *pointwise commutative* if, for all $(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_r)$, and all elements $x_1 \in F[\Omega_1]$, $x_2 \in F[\Omega_2]$, we have

$$\eta_{(\Omega_1, \Omega_2)}((x_1, x_2)) = \eta_{(\Omega_2, \Omega_1)}((x_2, x_1)).$$

The η -maps in Examples 1 and 2 are instances of pointwise associative and commutative composition operators, while the composition operator in Example 3 is neither pointwise associative nor pointwise commutative. The notion of pointwise commutativity and associativity as defined above should not be confused with the commutativity and the 3-associativity proved in Lemmas 6 and 7, respectively, which are (in general) strictly weaker assertions.

Before we are able to make the above statement precise, we need two preparatory results. Recall that a *species isomorphism* between two r -sort species F_1 and F_2 is a collection of maps $\varphi = (\varphi_\Omega)_{\Omega \in \text{Ob}(\mathbf{Set}^r)}$, where, for each $\Omega \in \text{Ob}(\mathbf{Set}^r)$,

$$\varphi_\Omega : F_1[\Omega] \rightarrow F_2[\Omega]$$

is a bijection, with the property that, for every morphism $f : \Omega \rightarrow \tilde{\Omega}$ in the category \mathbf{Set}^r , the diagram

$$\begin{array}{ccc} F_1[\Omega] & \xrightarrow{F_1[f]} & F_1[\tilde{\Omega}] \\ \varphi_\Omega \downarrow & & \downarrow \varphi_{\tilde{\Omega}} \\ F_2[\Omega] & \xrightarrow{F_2[f]} & F_2[\tilde{\Omega}] \end{array} \quad (9.1)$$

commutes. If F_1 carries a weak Λ_1 -weight \mathbf{w}_1 and F_2 carries a weak Λ_2 -weight \mathbf{w}_2 , where, by “weak,” we mean that \mathbf{w}_1 and \mathbf{w}_2 satisfy Axioms (W0) and (W1), but not necessarily (W2) (cf. Section 2), then an isomorphism $\varphi : F_1 \rightarrow F_2$ is called *weight-preserving*, if there exists a ring homomorphism $\lambda : \Lambda_1 \rightarrow \Lambda_2$ such that the diagram

$$\begin{array}{ccc} F_1[\Omega] & \xrightarrow{(w_1)_\Omega} & \Lambda_1 \\ \varphi_\Omega \downarrow & & \downarrow \lambda \\ F_2[\Omega] & \xrightarrow{(w_2)_\Omega} & \Lambda_2 \end{array} \quad (9.2)$$

commutes.

The lemma below tells us that, if F_1 and F_2 are two isomorphic r -sort species, where F_1 is decomposable with composition operator η_1 , then η_1 can be lifted to a composition operator for F_2 , demonstrating that F_2 is decomposable as well.

Lemma 20. *Let F_1 and F_2 be two isomorphic r -sort species, where F_1 is decomposable with composition operator η_1 . Furthermore, let φ be an isomorphism between F_1 and F_2 . Then F_2 is decomposable, and the family of maps $\eta_2 = ((\eta_2)_{(\Omega_1, \Omega_2)})_{(\Omega_1, \Omega_2) \in \text{Ob}(\mathfrak{D}_r)}$ defined by*

$$(\eta_2)_{(\Omega_1, \Omega_2)}(x_1, x_2) := \varphi_{\Omega_1 \amalg \Omega_2} \left((\eta_1)_{(\Omega_1, \Omega_2)} \left((\varphi_{\Omega_1}^{-1}(x_1), \varphi_{\Omega_2}^{-1}(x_2)) \right) \right),$$

$$x_1 \in F_2[\Omega_1], \quad x_2 \in F_2[\Omega_2],$$

is a composition operator for F_2 .

Proof. We have to show that η_2 is a natural transformation from $F_2 \times F_2$ to $F_2 \circ \amalg$, and that the pair (F_2, η_2) satisfies Axiom (D1). The former follows immediately from the corresponding property for (F_1, η_1) and the naturality condition (9.1). In order to verify (D1), we start with the left-hand side of (2.2) for the pair (F_2, η_2) , suppressing the indices of η_1, η_2, φ for better readability:

$$\begin{aligned} & \eta_2(F_2[\Omega_1] \times F_2[\Omega_2]) \cap \eta_2(F_2[\tilde{\Omega}_1] \times F_2[\tilde{\Omega}_2]) \\ &= \varphi \left(\eta_1(\varphi^{-1}(F_2[\Omega_1]) \times \varphi^{-1}(F_2[\Omega_2])) \right) \cap \varphi \left(\eta_1(\varphi^{-1}(F_2[\tilde{\Omega}_1]) \times \varphi^{-1}(F_2[\tilde{\Omega}_2])) \right) \\ &= \varphi \left(\eta_1(F_1[\Omega_1] \times F_1[\Omega_2]) \right) \cap \varphi \left(\eta_1(F_1[\tilde{\Omega}_1] \times F_1[\tilde{\Omega}_2]) \right) \\ &= \varphi \left(\eta_1(F_1[\Omega_1] \times F_1[\Omega_2]) \cap \eta_1(F_1[\tilde{\Omega}_1] \times F_1[\tilde{\Omega}_2]) \right). \end{aligned}$$

Here we have used the injectivity of φ to obtain the last line. Now we substitute the right-hand side of (2.2) for the pair (F_1, η_1) , to obtain

$$\begin{aligned} & \eta_2(F_2[\Omega_1] \times F_2[\Omega_2]) \cap \eta_2(F_2[\tilde{\Omega}_1] \times F_2[\tilde{\Omega}_2]) \\ &= \varphi \left(\eta_1(\eta_1(F_1[\Omega_{11}] \times F_1[\Omega_{12}]) \times \eta_1(F_1[\Omega_{21}] \times F_1[\Omega_{22}])) \right), \end{aligned}$$

where $\Omega_{ij} := \Omega_i \cap \tilde{\Omega}_j$ for $i, j \in \{1, 2\}$. Using $F_1[\Omega_{ij}] = \varphi^{-1}(F_2[\Omega_{ij}])$ at each possible place, and inserting $\text{id} = \varphi^{-1} \circ \varphi$ at two places, we arrive at

$$\begin{aligned} & \eta_2(F_2[\Omega_1] \times F_2[\Omega_2]) \cap \eta_2(F_2[\tilde{\Omega}_1] \times F_2[\tilde{\Omega}_2]) \\ &= \varphi \left(\eta_1(\varphi^{-1}(\varphi(\eta_1(\varphi^{-1}(F_2[\Omega_{11}]) \times \varphi^{-1}(F_2[\Omega_{12}])))) \right. \\ & \quad \left. \times \varphi^{-1}(\varphi(\eta_1(\varphi^{-1}(F_2[\Omega_{21}]) \times \varphi^{-1}(F_2[\Omega_{22}])))) \right) \\ &= \eta_2(\eta_2(F_2[\Omega_{11}] \times F_2[\Omega_{12}]) \times \eta_2(F_2[\Omega_{21}] \times F_2[\Omega_{22}])), \end{aligned}$$

which is exactly (2.2) for the pair (F_2, η_2) . \square

The second preparatory result, Proposition 21 below, states that, given a decomposable r -sort species F with pointwise associative and commutative composition operator η , F is isomorphic to $E(F_\eta)$, where $E(F_\eta)$ denotes the species of sets of F_η -structures (cf. [7, p. 8] for the definition of the species of sets, E , and [7, p. 41] for the definition of composition of species). In rigorous terms, for $\Omega \in \text{Ob}(\mathbf{Set}^r)$, the set $E(F_\eta)[\Omega]$ can be defined by

$$E(F_\eta)[\Omega] := \left\{ \left\{ (x_1, \Omega_1), \dots, (x_k, \Omega_k) \right\} : x_i \in F_\eta[\Omega_i], \quad i = 1, \dots, k, \right. \\ \left. \text{for some } k \in \mathbb{N}_0 \text{ and } \Omega_1 \amalg \dots \amalg \Omega_k = \Omega, \text{ all } \Omega_i \text{'s being non-empty} \right\},$$

with the obvious notion of induced morphisms. If F carries a weak Λ -weight \mathbf{w} , then \mathbf{w} can be lifted to a weak Λ -weight of $E(F_\eta)$ by setting

$$w_\Omega \left(\{ (x_1, \Omega_1), \dots, (x_k, \Omega_k) \} \right) := w_{\Omega_1}(x_1) \cdots w_{\Omega_k}(x_k).$$

Proposition 21. *Let F be a decomposable weighted r -sort species with composition operator η , where η is pointwise associative and commutative. Then there exists a weight-preserving isomorphism between F and $E(F_\eta)$.*

Proof. The starting point is the combination of Lemmas 12 and 13. It says that, for each non-empty $\Omega \in \text{Ob}(\text{Set}^r)$ and every choice of base point $(\omega, \rho) \in \Omega$, we have

$$F[\Omega] = \coprod_{\substack{\Omega_1 \in \text{Ob}(\text{Set}^r) \\ (\omega, \rho) \in \Omega_1 \subseteq \Omega}} \eta_{(\Omega_1, \Omega - \Omega_1)}(F_\eta[\Omega_1] \times F[\Omega - \Omega_1]). \quad (9.3)$$

We are now going to construct bijective maps $\psi_\Omega : F[\Omega] \rightarrow E(F_\eta)[\Omega]$ by induction on $\|\Omega\|$, where $\|\cdot\|$ has been defined in (2.5). For $\Omega = \emptyset$, we have $|F[\Omega]| = |E(F_\eta)[\Omega]| = 1$ by Lemma 5 respectively the definition of $E(F_\eta)$, whence the construction of ψ_\emptyset is trivial. Henceforth, we shall suppose that $\|\Omega\| \geq 1$, and we assume that we have constructed maps $\psi_{\tilde{\Omega}}$ for all $\tilde{\Omega} \in \text{Ob}(\text{Set}^r)$ with $\|\tilde{\Omega}\| < N$.

Now let $\|\Omega\| = N$. Choose a base point $(\omega, \rho) \in \Omega$, and let $x \in F[\Omega]$. By (9.3), there is a unique Ω_1 such that $x \in \eta_{(\Omega_1, \Omega - \Omega_1)}(F_\eta[\Omega_1] \times F[\Omega - \Omega_1])$ and $(\omega, \rho) \in \Omega_1$. Let (y_1, x_1) be the uniquely determined pair with $y_1 \in F_\eta[\Omega_1]$ and $x_1 \in F[\Omega - \Omega_1]$, such that

$$(y_1, x_1) := \eta_{(\Omega_1, \Omega - \Omega_1)}^{-1}(x). \quad (9.4)$$

By the inductive hypothesis, there exist uniquely determined elements y_2, \dots, y_k , for some $k \in \mathbb{N}$ and $y_i \in F_\eta[\Omega_i]$, $i = 2, \dots, k$, with $\Omega_2 \amalg \cdots \amalg \Omega_k = \Omega - \Omega_1$, such that

$$\psi_{\Omega - \Omega_1}(x_1) = \{ (y_2, \Omega_2), \dots, (y_k, \Omega_k) \}. \quad (9.5)$$

Define

$$\psi_\Omega(x) := \{ (y_1, \Omega_1), (y_2, \Omega_2), \dots, (y_k, \Omega_k) \}.$$

We claim that this yields a well-defined bijection $\psi_\Omega : F[\Omega] \rightarrow E(F_\eta)[\Omega]$. What needs to be checked here first of all is that different choices of base points would always lead to the same result. So, let us suppose, that, by choosing a different base point, we would have obtained

$$\bar{\psi}_\Omega(x) := \{ (\bar{y}_1, \bar{\Omega}_1), (\bar{y}_2, \bar{\Omega}_2), \dots, (\bar{y}_l, \bar{\Omega}_l) \},$$

for some l , instead. Since we must have

$$\Omega = \Omega_1 \amalg \Omega_2 \amalg \cdots \amalg \Omega_k = \bar{\Omega}_1 \amalg \bar{\Omega}_2 \amalg \cdots \amalg \bar{\Omega}_l,$$

there is a j such that $(\omega, \rho) \in \bar{\Omega}_j$. By our inductive construction via (9.4) and (9.5), we have

$$x \in \eta \left(F_\eta[\bar{\Omega}_1] \times \eta \left(F_\eta[\bar{\Omega}_2] \times \cdots \times \left(F_\eta[\bar{\Omega}_{l-1}] \times F_\eta[\bar{\Omega}_l] \right) \cdots \right) \right).$$

By Lemma 15 (m -permutability for (F_η, η)), this is equivalent to saying that

$$x \in \eta \left(F_\eta[\bar{\Omega}_j] \times \eta \left(F_\eta[\bar{\Omega}_{\sigma(2)}] \times \cdots \times \left(F_\eta[\bar{\Omega}_{\sigma(l-1)}] \times F_\eta[\bar{\Omega}_{\sigma(l)}] \right) \cdots \right) \right), \quad (9.6)$$

where $\sigma(2), \dots, \sigma(l-1), \sigma(l)$ is some permutation of $\{1, \dots, j-1, j+1, \dots, l\}$. If $\bar{\Omega}_j \neq \Omega_1$, then (9.4) and (9.6) would contradict the disjointness in (9.3). Hence, we must have $\bar{\Omega}_j = \Omega_1$, and, by our assumption that η be pointwise associative and

commutative, we even must have $\bar{y}_j = y_1$. The inductive hypothesis applied to $\Omega - \Omega_1$ then guarantees that, moreover,

$$\{(y_2, \Omega_1), \dots, (y_k, \Omega_k)\} = \{(\bar{y}_1, \Omega_1), \dots, (\bar{y}_{j-1}, \bar{\Omega}_{j-1}), (\bar{y}_{j+1}, \bar{\Omega}_{j+1}), \dots, (\bar{y}_l, \Omega_l)\}.$$

This proves that ψ_Ω is indeed well-defined.

The facts that each map ψ_Ω is a bijection, and that the family $\psi = (\psi_\Omega)_{\Omega \in \text{Ob}(\text{Set}^r)}$ is an isomorphism between F and $E(F_\eta)$, are not hard to verify. The fact that ψ is weight-preserving is obvious from the definition of ψ and Axiom (W2) for (F, η, \mathbf{w}) . This completes the proof of the proposition. \square

If we combine Lemma 20 and Proposition 21, then we can say exactly how a decomposable r -sort species F with pointwise associative and commutative composition operator η arises from the composition of the species of sets with the species of F_η -structures (“components”). We should point out here that, clearly, a natural composition operator for $E(F_\eta)$ is given by

$$(y_1, y_2) \mapsto y_1 \amalg y_2, \quad y_1 \in E(F_\eta)[\Omega_1], \quad y_2 \in E(F_\eta)[\Omega_2], \quad \Omega_1 \amalg \Omega_2 = \Omega. \quad (9.7)$$

Theorem 22. *Let F be a decomposable r -sort species with composition operator η , where η is pointwise associative and commutative. Then, for all $(\Omega_1, \Omega_2) \in \text{Ob}(\mathcal{D}_r)$, the composition operator η can be expressed as follows:*

$$\eta_{(\Omega_1, \Omega_2)}(x_1, x_2) = \psi_{\Omega_1 \amalg \Omega_2}^{-1} \left(\psi_{\Omega_1}(x_1) \amalg \psi_{\Omega_2}(x_2) \right), \quad x_1 \in F[\Omega_1], \quad x_2 \in F[\Omega_2], \quad (9.8)$$

where ψ is the isomorphism between F and $E(F_\eta)$ constructed in the proof of Proposition 21.

Proof. One combines Lemma 20 with Proposition 21, where the role of the isomorphism φ in Lemma 20 is played by the family of maps ψ^{-1} constructed in the proof of Proposition 21. \square

In summary, all decomposable r -sort species with pointwise associative and commutative composition operator can be constructed from $E(G)$, for some species G , equipped with the natural composition operator as given in (9.7) (in the case where $G = F_\eta$), by applying a lift in the sense of Lemma 20 via a species isomorphism. In order to see how Example 2 in Section 7 fits into the setting of Theorem 22, recall that the isomorphism between F and $E(F_\eta)$ in that example can be defined by mapping the bipartite graph $b \in F[\Omega]$ to its complement b^c , identifying the connected components (in the classical sense of graph theory) of b^c , and forming the set of complements of these connected components (restricted to the set of vertices which a component involves). If this isomorphism is inserted in (9.8), the result is (7.5).

We conclude our paper by pointing out that the construction in Theorem 22 can be “twisted” to produce pointwise non-associative and non-commutative composition operators as well.

Theorem 23. *Let G be a weighted r -sort species, and let $\mathbf{g} : G \rightarrow G$ be a weight-preserving isomorphism. We extend \mathbf{g} to $E(G)$ by setting*

$$g_{\Omega_1 \amalg \dots \amalg \Omega_k} \left(\{(y_1, \Omega_1), \dots, (y_k, \Omega_k)\} \right) = \{(g_{\Omega_1}(y_1), \Omega_1), \dots, (g_{\Omega_k}(y_k), \Omega_k)\}.$$

Then the family $\eta = (\eta_{(\Omega_1, \Omega_2)})_{(\Omega_1, \Omega_2) \in \text{Ob}(\mathcal{D}_r)}$ of maps defined by

$$\eta_{(\Omega_1, \Omega_2)}(x_1, x_2) = x_1 \amalg g_{\Omega_2}(x_2), \quad x_1 \in E(G)[\Omega_1], \quad x_2 \in E(G)[\Omega_2],$$

where $(\Omega_1, \Omega_2) \in \text{Ob}(\mathcal{D}_r)$, is a composition operator for the weighted species $E(G)$.

It is obvious from the definition that the composition operator η of Theorem 23 will, in general, be neither pointwise associative nor pointwise commutative. Example 3 in Section 7 provides a typical example of the above construction, with G given by

$$G[\Omega] = \begin{cases} \{0_\Omega, 1_\Omega\}, & \text{if } |\Omega| = 1, \\ \{\}, & \text{otherwise,} \end{cases}$$

where 0_Ω and 1_Ω are the constant functions on Ω taking the value 0 and 1, respectively, and where the isomorphism g is given by $g_\Omega(0_\Omega) = 1_\Omega$ and $g_\Omega(1_\Omega) = 0_\Omega$ for $|\Omega| = 1$. However, we expect that there are many composition operators η not obtainable in this way.

REFERENCES

- [1] M. Ahmed, J. De Loera and R. Hemmecke, Polyhedral Cones of Magic Cubes and Squares, in: Discrete and computational geometry. The Goodman–Pollack Festschrift (B. Aronov, S. Basu, J. Pach and M. Sharir, eds.), Springer–Verlag, Berlin, Algorithms Comb. 25, 2003, pp. 25–41.
- [2] M. Aigner, *Combinatorial Theory*, Springer–Verlag, Berlin, 1979.
- [3] H. Anand, V. C. Dumir, and H. Gupta, A combinatorial distribution problem, *Duke Math. J.* **33** (1966), 757–769.
- [4] M. Beck, M. Cohen, J. Cuomo and P. Gribelyuk, The number of “magic” squares, cubes, and hypercubes, *Amer. Math. Monthly* **110** (2003), 707–717.
- [5] E. Bender and J. Goldman, Enumerative uses of generating functions, *Indiana Univ. Math. J.* **20** (1971), 753–764.
- [6] G. Birkhoff, Tres observaciones sobre el algebra lineal, *Univ. Nac. Tucumán Rev. Ser. A* **5** (1946), 147–150.
- [7] F. Bergeron, G. Labelle, and P. Leroux, *Combinatorial Species and Tree-Like Structures*, Cambridge University Press, Cambridge, 1998.
- [8] L. Comtet, *Advanced Combinatorics*, Reidel, Dordrecht and Boston, 1974.
- [9] J. A. De Loera, F. Liu and R. Yoshida, A generating function for all semi-magic squares and the volume of the Birkhoff polytope, *J. Algebraic Combin.* **30** (2009), 113–139.
- [10] A. Dress and T. W. Müller, Decomposable functors and the exponential principle, *Adv. in Math.* **129** (1997), 188–221.
- [11] C. Ehresmann, *Catégories et structures*, Dunod, Paris, 1965.
- [12] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, Cambridge, 2009.
- [13] D. Foata, *La série génératrice exponentielle dans les problèmes d’énumération*, Séminaire de mathématiques supérieures — été 1971, no. 54, Montréal, Les Presses de l’Université de Montréal, 1974.
- [14] I. P. Goulden and D. M. Jackson, *Combinatorial Enumeration*, John Wiley & Sons, New York, 1983.
- [15] H. Gupta, Enumeration of symmetric matrices, *Duke Math. J.* **35** (1968), 653–659.
- [16] A. Joyal, Une théorie combinatoire des séries formelles, *Adv. in Math.* **42** (1981), 1–82.
- [17] G. Labelle and P. Leroux, An extension of the exponential formula in enumerative combinatorics, *Electron. J. Comb.* **3**(2) (1996), Article R12, 14 pp.
- [18] P. A. MacMahon, *Combinatory Analysis*, vol. 2, Cambridge University Press, 1916; reprinted by Chelsea, New York, 1960.
- [19] M. Menni, *Symmetric monoidal completions and the exponential principle among labeled combinatorial structures*, Theory and Appl. Categories **11** (2003), 397–419.
- [20] M. Menni, *Algebraic categories whose projections are explicitly free*, Theory and Appl. Categories **22** (2009), 509–541.
- [21] A. D. Scott and A. D. Sokal, Some variants of the exponential formula, with application to the multivariate Tutte polynomial (alias Potts model), *Séminaire Lotharingien Combin.* **61A** (2009), Article B61Ae, 33 pp.
- [22] L. J. Slater, *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, 1966.

- [23] N. J. A. Sloane (editor), *The On-Line Encyclopedia of Integer Sequences*, available at <http://www.research.att.com:80/~njas/sequences/>.
- [24] R. P. Stanley, Linear homogeneous diophantine equations and magic labelings of graphs, *Duke Math. J.* **40** (1973), 607–632.
- [25] R. P. Stanley, Generating functions, in: *MAA Studies in Mathematics*, vol. 17 (G.-C. Rota, ed.), Math. Assoc. Am., Washington, 1978, 100–141.
- [26] R. P. Stanley, *Enumerative Combinatorics*, vol. 1, Wadsworth & Brooks/Cole, Pacific Grove, California, 1986; reprinted by Cambridge University Press, Cambridge, 1998.
- [27] R. P. Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge University Press, Cambridge, 1999.
- [28] H. S. Wilf, *generatingfunctionology*, 2nd edition, Academic Press, San Diego, 1994.

SCHOOL OF MATHEMATICAL SCIENCES, QUEEN MARY & WESTFIELD COLLEGE, UNIVERSITY OF LONDON, MILE END ROAD, LONDON E1 4NS, UNITED KINGDOM.

WWW: <http://www.maths.qmw.ac.uk/~pjc/>, <http://www.maths.qmw.ac.uk/~twm/>.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN, NORDBERGSTRASSE 15, A-1090 VIENNA, AUSTRIA. WWW: <http://www.mat.univie.ac.at/~kratt>.