



MTH4100 Calculus I

Lecture notes for Week 11

**Thomas' Calculus, Sections 7.1 to 7.8
(except Sections 7.5, 7.6)**

Rainer Klages

School of Mathematical Sciences
Queen Mary, University of London

Autumn 2008

Lecture 28

Definite integrals of symmetric functions


Theorem 1 Let f be continuous on the symmetric interval $[-a, a]$.

(a) If f is even, then $\int_{-a}^a f(x)dx = 2 \int_0^a f(x)dx$.

(b) If f is odd, then $\int_{-a}^a f(x)dx = 0$.


(proof by splitting the integrals and straightforward formal manipulations, see book p.379 for part (a))

example:



Areas between curves

example:




DEFINITION Area Between Curves

If f and g are continuous with $f(x) \geq g(x)$ throughout $[a, b]$, then the **area of the region between the curves $y = f(x)$ and $y = g(x)$ from a to b** is the integral of $(f - g)$ from a to b :

$$A = \int_a^b [f(x) - g(x)] dx.$$


example: Find the area that is enclosed above by $y = \sqrt{x}$ and below by $y = 0$ and $y = x - 2$.
 Two solutions:
 (a) by definition:



Split total area into area A + area B.
 Find right-hand limit for B by solving $\sqrt{x} = x - 2 \Rightarrow x = 4$.

$$\begin{aligned} \text{total area} &= \int_0^2 \sqrt{x} - 0 \, dx + \int_2^4 \sqrt{x} - (x - 2) \, dx \\ &= \left. \frac{2}{3} x^{3/2} \right|_0^2 + \left. \left(\frac{2}{3} x^{3/2} - \frac{1}{2} x^2 + 2x \right) \right|_2^4 \\ &= \frac{10}{3} \end{aligned}$$

(b) the clever way:



The area below the parabola is

$$A_1 = \int_0^4 \sqrt{x} \, dx = \left. \frac{2}{3} x^{3/2} \right|_0^4 = \frac{16}{3} .$$

The area of the triangle is $A_2 = 2 \cdot 2 / 2 = 2$ so that

$$\text{total area} = A_1 - A_2 = \frac{16}{3} - 2 = \frac{10}{3} .$$


Inverse functions and their derivatives

DEFINITION One-to-One Function

A function $f(x)$ is **one-to-one** on a domain D if $f(x_1) \neq f(x_2)$ whenever $x_1 \neq x_2$ in D .

These functions take on any value in their range *exactly once*.

examples:




Both functions are one-to-one on \mathbb{R} , respectively on \mathbb{R}_0^+ .

The Horizontal Line Test for One-to-One Functions

A function $y = f(x)$ is one-to-one if and only if its graph intersects each horizontal line at most once.

examples:



$y = x^2$ is one-to-one on, e.g., \mathbb{R}_0^+ but not \mathbb{R} .

$y = \sin x$ is one-to-one on, e.g., $[0, \pi/2]$ but not \mathbb{R} .

DEFINITION Inverse Function

Suppose that f is a one-to-one function on a domain D with range R . The **inverse function** f^{-1} is defined by

$$f^{-1}(a) = b \text{ if } f(b) = a.$$


The domain of f^{-1} is R and the range of f^{-1} is D .

note:


- f^{-1} reads f *inverse*
- $f^{-1}(x) \neq (f(x))^{-1} = 1/f(x)!$ (not an exponent)
- $(f^{-1} \circ f)(x) = x$ for all $x \in D(f)$
- $(f \circ f^{-1})(x) = x$ for all $x \in R(f)$

Read off inverse from graph of $f(x)$, as follows:


usual procedure $x \mapsto y = f(x)$:




for inverse $y \mapsto x = f^{-1}(y)$:



Note that $D(f) = R(f^{-1})$ and $R(f) = D(f^{-1})$, which suggests to *reflect* $x = f^{-1}(y)$ along $y = x$:



After reflection, x and y have changed places. Therefore, swap x and y ...




... and we have found $y = f^{-1}(x)$ graphically.

method for finding inverses *algebraically*:

1. solve $y = f(x)$ for x : $x = f^{-1}(y)$
2. interchange x and y : $y = f^{-1}(x)$

example: Find the inverse of $y = x^2, x \geq 0$.

1. solve $y = f(x)$ for x : $\sqrt{y} = \sqrt{x^2} = |x| = x$, as $x \geq 0$.
2. interchange x and y : $y = \sqrt{x}$.



Lecture 29

Calculate derivatives of inverse functions.

Differentiate $y = f^{-1}(x)$, or $x = f(y)$:

$$\frac{dx}{dy} = 1 = \frac{d}{dy}f(y) = f'(y)\frac{dy}{dy}.$$

Therefore,

$$\frac{dy}{dx} = \frac{1}{f'(y)} = \frac{1}{\frac{dx}{dy}}$$

The derivatives are reciprocals of one another.

Be precise: $x = f(y)$ means $y = f^{-1}(x)$ so that

$$\frac{dy}{dx} = \frac{1}{f'(f^{-1}(x))}$$

Be more precise:

THEOREM 1 The Derivative Rule for Inverses

If f has an interval I as domain and $f'(x)$ exists and is never zero on I , then f^{-1} is differentiable at every point in its domain. The value of $(f^{-1})'$ at a point b in the domain of f^{-1} is the reciprocal of the value of f' at the point $a = f^{-1}(b)$:

$$(f^{-1})'(b) = \frac{1}{f'(f^{-1}(b))}$$


or

$$\left. \frac{df^{-1}}{dx} \right|_{x=b} = \frac{1}{\left. \frac{df}{dx} \right|_{x=f^{-1}(b)}}$$

example: $f(x) = x^2, x \geq 0$ continued.

$f^{-1}(x) = \sqrt{x}$ and $f'(x) = 2x$ so that

$$(f^{-1})'(x) = \frac{1}{f'(f^{-1}(x))} = \frac{1}{2f^{-1}(x)} = \frac{1}{2\sqrt{x}}$$



note: The theorem can be used pointwise to find a value of the inverse derivative without calculating any formula for the inverse (see the book p.472 for an example). Otherwise, simply differentiate the inverse.

Natural Logarithms

For $a \in \mathbb{Q} \setminus \{-1\}$ we know that


$$\int_1^x t^a dt = \frac{1}{a+1} (x^{a+1} - 1)$$

(fundamental theorem of calculus part 2).

What happens if $a = -1$? $\int_1^x \frac{1}{t} dt$ is well defined for $x > 0$:

DEFINITION The Natural Logarithm Function

$$\ln x = \int_1^x \frac{1}{t} dt, \quad x > 0$$



The range of $\ln x$ is \mathbb{R} .

A special value: the number $e = 2.718281828459\dots$ (sometimes called *Euler's number*), satisfying

$$\ln e = 1.$$

Differentiate $\ln x$ (according to the fundamental theorem of calculus part 1):

$$\frac{d}{dx} \ln x = \frac{d}{dx} \int_1^x \frac{1}{t} dt = \frac{1}{x}.$$

If $u(x) > 0$, by the chain rule

$$\frac{d}{dx} \ln u = \frac{1}{u} u'.$$

If $u(x) = ax$ with $a > 0$,

$$\frac{d}{dx} \ln ax = \frac{1}{ax} a = \frac{1}{x}$$

Since $\ln ax$ and $\ln x$ have the same derivative (!),

$$\ln ax = \ln x + C.$$

For $x = 1$ we get $C = \ln a1 - \ln 1 = \ln a$ and therefore

$$\ln ax = \ln a + \ln x.$$

We have shown rule 1 in the following table:

THEOREM 2 Properties of Logarithms

For any numbers $a > 0$ and $x > 0$, the natural logarithm satisfies the following rules:

1. *Product Rule:* $\ln ax = \ln a + \ln x$
2. *Quotient Rule:* $\ln \frac{a}{x} = \ln a - \ln x$
3. *Reciprocal Rule:* $\ln \frac{1}{x} = -\ln x$ Rule 2 with $a = 1$
4. *Power Rule:* $\ln x^r = r \ln x$ r rational

(For the proof of rule 4 see book p.480.)

examples: Apply the logarithm properties to function formulas by replacing $a \rightarrow f(x)$, $x \rightarrow g(x)$.

1. $\ln 8 + \ln \cos x = \ln(8 \cos x)$
2. $\ln \frac{z^2 + 3}{2z - 1} = \ln(z^2 + 3) - \ln(2z - 1)$
3. $\ln \cot x = \ln \frac{1}{\tan x} = -\ln \tan x$
4. $\ln \sqrt[5]{x - 3} = \ln(x - 3)^{1/5} = \frac{1}{5} \ln(x - 3)$

For $t > 0$, the fundamental theorem of calculus tells us that

$$\int \frac{1}{t} dt = \ln t + C.$$

For $t < 0$, $(-t)$ is positive, and we find analogously

$$\int \frac{1}{(-t)} d(-t) = \ln(-t) + C.$$

For $t \neq 0$, together this gives

$$\int \frac{1}{t} dt = \ln |t| + C$$

Substituting $t = f(x)$, $dt = f'(x)dx$ leads to

$$\boxed{\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + C}$$

(for all $f(x)$ that maintain a constant sign on the range of integration).

example:

$$\int \tan x dx = \int \frac{\sin x}{\cos x} dx$$

Substitute $t = \cos x$, $dt = -\sin x dx$ on $(-\pi/2, \pi/2)$:

$$\int \tan x dx = - \int \frac{1}{t} dt = -\ln |t| + C = -\ln |\cos x| + C$$

Analogously for $\cot x$:


$$\int \tan u du = -\ln |\cos u| + C = \ln |\sec u| + C$$

$$\int \cot u du = \ln |\sin u| + C = -\ln |\csc u| + C$$

Lecture 30

The exponential function

$\ln x$ is strictly increasing, therefore invertible:



Definition 1 (Exponential function) For every $x \in \mathbb{R}$, $\exp x = \ln^{-1} x$.

Recall that $1 = \ln e$ so that $\exp 1 = e$.

Apply the power rule:

$$\ln e^r = r \ln e = r$$

so that

$$e^r = \exp r, \quad r \in \mathbb{Q}.$$

But $\exp x$ is defined for *any real* x , which suggests to define *real* exponents for base e via $\exp x$:

Definition 2 For every $x \in \mathbb{R}$, $e^x = \exp x$.

It is

$$\ln(e^a) = a, \quad a \in \mathbb{R}$$

and

$$e^{\ln a} = a, \quad a > 0.$$

With

$$(e^{\ln a})^x = e^{x \ln a} = a^x$$

we can define *real powers of positive real numbers* a :

Definition 3 (General exponential functions) For every $x \in \mathbb{R}$ and $a > 0$, the exponential function with base a is

$$a^x = e^{x \ln a}.$$

note: By using $x^n = e^{n \ln x}$, it can be proven that

$$\frac{d}{dx} x^n = nx^{n-1}, x > 0,$$

for all real n . (see book p.492)

We have

THEOREM 3 Laws of Exponents for e^x

For all numbers x, x_1 , and x_2 , the natural exponential e^x obeys the following laws:

1. $e^{x_1} \cdot e^{x_2} = e^{x_1+x_2}$
2. $e^{-x} = \frac{1}{e^x}$
3. $\frac{e^{x_1}}{e^{x_2}} = e^{x_1-x_2}$
4. $(e^{x_1})^{x_2} = e^{x_1 x_2} = (e^{x_2})^{x_1}$

Proof of 1.:

$$\begin{aligned} \exp(x_1) \cdot \exp(x_2) &= \exp \ln(\exp(x_1) \cdot \exp(x_2)) \\ (\text{product rule for } \ln x) &= \exp(\ln \exp(x_1) + \ln \exp(x_2)) \\ &= \exp(x_1 + x_2) \end{aligned}$$

As $e^x = f^{-1}(x)$ with $f(x) = \ln x$ and $f'(x) = 1/x$, we find (by using the derivative rule for inverses)

$$\frac{d}{dx} e^x = \frac{1}{f'(f^{-1}(x))} = f^{-1}(x) = e^x$$

implying

$$\int e^x dx = e^x + C.$$

By the chain rule,

$$\frac{d}{dx} e^{f(x)} = e^{f(x)} f'(x)$$

so that

$$\int e^{f(x)} f'(x) dx = e^{f(x)} + C.$$

examples:

1.

$$\frac{d}{dx} e^{\sin x} = e^{\sin x} \frac{d}{dx} \sin x = e^{\sin x} \cos x$$

2.

$$\begin{aligned}
 \int_0^{\ln 2} e^{3x} dx &= \int_0^{\ln 8} e^u \frac{1}{3} du \\
 &= \frac{1}{3} e^u \Big|_0^{\ln 8} \\
 &= \frac{7}{3}
 \end{aligned}$$

We defined e via $\ln e = 1$ and stated $e = 2.718281828459 \dots$

Theorem 2 (The number e as a limit)

$$e = \lim_{x \rightarrow 0} (1 + x)^{1/x}$$

Proof:

$$\begin{aligned}
 \ln \left(\lim_{x \rightarrow 0} (1 + x)^{1/x} \right) &= \\
 \text{(continuity of } \ln x \text{)} &= \lim_{x \rightarrow 0} (\ln(1 + x)^{1/x}) \\
 \text{(power rule)} &= \lim_{x \rightarrow 0} \left(\frac{1}{x} \ln(1 + x) \right) \\
 (\ln 1 = 0) &= \lim_{x \rightarrow 0} \frac{\ln(1 + x) - \ln(1)}{x} \\
 (f(t) = \ln t) &= \lim_{h \rightarrow 0} \frac{f(1 + h) - f(1)}{h} \\
 &= f'(1) \\
 &= 1 \\
 &= \ln(e)
 \end{aligned}$$

q.e.d.

Differentiate general exponential functions of base $a > 0$:

$$\frac{d}{dx} a^x = \frac{d}{dx} e^{x \ln a} = e^{x \ln a} \ln a = a^x \ln a$$

implying

$$\int a^x dx = \frac{a^x}{\ln a} + C, \quad a \neq 1$$

example:

$$\frac{d}{dx} x^x = \frac{d}{dx} e^{x \ln x} = e^{x \ln x} \frac{d}{dx} (x \ln x) = x^x (1 + \ln x)$$

Definition 4 ($\log_a x$) *The inverse of $y = a^x$ is*

$\log_a x$, *the logarithm of x with base a ,*

provided $a > 0$ and $a \neq 1$.

It is

$$\log_a(a^x) = x, \quad x \in \mathbb{R}$$

and

$$x = a^{\log_a x}, \quad x > 0.$$

Furthermore,

$$\ln x = \ln(a^{\log_a x}) = \log_a x \cdot \ln a.$$

yielding

$$\log_a x = \frac{\ln x}{\ln a}$$

note: The algebra for $\log_a x$ is precisely the same as that for $\ln x$.

Read

Thomas' Calculus:

Section 7.7 Inverse trigonometric functions,
and Section 7.8, Hyperbolic functions


You will need this information for coursework 10!

In the following two sections I will explain some very bare essentials that can be found on these pages.

Inverse trigonometric functions

note: \sin , \cos , \sec , \csc , \tan , \cot are not one-to-one *unless* the domain is restricted.

example:




Once the domains are suitably restricted, we can define:

$$\begin{aligned} \arcsin x &= \sin^{-1} x & \operatorname{arccsc} x &= \csc^{-1} x \\ \arccos x &= \cos^{-1} x & \operatorname{arcsec} x &= \sec^{-1} x \\ \arctan x &= \tan^{-1} x & \operatorname{arccot} x &= \cot^{-1} x \end{aligned}$$


examples:

Domain: $-1 \leq x \leq 1$
 Range: $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$



(a)

Domain: $-1 \leq x \leq 1$
 Range: $0 \leq y \leq \pi$



(b)

... and so on.

caution:

$$\sin^{-1} x \neq (\sin x)^{-1}$$

Unfortunately this is inconsistent, since $\sin^2 x = (\sin x)^2$. Best to avoid $\sin^{-1} x$ and use $\arcsin x$ etc. instead.

How to differentiate inverse trigonometric functions?

example: Differentiate $\arcsin x$.

Start with implicit differentiation of $\sin y = x$,

$$\cos y \frac{dy}{dx} = 1 .$$

Solve for $\frac{dy}{dx}$:

$$\frac{dy}{dx} = \frac{1}{\cos y} = \frac{1}{\sqrt{1 - \sin^2 y}}$$

for $-\pi/2 < y < \pi/2$ ($\cos x = 0$ for $x = \pm\pi/2$). Therefore, for $|x| < 1$,

$$\frac{d}{dx} \arcsin x = \frac{1}{\sqrt{1 - x^2}}$$

and, conversely,

$$\int \frac{dx}{\sqrt{1 - x^2}} = \arcsin x + C .$$

example: Evaluate

$$\int \frac{dx}{\sqrt{4x - x^2}} .$$

Trick: complete the square!

$$4x - x^2 = 4 - (x - 2)^2$$

Now integrate

$$\begin{aligned}\int \frac{dx}{\sqrt{4x-x^2}} &= \int \frac{dx}{\sqrt{4-(x-2)^2}} \\ (u=x-2) &= \int \frac{du}{\sqrt{4-u^2}} \\ &= \arcsin \frac{u}{2} + C \\ &= \arcsin \left(\frac{x}{2} - 1 \right) + C\end{aligned}$$

Hyperbolic functions

Every function f on $[-a, a]$ can be decomposed into

$$f(x) = \underbrace{\frac{f(x) + f(-x)}{2}}_{\text{even function}} + \underbrace{\frac{f(x) - f(-x)}{2}}_{\text{odd function}}$$


For $f(x) = e^x$:

$$e^x = \underbrace{\frac{e^x + e^{-x}}{2}}_{=\cosh x} + \underbrace{\frac{e^x - e^{-x}}{2}}_{=\sinh x},$$


called *hyperbolic sine* and *hyperbolic cosine*.

Define \tanh , \coth , sech , and csch in analogy to trigonometric functions.

examples:



$$\sinh x = \frac{e^x - e^{-x}}{2}$$



$$\cosh x = \frac{e^x + e^{-x}}{2}$$

Compare the following with trigonometric functions:

TABLE 7.6 Identities for hyperbolic functions

$$\cosh^2 x - \sinh^2 x = 1$$

$$\sinh 2x = 2 \sinh x \cosh x$$

$$\cosh 2x = \cosh^2 x + \sinh^2 x$$

$$\cosh^2 x = \frac{\cosh 2x + 1}{2}$$

$$\sinh^2 x = \frac{\cosh 2x - 1}{2}$$

$$\tanh^2 x = 1 - \operatorname{sech}^2 x$$

$$\operatorname{coth}^2 x = 1 + \operatorname{csch}^2 x$$

How to differentiate hyperbolic functions?

example:

$$\frac{d}{dx} \sinh x = \frac{d}{dx} \frac{e^x - e^{-x}}{2} = \frac{e^x + e^{-x}}{2} = \cosh x$$

$$\frac{d}{dx} \cosh x = \frac{d}{dx} \frac{e^x + e^{-x}}{2} = \frac{e^x - e^{-x}}{2} = \sinh x$$

Inverse hyperbolic functions defined in analogy to trigonometric functions.