

Uniform asymptotics of area-weighted Dyck paths

Nils Haug and Thomas Prellberg

Citation: *Journal of Mathematical Physics* **56**, 043301 (2015); doi: 10.1063/1.4917052

View online: <http://dx.doi.org/10.1063/1.4917052>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jmp/56/4?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[A note on asymptotic uniform confidence bands in a multivariate statistical deconvolution problem](#)
AIP Conf. Proc. **1479**, 438 (2012); 10.1063/1.4756159

[Asymptotic evolution of quantum walks with random coin](#)
J. Math. Phys. **52**, 042201 (2011); 10.1063/1.3575568

[Diffusion in a half-space: From Lord Kelvin to path integrals](#)
Am. J. Phys. **73**, 308 (2005); 10.1119/1.1842734

[Singularities in the complex temperature plane at the first order phase transitions and critical points](#)
AIP Conf. Proc. **519**, 314 (2000); 10.1063/1.1291573

[Random walks and effective resistances on toroidal and cylindrical grids](#)
Am. J. Phys. **68**, 37 (2000); 10.1119/1.19370



Uniform asymptotics of area-weighted Dyck paths

Nils Haug^{a)} and Thomas Prellberg

School of Mathematical Sciences, Queen Mary University of London, London E1 4NS, United Kingdom

(Received 11 January 2015; accepted 21 March 2015; published online 13 April 2015)

Using the generalized method of steepest descents for the case of two coalescing saddle points, we derive an asymptotic expression for the bivariate generating function of Dyck paths, weighted according to their length and their area in the limit of the area generating variable tending towards 1. The result is valid uniformly for a range of the length generating variable, including the tricritical point of the model. © 2015 AIP Publishing LLC. [<http://dx.doi.org/10.1063/1.4917052>]

I. INTRODUCTION

A Dyck path is a trajectory of a directed random walk on the two-dimensional square lattice above the diagonal, starting at the origin and ending on this diagonal. More precisely, the random walk starts at the point $(0,0)$ and, from any given point (x,y) , the random walker can only step towards $(x+1,y)$ and $(x,y+1)$. Steps from (x,y) to $(x-1,y)$ or to $(x,y-1)$ are forbidden. Furthermore, each point on the trajectory (x,y) must satisfy $x \leq y$, which means that the random walker always stays above the main diagonal $x = y$, and its final position (x,y) must be on the main diagonal. Figure 1 shows an example of a Dyck path as it is usually drawn with the lattice oriented such that the main diagonal lies horizontally in the image. As a possible physical system which can be described by Dyck paths, one can think of two-dimensional vesicles attached to a surface¹ or of discrete trajectories of charged particles moving in an external magnetic field.² Since we will only deal with Dyck paths in the following, we will refer to them simply as “paths.”

The *length* of a path is the number of steps it consists of. Since the number of horizontal steps must equal the number of vertical steps, it follows that the length of a path is always even. Further, the *area* of a path is the number of entire unit squares enclosed between the trajectory and the main diagonal. For example, the path in Figure 1 has length $n = 18$ and area $m = 10$ (occasionally, an alternative definition of the area under a path as the number of triangular plaquettes enclosed by the trajectory and the main diagonal is used; see, e.g., Ref. 3).

The generating function of paths of length $2n$, weighted according to their area is defined as

$$Z_n(q) = \sum_{m=1}^{\infty} c_{m,n} q^m, \quad (1)$$

where $c_{m,n}$ is the number of paths of length $2n$ and area m and q is the weight associated to the area. In physical terms, one can interpret $Z_n(q)$ as the canonical partition function of two-dimensional surface-attached vesicles with perimeter $4n$ and area-fugacity q . Note that the maximum area of a path of given finite length is bounded; therefore, the sum on the RHS of (1) is finite for finite values of n . As a physical consequence, phase transitions can only occur in the thermodynamic limit $n \rightarrow \infty$.

One can also define the generating function of paths of area m , weighted with respect to their length as

$$Q_m(t) = \sum_{n=0} c_{m,n} t^n, \quad (2)$$

^{a)}n.a.haug@qmul.ac.uk




FIG. 1. A Dyck path of area $m = 10$ and length $n = 18$. The shaded squares have unit area. The white triangles on the bottom do not contribute to the area.

where t is the weight conjugate to the length. The generating function of paths weighted according to both their area *and* their length is defined as

$$G(t, q) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} c_{m,n} q^m t^n. \quad (3)$$

With (1), this can be rewritten as

$$G(t, q) = \sum_{n=1}^{\infty} Z_n(q) t^n \quad (4)$$

and with (2), we can write

$$G(t, q) = \sum_{m=1}^{\infty} G_m(t) q^m. \quad (5)$$

By a standard factorization argument,⁴ one obtains the functional equation

$$G(t, q) = 1 + t G(t, q) G(qt, q), \quad (6)$$

which can be solved by using the ansatz,

$$G(t, q) = \frac{H(qt)}{H(t)}. \quad (7)$$

Here,

$$H(t) = \sum_{n=0}^{\infty} \frac{q^{n^2-n} (-t)^n}{(q; q)_n}, \quad (8)$$

and we have used the standard notation for the q-Pochhammer symbol,

$$(z; q)_n = \prod_{k=0}^{n-1} (1 - zq^k), \quad (9)$$

which is a q-generalization of the Pochhammer symbol. The function $H(qt) = \text{Ai}_q(t)$ is a q-Airy function.⁵

For $q = 1$, (6) is solved by the generating function of the Catalan numbers (see, e.g., Ref. 14),

$$G(t, 1) = \frac{1}{2t} (1 - \sqrt{1 - 4t}). \quad (10)$$

In Figure 2, we show how the radius of convergence t_{∞} of the series on the RHS of Eq. (4) behaves qualitatively as a function of q . This picture, which is also called the “phase diagram” of the system, is typical for lattice polygon models.⁶ The radius of convergence is defined by a decreasing line of pole singularities for $q < 1$ and zero for $q > 1$. From (10), the value of t_{∞} for $q = 1$ can be deduced to be $t_c = 1/4$. The point $(t, q) = (t_c, 1)$ is called the *tricritical point* of the model (see, e.g., Ref. 7) and the area below $t_{\infty}(q)$ is called the *finite size region*.




FIG. 2. The qualitative behaviour of the radius of convergence t_∞ of $G(t, q)$ as a function of q . The small circle marks the tricritical point.

In Ref. 3, the general form of continued fraction expressions for generating functions of Dyck and Motzkin paths has been discussed. In particular,

$$G(t, q) = \frac{1}{1 - \frac{t}{1 - \frac{tq}{1 - \frac{tq^2}{1 - \frac{tq^3}{1 - \dots}}}}}. \quad (11)$$

This expression enables us to continue $G(t, q)$ analytically beyond the finite size region.

The asymptotic behaviour of $G(t, q)$ for $q \rightarrow 1^-$ as one approaches the tricritical point has so far not been derived rigorously. The aim of this paper is to close this gap by rigorously deriving an asymptotic expression for the generating function $G(t, q)$ in the limit $q \rightarrow 1^-$ which is valid uniformly for a range of values of t including the critical point t_c . A similar calculation has been carried out in Ref. 8 for staircase polygons.

Our main result is given in Proposition 2.1 and Corollary 2.2.

II. RESULTS

According to Eq. (7), the function $G(t, q)$ is given as a quotient of two alternating q -series. In order to obtain its asymptotic behaviour, we first derive the asymptotic behaviour of both the numerator and the denominator separately. Taking the fraction of the two obtained expressions will then lead us to the asymptotic behaviour of $G(t, q)$. We will start with the asymptotic expansion of $H(t)$.

A. Uniform asymptotic expansion of $H(t)$


The first step in our calculation is to express $H(t)$ as a contour integral. We prove

Lemma 2.1. For complex t and $0 < q < 1$,

$$H(t) = \frac{(q; q)_\infty}{2\pi i} \int_C \frac{z^{(1+\log_q z)/2 - \log_q t}}{(z; q)_\infty} dz, \quad (12)$$

where C is a contour as shown in Figure 3(b).

Proof. For complex $q \neq 0$ and $n \in \mathbb{N}_0$, we have

FIG. 3. The contours C_1 (a) and C (b).

$$\frac{(-1)^n q^{\binom{n}{2}}}{(q; q)_n (q; q)_\infty} = -\text{Res} \left[(z; q)_\infty^{-1}; z = q^{-n} \right], \quad (13)$$

from which it follows that

$$\frac{(-t)^n q^{n^2-n}}{(q; q)_n (q; q)_\infty} = -\text{Res} \left[\frac{z^{(1+\log_q z)/2 - \log_q t}}{(z; q)_\infty}; z = q^{-n} \right]. \quad (14)$$

Suppose now that $0 < q < 1$. Then, the contour $C_N = C_N^1 \cup C_N^2 \cup C_N^3$, where

$$\left. \begin{aligned} C_N^1 &= \{ \rho + \lambda e^{-i\psi} \mid 0 < \lambda < q^{-N-1/2} \} \\ C_N^2 &= \{ \rho + \lambda e^{i\varphi} \mid 0 < \lambda < q^{-N-1/2} \} \\ C_N^3 &= \{ \rho + q^{-N-1/2} e^{i\theta} \mid -\psi < \theta < \varphi \} \end{aligned} \right\}, \quad (15)$$

$0 < \rho < 1$ and $(\varphi, \psi) \in]0, \pi[^2$, surrounds exactly the N leftmost singularities of the integrand on the RHS of Eq. (12)—see Figure 3(a). We can therefore write

$$\sum_{n=0}^N \frac{q^{n^2-n} (-t)^n}{(q; q)_n} = \frac{(q; q)_\infty}{2\pi i} \oint_{C_N} \frac{z^{(1+\log_q z)/2 - \log_q t}}{(z; q)_\infty} dz, \quad (16)$$

where the integration is performed in clockwise sense, as indicated by the arrows in Figure 3(a). Combining (8) and (16), we obtain

$$H(t) = \lim_{N \rightarrow \infty} \frac{(q; q)_\infty}{2\pi i} \oint_{C_N} \frac{z^{(1+\log_q z)/2 - \log_q t}}{(z; q)_\infty} dz. \quad (17)$$

It is left to show that in the limit $N \rightarrow \infty$, the contribution of the circle segment C_N^3 to contour integral (17) vanishes, such that the contour C_N can be replaced by the contour shown in Figure 3(b). On C_N^3 , we can estimate the denominator of the integrand on the RHS of (17) as

$$\begin{aligned} |(z, q)_\infty| &= \left| \prod_{n=0}^{\infty} (1 - q^{-N-1/2+n} e^{i\varphi}) \right| \geq \left| \prod_{n=0}^{\infty} (1 - q^{-N-1/2+n}) \right| \\ &= \left| \prod_{n=0}^{\infty} (1 - q^{-1/2+n}) \right| \cdot \left| \prod_{n=1}^N (1 - q^{-1/2-n}) \right| \geq c_1 \left| \prod_{n=1}^N q^{-1/2-n} \right| = c_1 |q^{-N^2/2-N}|, \end{aligned} \quad (18)$$

where c_1 is a constant independent of N . Furthermore, the absolute value of the numerator has for $z \in C_N^3$ the upper bound

$$|z^{(1+\log_q z)/2 - \log_q t}| \leq c_2 |q^{N^2/2}| |t|^N, \quad (19)$$

where c_2 is another constant independent of N . Therefore, we can estimate

$$\frac{1}{2\pi i} \oint_{C_N^3} \frac{z^{(1+\log_q z)/2 - \log_q t}}{(z; q)_\infty} dz \leq c_3 |q|^{N^2} |t|^N, \quad (20)$$

where c_3 is a third constant independent of N . Since the expression on the right hand side tends to zero as $N \rightarrow \infty$ for $q < 1$, it follows that

$$\lim_{N \rightarrow \infty} \frac{1}{2\pi i} \oint_{C_N} \frac{z^{(1+\log_q z)/2 - \log_q t}}{(z; q)_\infty} dz = \frac{1}{2\pi i} \int_C \frac{z^{(1+\log_q z)/2 - \log_q t}}{(z; q)_\infty} dz, \quad (21)$$

where

$$C = \{\rho + \lambda e^{-i\psi} \mid 0 < \lambda < \infty\} \cup \{\rho + \lambda e^{i\psi} \mid 0 < \lambda < \infty\}, \quad (22)$$

and where the integration is carried out as indicated by the arrows in Figure 3(b). Combining (21) with (17), we obtain (12). $\square$

In Ref. 8, it was shown by applying the Euler-Maclaurin summation formula that

$$\ln(z; q)_\infty = \frac{1}{\log(q)} \text{Li}_2(z) + \frac{1}{2} \ln(1-z) + \ln(q) R(z, q), \quad (23)$$

where the remainder satisfies the non-uniform bound

$$|R(z, q)| \leq \frac{1}{6} \left(\ln |1-z| + \frac{\text{Re}(z)}{\text{Im}(z)} \arctan \frac{\text{Im}(z)}{1 - \text{Re}(z)} \right). \quad (24)$$

Here, Li_2 denotes the Euler dilogarithm,⁹ which can be defined as

$$\text{Li}_2(z) = - \int_0^z \frac{\ln(1-s)}{s} ds. \quad (25)$$

Combining (12) with (23) and defining

$$f(z, t) = \ln(t) \ln(z) + \text{Li}_2(z) - \frac{1}{2} \ln(z)^2, \quad (26a)$$

$$g(z) = \left(\frac{z}{1-z} \right)^{1/2}, \quad (26b)$$

we can write

$$H(t) = \frac{(q; q)_\infty}{2\pi i} \int_C \exp \left(\frac{1}{\epsilon} f(z, t) + \epsilon R(z, q) \right) g(z) dz, \quad (27)$$

where $\epsilon = -\ln(q)$.

The function $f(z, t)$ is analytic for $\arg(z) < \pi$ and $\arg(1-z) < \pi$ and has the two saddle points

$$z_1(t) = \frac{1}{2} (1 + \sqrt{1-4t}), \quad z_2(t) = \frac{1}{2} (1 - \sqrt{1-4t}), \quad (28)$$

which coalesce for $t = t_c = 1/4$.

From the identity

$$\text{Li}_2(\lambda e^{i\phi}) = -\frac{1}{2} \ln(-\lambda e^{i\phi})^2 - \frac{\pi^2}{6} - \text{Li}_2\left(\frac{1}{\lambda} e^{-i\phi}\right) \quad (29)$$

(see, e.g., Ref. 10), we obtain

Lemma 2.2. For $0 < |\phi| \leq \pi$,

$$\text{Li}_2(\lambda e^{i\phi}) \sim -\frac{1}{2} \ln(-\lambda e^{i\phi})^2, \quad (30a)$$

$$\text{Re}[\text{Li}_2(\lambda e^{i\phi})] \sim -\frac{1}{2} \ln(\lambda)^2, \quad (30b)$$

$$\text{Im}[\text{Li}_2(\lambda e^{i\phi})] \sim -\frac{1}{2} \text{Im}[(\ln(-\lambda e^{i\phi}))^2] \quad (30c)$$

as $\lambda \rightarrow \infty$.

Consequently, we have the following.

Corollary 2.1. For complex t and $0 < |\phi| < \pi$,

$$f(\lambda e^{i\phi}, t) \sim -\ln(\lambda)^2 - i\psi \ln(\lambda) \quad \text{as } \lambda \rightarrow \infty, \quad (31)$$

where $\psi = 2\phi + \pi$ for $\phi < 0$ and $\psi = 2\phi - \pi$ for $\phi > 0$.

The remainder $R(z, q)$ is not uniformly bounded with respect to z ; therefore, it is not immediately clear that it can be neglected in the limit $\epsilon \rightarrow 0^+$. However, from the asymptotic behaviour of $f(z, t)$, one can conclude that the tails of the integration contour do not contribute to the asymptotics of the integral. Therefore, we have

Lemma 2.3. For complex t and $0 < q < 1$,

$$H(t) \sim \frac{(q; q)_\infty}{2\pi i} \int_C \exp\left(\frac{1}{\epsilon} f(z, t)\right) g(z) dz \quad \text{as } \epsilon \rightarrow 0^+. \quad (32)$$

It was shown in Ref. 11 (see also Refs. 12 and 13) that for a function $f(z, t)$, which is analytic with respect to both z and t and which has two saddle points $z_1(t)$ and $z_2(t)$, there is a unique transformation $u : z \mapsto u(z)$, such that

$$f(z, t) = \frac{1}{3}u^3 - \alpha(t)u + \beta(t), \quad (33)$$

which is regular and one-to-one in a domain containing $z_1(t)$ and $z_2(t)$ if t lies in some small domain containing t_c . Moreover,

$$u(z_1(t)) = |\alpha(t)^{1/2}|, \quad u(z_2(t)) = -|\alpha(t)^{1/2}|. \quad (34)$$

Combining (33) with (34), one gets the explicit form

$$u(z) = \left[\left(\frac{3}{2}(f(z, t) - \beta) \right)^2 + \left(\left(\frac{3}{2}(f(z, t) - \beta) \right)^2 - \alpha^3 \right)^{1/2} \right]^{1/3} + \alpha \left[\left(\frac{3}{2}(f(z, t) - \beta) \right)^2 + \left(\left(\frac{3}{2}(f(z, t) - \beta) \right)^2 - \alpha^3 \right)^{1/2} \right]^{-1/3}, \quad (35)$$

where the two parameters are obtained by inserting (34) into (33) as

$$\begin{aligned} \alpha(t) &= \left(\frac{3}{4} [f(z_2, t) - f(z_1, t)] \right)^{2/3} \\ \beta(t) &= \frac{1}{2} (f(z_1, t) + f(z_2, t)) = \frac{1}{2} \ln(t)^2 + \frac{\pi^2}{6} \end{aligned} \quad (36)$$

Note that for $t \rightarrow t_c$, one has $\alpha(t) \sim 1 - 4t$. From Corollary 2.1, it follows that

$$u(\rho + i\lambda) \sim \exp\left(\pm i \frac{\pi}{3}\right) \left(\frac{3}{2} \ln |\lambda| \right)^{1/3}, \quad (37)$$

for $\lambda \rightarrow \pm\infty$. This leads us to

Lemma 2.4. For complex t ,

$$H(t) \sim \frac{(q; q)_\infty}{2\pi i} \int_C e^{\frac{1}{\epsilon} \left(\frac{1}{3}u^3 - \alpha u + \beta \right)} g(z(u)) \frac{dz}{du} du \quad (38)$$

as $\epsilon \rightarrow 0^+$, where C is a contour as shown in Figure 3(b) with $\varphi = \psi = \pi/3$.

It is possible to write

$$g(z(u)) \frac{dz}{du} = \sum_{m=0}^{\infty} (p_m + uq_m)(u^2 - \alpha)^m \quad (39)$$

and insert this expansion into (38). Interchanging the order of integration and summation, we obtain the asymptotic expansion,

$$H(t) \sim \frac{(q; q)_{\infty}}{2\pi i} \sum_{m=0}^{\infty} \int_C (p_m + uq_m)(u^2 - \alpha)^m e^{\frac{1}{\epsilon} \left(\frac{1}{3}u^3 - \alpha u + \beta \right)} du. \quad (40)$$

The two leading coefficients can be obtained from (33) and (34). We get

$$2g(z_1) \sqrt{\frac{\alpha}{f''(z_1)}} = p_0 + \alpha^{1/2} q_0, \quad 2g(z_2) \sqrt{\frac{\alpha}{f''(z_2)}} = p_0 - \alpha^{1/2} q_0, \quad (41)$$

and these two equations can be solved with respect to p_0 and q_0 , respectively, to obtain

$$p_0 = \left(\frac{\alpha}{d} \right)^{\frac{1}{4}} (z_1^{3/2} + z_2^{3/2}), \quad q_0 = \left(\frac{1}{\alpha d} \right)^{\frac{1}{4}} (z_1^{3/2} - z_2^{3/2}). \quad (42)$$

Here, we have set $d = 1 - 4t$.

Inserting (42) into (40), we arrive at an asymptotic expression for $H(t)$ in terms of the Airy function,

$$\text{Ai}(z) = \int_C \exp \left(\frac{w^3}{3} - zw \right) dw. \quad (43)$$

This presents the main result of this section.

Lemma 2.5. For complex t ,

$$H(t) \sim \frac{(q; q)_{\infty}}{2\pi i} \left(\frac{1}{\alpha d} \right)^{\frac{1}{4}} \exp \left(\frac{\beta}{\epsilon} \right) \left(\alpha^{1/2} \epsilon^{1/3} (z_1^{3/2} + z_2^{3/2}) \text{Ai}(\alpha \epsilon^{-2/3}) + \right. \\ \left. + \epsilon^{2/3} (z_2^{3/2} - z_1^{3/2}) \text{Ai}'(\alpha \epsilon^{-2/3}) \right) \quad \text{as } \epsilon \rightarrow 0^+. \quad (44)$$

B. Uniform asymptotic expansion of $H(qt)$

The approach used in Sec. II A can be applied in a completely analogous way to $H(qt)$. The function $f(z, t)$ remains the same, whereas now

$$g(z) = \left(\frac{1}{z(1-z)} \right)^{1/2}. \quad (45)$$

This changes the leading coefficients of the expansion towards

$$p_0 = \left(\frac{\alpha}{d} \right)^{\frac{1}{4}} (z_1^{1/2} + z_2^{1/2}), \quad q_0 = \left(\frac{1}{\alpha d} \right)^{\frac{1}{4}} (z_1^{1/2} - z_2^{1/2}), \quad (46)$$

and we obtain the following.

Lemma 2.6. For complex t ,

$$H(qt) \sim \frac{(q; q)_{\infty}}{2\pi i} \left(\frac{1}{\alpha d} \right)^{\frac{1}{4}} \exp \left(\frac{\beta}{\epsilon} \right) \left(\alpha^{1/2} \epsilon^{1/3} (z_1^{1/2} + z_2^{1/2}) \text{Ai}(\alpha \epsilon^{-2/3}) + \right. \\ \left. + \epsilon^{2/3} (z_2^{1/2} - z_1^{1/2}) \text{Ai}'(\alpha \epsilon^{-2/3}) \right) \quad \text{as } \epsilon \rightarrow 0^+. \quad (47)$$




FIG. 4. Plot of $G(t, q)$ (black) against uniform asymptotic expression (48) (grey) for $\epsilon = 10^{-2}$ and t ranging between 0 and $1/2$ (horizontal axis).

C. Uniform asymptotics of $G(t, q)$

Combining Lemmas 2.5 and 2.6, we arrive at the following.

Proposition 2.1. For complex t and $q \rightarrow 1^-$,

$$G(t, q) \sim \frac{\alpha^{1/2}(z_1^{1/2} + z_2^{1/2})\text{Ai}(\alpha\epsilon^{-2/3}) + (z_2^{1/2} - z_1^{1/2})\epsilon^{1/3}\text{Ai}'(\alpha\epsilon^{-2/3})}{\alpha^{1/2}(z_1^{3/2} + z_2^{3/2})\text{Ai}(\alpha\epsilon^{-2/3}) + (z_2^{3/2} - z_1^{3/2})\epsilon^{1/3}\text{Ai}'(\alpha\epsilon^{-2/3})}, \quad (48)$$

where $\epsilon = -\ln(q)$,

$$\left. \begin{aligned} z_1 &= \frac{1}{2}(1 + \sqrt{1 - 4t}) \\ z_2 &= \frac{1}{2}(1 - \sqrt{1 - 4t}) \end{aligned} \right\},$$

and $\alpha(t)$ is given by Eq. (36).

One easily shows that for $t \leq t_c$ and $q \rightarrow 1^-$, $G(t, q)$ tends towards the generating function of the Catalan numbers; hence, our result is consistent with (10). By applying Dini's theorem, one can further show that the convergence is *uniform* for $t \in [0, t_c]$. However, (48) is also valid for $t > t_c$, though not in a uniform sense due to the occurrence of poles in the denominator. This fact is illustrated in Figure 4, where we have plotted $G(t, q)$ against the uniform asymptotic expression (48) for $\epsilon = 10^{-2}$ and $t \in [0, 1/2]$.

Defining the tricritical scaling function

$$F(s) := \frac{\text{Ai}'(s)}{\text{Ai}(s)}, \quad (49)$$

we can conclude

Corollary 2.2. For fixed $s = (1 - 4t) \cdot (1 - q)^{-\phi}$ and $q \rightarrow 1^-$,

$$G(t, q) \sim 2 \left[1 + (1 - q)^{-\gamma_0} F((1 - 4t)(1 - q)^{-\phi}) \right], \quad (50)$$

where the critical exponents are $\phi = 2/3$ and $\gamma_0 = -1/3$.

In particular,

$$G(t_c, q) \sim 2 \cdot \left(1 + A_0 \cdot (1 - q)^{-\gamma_0} \right) \quad (q \rightarrow 1^-), \quad (51)$$

where $A_0 = \text{Ai}'(0)/\text{Ai}(0) = -0.72 \dots$.

Both (48) and (50) can be rearranged in order to obtain an asymptotic expression for $F(s)$. In Figure 5, we have plotted $-2F(s)$ against the expressions obtained from (48) and (50) for $\epsilon = 10^{-3}$, 10^{-4} , and 10^{-5} . For fixed value of q , (48) provides a more accurate approximation than (50).




FIG. 5. Plot of the scaling function $F(s)$ (black) against asymptotic expressions (48) (a) and (50) (b) (grey) for $\epsilon = 10^{-3}$, 10^{-4} , and 10^{-5} (the smallest value of ϵ corresponds to the closest approximation).

D. Finite size scaling

Given observed scaling behaviour (50) of the generating function, we now aim to calculate an expression for the finite size scaling function of fixed-area partition function (2). For this purpose, we first derive a further asymptotic expression for $G(t, q)$.

We define the *singular part* of $G(t, q)$ as

$$G_{\text{sing}}(t, q) := G(t, q) - \frac{1}{2t} \quad (52)$$

and prove

Proposition 2.2. Let $I = [t_0, t_c]$, where $0 < t_0 \leq t_c$. Then,

$$G_{\text{sing}}(t, q) \sim \frac{1}{2t}(1-q)^{-\gamma_0} F((1-4t)(1-q)^{-\phi}) \quad (q \rightarrow 1^-), \quad (53)$$

uniformly for $t \in I$.

Proof. By using that $F(s) \sim -\sqrt{s}$ for $s \rightarrow \infty$, one easily sees that for $t \in I$, the RHS of (53) converges in a pointwise sense towards $C_{\text{sing}} := -\sqrt{1-4t}/2t$, which is a continuous function on I . One also proves easily that for all $t \in I$, the RHS of (53) decreases monotonically with q . It therefore follows from Dini's theorem that the RHS converges uniformly to $C_{\text{sing}}(t)$. It is also clear that the same holds for the LHS, $G_{\text{sing}}(t, q)$. Hence, both sides of (53) have the same uniform asymptotic expression and are therefore uniformly asymptotic to each other. $\square$

It is possible to derive the Hadamard product expression,¹⁵

$$\text{Ai}(s) = \text{Ai}(0) \exp(-A_0 s) \prod_{k=1}^{\infty} \left(1 - \frac{s}{s_k}\right), \quad (54)$$

where s_k is the k th zero of the Airy function. Taking the derivative of the logarithm of this expression and exercising some careful analysis, we get

$$F(s) = -\frac{1}{s} \sum_{j=1}^{\infty} Z(j) s^j, \quad (55)$$

where we have used the Airy Zeta function,

$$Z(j) = \sum_{k=1}^{\infty} \left(\frac{1}{s_k}\right)^j \quad (56)$$

and inserted the conjectured value $Z(1) = -\text{Ai}'(0)/\text{Ai}(0)$.¹⁶

Inserting (55) into (53), we obtain that for $m \rightarrow \infty$,

$$\begin{aligned} G_{\text{sing}}(t, q) &\sim - \sum_{j=0}^{\infty} Z(j+1)(1-4t)^j(1-q)^{-2/3j+1/3} \\ &= - \sum_{m=0}^{\infty} \sum_{j=0}^{\infty} Z(j+1)(1-4t)^j \binom{m + \frac{2}{3}j - \frac{4}{3}}{m} q^m. \end{aligned} \quad (57)$$

For $n \in \mathbb{N}$ and $\alpha \in \mathbb{C} \setminus \mathbb{Z}_{\leq 0}$, Theorem VI.1 from Ref. 4 states that

$$[z^n](1-z)^{-\alpha} = \binom{n+\alpha-1}{n} \sim \frac{n^{\alpha-1}}{\Gamma(\alpha)} \left(1 + \frac{\alpha(\alpha-1)}{2n}\right). \quad (58)$$

Inserting the leading order of this expansion into (57) and extracting the m th coefficient, one formally arrives at

$$Q_m(t) \sim m^{-4/3} \sum_{j=0}^{\infty} \frac{Z(j+1)}{\Gamma(\frac{2}{3}j - \frac{1}{3})} m^{2/3j} (1-4t)^j \quad (m \rightarrow \infty). \quad (59)$$

Defining the finite size scaling function

$$\phi(s) := \sum_{j=0}^{\infty} \frac{Z(j)}{\Gamma(\frac{2}{3}j - \frac{1}{3})} s^j, \quad (60)$$

we can rewrite (59) as

$$Q_m(t) \sim m^{-4/3} \phi((1-4t)m^{2/3}). \quad (61)$$

This expression is of the generic form expected for models which exhibit tricritical scaling.^{7,17}

III. CONCLUSION

We have calculated an asymptotic expression for the generating function of Dyck paths, weighted with respect to both their perimeter and their area in the limit of the area generating variable tending towards 1. The result is valid uniformly for a range of values of the perimeter generating variable, including the tricritical point.

In the limit of both the perimeter and the area generating variable tending towards their critical values, we have shown the existence of a scaling function, expressible via Airy functions and their derivatives. The same type of scaling expression has been proven before to hold in the case of staircase polygons.⁸

Note in particular that the scaling function is obtained as a particular limit of the uniform asymptotic expansion. This is in contrast to the behaviour found in Ref. 8, where the scaling function and the uniform asymptotic expansion are related by a local variable transformation. In Ref. 17, uniform asymptotic expansions for tricritical phase transitions were also constructed from scaling functions using such transformations.

From the scaling function of the generating function, we derived an expression for the scaling function of the finite-area partition function.

¹ A. L. Owczarek and T. Prellberg, "Pressure exerted by a vesicle on a surface," *J. Phys. A: Math. Theor.* **47**, 215001 (2014).

² K. Bulycheva, A. Gorsky, and S. Nechaev, "Critical behavior in topological ensembles," <http://arxiv.org/pdf/1409.3350.pdf> (2014).

³ P. Flajolet, "Combinatorial aspects of continued fractions," *Discrete Math.* **32**, 125–161 (1980).

⁴ P. Flajolet and R. Sedgewick, *Analytic Combinatorics* (Cambridge University Press, 2009).

⁵ M. E. H. Ismail, "Asymptotics of q-orthogonal polynomials and a q-airy function," *Int. Math. Res. Not.* **18**, 1063–1088 (2005).

⁶ C. Richard, in *Polygons, Polyominoes and Polycubes*, edited by A. J. Guttmann (Springer, Netherlands, 2009), Vol. 775, pp. 247–299.

⁷ E. J. J. V. Rensburg, *The Statistical Mechanics of Interacting Walks, Polygons, Animals and Vesicles*, Oxford Lecture Series in Mathematics Vol. 18 (Oxford University Press, 2000).

- ⁸ T. Prellberg, “Uniform q-series asymptotics for staircase polygons,” *J. Phys. A: Math. Gen.* **28**, 1289–1304 (1995).
- ⁹ NIST Digital Library of Mathematical Functions, <http://dlmf.nist.gov/>.
- ¹⁰ L. C. Maximon, “The dilogarithm function for complex argument,” *Proc. R. Soc. London, Ser. A* **459**, 2807–2819 (2003).
- ¹¹ C. Chester, B. Friedman, and F. Ursell, “An extension of the method of steepest descents,” *Math. Proc. Cambridge Philos. Soc.* **53**, 599–611 (1957).
- ¹² N. Bleistein and R. A. Handelsman, *Asymptotic Expansions of Integrals* (Holt, Rinehart and Winston, 1975).
- ¹³ R. Wong, *Asymptotic Approximations of Integrals*, Classics in Applied Mathematics (Society for Industrial and Applied Mathematics, 2001).
- ¹⁴ M. Aigner, *A Course in Enumeration* (Springer–Verlag, Berlin Heidelberg, 2007).
- ¹⁵ P. Flajolet and G. Louchard, “Analytic variations on the airy distribution,” *Algorithmica* **31**, 361–377 (2001).
- ¹⁶ R. E. Crandall, “On the quantum zeta function,” *J. Phys. A: Math. Gen.* **29**, 6795–6816 (1996).
- ¹⁷ R. Brak and A. L. Owczarek, “On the analyticity properties of scaling functions in models of polymer collapse,” *J. Phys. A: Math. Gen.* **28**, 4709–4725 (1995).