

PAPER

## Multicritical scaling in a lattice model of vesicles

To cite this article: N Haug and T Prellberg 2020 *J. Phys. A: Math. Theor.* **53** 265003

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

# Multicritical scaling in a lattice model of vesicles

N Haug<sup>1,2,3,4</sup>  and T Prellberg<sup>1</sup> 

<sup>1</sup> School of Mathematical Sciences, Queen Mary University of London, London, United Kingdom

<sup>2</sup> CeMSIIS, Section for Science of Complex Systems, Medical University of Vienna, Vienna, Austria

<sup>3</sup> Complexity Science Hub Vienna, Vienna, Austria

E-mail: [nina.haug@meduniwien.ac.at](mailto:nina.haug@meduniwien.ac.at)

Received 28 January 2020, revised 6 May 2020

Accepted for publication 12 May 2020

Published 9 June 2020



CrossMark

## Abstract

Vesicles, or closed fluctuating membranes, have been modelled in two dimensions by self-avoiding polygons, weighted with respect to their perimeter and enclosed area, with the simplest model given by area-weighted excursions (Dyck paths). These models generically show a tricritical phase transition between an inflated and a crumpled phase, with a scaling function given by the logarithmic derivative of the Airy function. Extending such a model, we find realisations of multicritical points of arbitrary order, with the associated multivariate scaling functions expressible in terms of generalised Airy integrals, as previously conjectured by John Cardy. This work therefore adds to the small list of models with a critical phase transition, for which exponents and the associated scaling functions are explicitly known.

Keywords: statistical mechanics, phase transitions, multicritical points, vesicles

(Some figures may appear in colour only in the online journal)

## 1. Introduction

Obtaining a thorough understanding of phase transitions is one of the main aims of statistical physics. For a continuous transition one would like to know the critical exponents describing the singular power-law behaviour of thermodynamic quantities as the transition is approached. Moreover, in the vicinity of such a transition it is generally believed that the thermodynamic quantities depend only on a suitably scaled combination of the parameters in terms of a universal scaling function [1]. Most progress has been made in two dimensions

<sup>4</sup> Author to whom any correspondence should be addressed.

with the help of conformal invariance [2]. While these scaling functions can be easily obtained numerically, there are only few instances for which one knows precise expressions, one classical example being the spin–spin correlations of the two-dimensional Ising model [3]. Based on field-theoretic arguments, John Cardy postulated that by including many-body interactions in a model of vesicles, a hierarchy of scaling functions could be found, but cautioned that due to the technical limitations of the method used, ‘it is very difficult to say to what these higher multicritical points might correspond physically’ [4]. In this letter, we give explicit examples of a statistical mechanical model having precisely these scaling functions, thereby providing a resolution to this problem.

## 2. Vesicles and self-avoiding polygons

A vesicle consists of a closed membrane formed from a lipid bilayer inside a watery solution. Depending on parameters such as the temperature and the osmotic pressure difference between the outside and the inside of the membrane, vesicles are found in different typical conformations [5]. Subject to thermal fluctuations, a vesicle of fixed surface area favours ‘crumpled’ configurations with relatively small volume if there is a large net pressure acting onto the outside of the membrane. On the other hand, if there is a net pressure acting onto the inside of the membrane, then the vesicle tends to appear in an inflated shape with larger volume.


In [6], a two-dimensional model of vesicles was proposed in terms of ring polymers enclosing an area. In this case, the length of the polymer plays the role of the surface area of the vesicle, and the volume of the vesicle becomes the enclosed area. In [7, 8] the vesicles were modelled as self-avoiding polygons (SAP) on the square lattice—see figure 1 for an example. Note that any intrinsic property of the vesicle membrane such as stiffness is neglected in that lattice model.

In order to analyse the model of SAP, one defines the area–perimeter generating function


$$P(x, q) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} p_{m,n} x^m q^n, \quad (1)$$

where  $p_{m,n}$  is the number of SAP with perimeter  $m$  and area  $n$ , with two SAP being considered identical if they are the same up to translation. The qualitative behaviour of the radius of convergence  $q_c(x)$  of  $G(x, q)$ , seen as a series in  $q$  for fixed values of  $x$ , was discussed in [7, 8]. This quantity is closely related to the asymptotic growth rate of the partition function  $Z_n(x) = \sum_{m=0}^{\infty} p_{m,n} x^m$ , and thus physically to the free energy per unit area in the thermodynamic limit of infinite area. The model exhibits a phase transition at a value  $x_c$  at which  $q_c(x)$  is not analytic. More precisely, it was shown that there exists a value  $x_c > 0$  such that for  $0 \leq x \leq x_c$ ,  $q_c(x) = 1$ ; for  $x > x_c$ ,  $q_c(x)$  is a continuous function of  $x$ ,  $q_c(x) < 1$  and  $\lim_{x \rightarrow \infty} q_c(x) = 0$ —see figure 2. For  $q < q_c(x)$ , polygons with relatively small area dominate the sum (1). This part of the  $(x, q)$ -plane is called the droplet phase. Analogously, the region  $q > 1$  is labelled the inflated phase. The region where  $q_c(x) < q < 1$  is described as the ‘seaweed’ phase, in which the typical conformation consists of a space-filling, convoluted polygon. Exact enumerations yield the estimate  $x_c \simeq 0.379$  [9]. The point  $(x, q) = (x_c, 1)$  is called a tricritical point [10].

In [11, 12], exact enumeration data was used to analyse the singular behaviour of the area–perimeter generating function of *rooted* SAP. In rooted SAP, there exists one distinguished point on the perimeter of the SAP, therefore the number of rooted SAP with perimeter  $m$  and area  $n$  is  $mp_{m,n}$ , and the area–perimeter generating function is  $R(x, q) = x \frac{d}{dx} P(x, q)$ . It



**Figure 1.** A self-avoiding polygon (SAP) on  $\mathbb{Z}^2$  of perimeter 52 and area 37.



**Figure 2.** Qualitative picture of the phase diagram of the SAP model of vesicles. The line  $q_c(x)$  is the boundary of the droplet phase. The exponents  $\gamma, \theta$  and  $\phi$  characterise the singular behaviour of the generating function  $P(x, q)$  around the tricritical point.

was conjectured that in the vicinity of the point  $(x, q) = (x_c, 1)$ , the singular part of this function satisfies the scaling relation

$$R^{\text{sing}}(x, e^{-\epsilon}) \sim \epsilon^\theta F((x_c - x)\epsilon^{-\phi}) \quad (\epsilon \rightarrow 0^+), \tag{2}$$

with the scaling function being, up to prefactors, given by the logarithmic derivative of the Airy function, which is defined for  $z \in \mathbb{C}$  as [13]

$$\text{Ai}(z) = \frac{1}{2\pi i} \int_{\infty e^{-i\pi/3}}^{\infty e^{i\pi/3}} \exp\left(\frac{u^3}{3} - zu\right) du. \tag{3}$$

Physically, the parameter  $\epsilon = -\ln(q)$  plays the role of an osmotic pressure acting onto the outside of the vesicle.

In [4] it was argued via field theoretic methods that, upon introducing further interactions into the SAP model, one should be able to observe multicritical points of higher order, described by scaling functions of more than one variable. More precisely, upon introducing  $\ell - 1$  further interactions  $(w_j)_{j=2}^\ell$ , there should exist multicritical points in the vicinity of which the singular part of the multivariate generating function of rooted SAP satisfies the scaling relation

$$R^{\text{sing}}(w_2, \dots, w_\ell, x, q) = \epsilon^\theta F(\alpha_1 \epsilon^{\phi_1}, \alpha_2 \epsilon^{\phi_2}, \dots, \alpha_\ell^{\phi_\ell}), \tag{4}$$

where the variables  $(\alpha_j)_{j=1}^\ell$  depend on the parameters of the generating function, the crossover exponents  $\phi_j$  are given by

$$\phi_j = \phi_j(\ell) = \frac{\ell + 2 - j}{\ell + 2} \quad (1 \leq j \leq \ell), \tag{5}$$

and  $\theta = \frac{1}{\ell+2}$  is a critical exponent. The scaling function  $F(s_1, s_2, \dots, s_\ell)$  is expressible in terms of generalised, higher-order Airy integrals, defined as

$$\Theta_\ell(s_1, s_2, \dots, s_\ell) = \frac{1}{2\pi i} \int_{\infty e^{-i\pi/(\ell+2)}}^{\infty e^{i\pi/(\ell+2)}} \exp\left(\frac{u^{\ell+2}}{\ell+2} - \sum_{j=1}^\ell s_j u^j\right) du, \tag{6}$$

However, no details of the interactions necessary to observe these multicritical points were given in that reference.

Progress in the study of SAPs can be made by considering directed subclasses such as staircase polygons. In [20] it was shown that staircase polygons show the same phase diagram as shown in figure 2, with a tricritical point around which the area–width generating function satisfies the same scaling behaviour as the one conjectured to hold for rooted SAPs. Dyck paths constitute an even more radical simplification of SAPs; their area–perimeter generating function has also been shown to satisfy the same scaling relation [17]. This motivates us to search for a model of one-dimensional lattice paths with a multicritical point as the one conjectured in [4] to hold for SAPs.


### 3. The model

One-dimensional lattice paths occur in many applications in probability theory, combinatorics and statistical physics. For  $m \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$ , a one-dimensional lattice path of length  $m$  is a sequence  $(\mathbf{r}_0, \mathbf{r}_1, \dots, \mathbf{r}_m)$  of points of  $\mathbb{Z}^2$ , where for  $0 < j \leq m$ ,  $\mathbf{r}_j - \mathbf{r}_{j-1} \in \{1\} \times \mathcal{S}$ , with  $\mathcal{S} \subseteq \mathbb{Z}$  [14]. One usually fixes  $\mathbf{r}_0 = (0, 0)$ . The path then stays in the right half-plane. Paths restricted further to stay in the upper right quarter plane  $\mathbb{N}_0 \times \mathbb{N}_0$  are called meanders, paths which end on the horizontal line  $\mathbb{N}_0 \times \{0\}$  are called bridges, and paths which are both meanders and bridges are called excursions. Łukasiewicz paths, which encode rooted ordered trees [15], are excursions with  $\mathcal{S} = \{k \in \mathbb{Z} \mid k \geq -1\}$ . Excursions with  $\mathcal{S} = \{-1\} \cup \{j \mid k \leq j \leq \ell\}$ , where  $k, \ell \in \mathbb{N}_0$ , are called  $(k, \ell)$ -Łukasiewicz paths [16]. Figure 3 shows a  $(1, 3)$ -Łukasiewicz path of length 13, with the horizontal and vertical axes of  $\mathbb{Z}^2$  labelled by  $x$  and  $y$ , respectively. A step in the direction  $(1, -1)$  is called a down-step and a step in direction  $(1, k)$ , where  $k \geq 1$  is called an up-step of length  $k$ . The *height* of a point is its distance from the  $x$ -axis, and the height of a step is the height of its starting point. The heights of the up-steps in figure 3 are marked by red dotted lines. Special subclasses of Łukasiewicz paths are Dyck and Motzkin paths, corresponding to  $(k, \ell) = (1, 1)$  and  $(k, \ell) = (0, 1)$ , respectively.

In this letter we consider the generating function

$$G_\ell(\mathbf{w}, x, q) = \sum_{m, n, j_1, \dots, j_\ell} c(j_1, \dots, j_\ell, m, n) w_1^{j_1} \dots w_\ell^{j_\ell} x^m q^n, \tag{7}$$

where we abbreviate  $w_1, \dots, w_\ell = \mathbf{w}$  and where  $c(j_1, \dots, j_\ell, m, n)$  is the number of  $(1, \ell)$ -Łukasiewicz paths with  $m$  up-steps, of which  $j_1$  have length one,  $j_2$  have length two etc, and the sum of the heights of all the up-steps, which is an area-like quantity, is  $n$ . For example, the path shown in figure 3 has the weight  $w_1^3 w_2 w_3 x^5 q^5$  in the generating function  $G_3(w_1, w_2, w_3, x, q)$ .



**Figure 3.** A (1, 3)-Łukasiewicz path of length 13.

One can set  $w_1 = 1$  without loss of generality. In the following we therefore write  $G_\ell(1, w_2, \dots, w_\ell, x, q) \equiv G_\ell(w_2, \dots, w_\ell, x, q)$ .

### 4. Previous results

In [17], the asymptotic behaviour of  $G_1(x, q)$  in the limit  $q \rightarrow 1^-$  was analysed, despite the parameters being interpreted slightly differently there. It was shown that, in the vicinity of the tricritical point  $(x, q) = (x_c, 1)$  with  $x_c = \frac{1}{4}$ , the singular part of the generating function satisfies the scaling relation

$$G_1^{\text{sing}}(x, e^{-\epsilon}) = \epsilon^\theta F((x_c - x)\epsilon^{-\phi}), \tag{8}$$

where  $\theta = \frac{1}{3}$ ,  $\phi = \frac{2}{3}$  and  $F(s) = b_0 \frac{d}{ds} \ln(\text{Ai}(b_1 s))$ , with positive constants  $b_0$  and  $b_1$ . Consistently with the solution  $G_1(x, 1) = \frac{1}{2x}(1 - \sqrt{1 - 4x})$ , equation (8) implies with the asymptotic expansions of the Airy function and its derivative [13] that  $G_1^{\text{sing}}(x, 1) \sim 2(x_c - x)^\gamma$ , where  $\gamma = \frac{\theta}{\phi} = \frac{1}{2}$ . Up to different constants  $b_0$  and  $b_1$ , the scaling relation (8) is identical to the one in equation (2) that was conjectured to hold for rooted self-avoiding polygons.

### 5. Result

In [18], it was shown rigorously that in the vicinity of the multicritical point  $(w_2, x, q) = (-\frac{1}{9}, \frac{1}{3}, 1)$ , the singular part of  $G_2(w_2, x, q)$  satisfies the scaling relation

$$G_2^{\text{sing}}(w_2, x, q) = \epsilon^\theta F(\alpha_1 \epsilon^{-\phi_1}, \alpha_2 \epsilon^{-\phi_2}), \tag{9}$$

where the scaling variables  $\alpha_1$  and  $\alpha_2$  are analytic functions of  $w_2$  and  $x$ ,  $\theta = \frac{1}{4}$ ,  $\phi_1 = \frac{3}{4}$ , and  $\phi_2 = \frac{1}{2}$ . The scaling function  $F$  is expressible via  $\Theta_2(s_1, s_2)$ , where the definition of  $\Theta_2$  is given by equation (6).

This result is generalised in this letter. More specifically, we show that in the model of  $(1, \ell)$ -Łukasiewicz paths for arbitrary  $\ell \geq 2$ , there exists a multicritical point  $(w_2, \dots, w_\ell, x, q)$ , with  $x = (\ell + 1)^{-1}$ ,  $q = 1$  and

$$w_k = \frac{2}{\ell(\ell + 1)} \left( \frac{-1}{\ell + 1} \right)^{k-1} \binom{\ell + 1}{k + 1} \quad (2 \leq k \leq \ell), \tag{10}$$

in the vicinity of which the generating function  $G_\ell(w_2, \dots, w_\ell, x, q)$  satisfies a scaling relation of the form of equation (4) with the same scaling function and the same critical exponents as

predicted in [4]. We thus present an exactly solvable model representing a concrete realisation of the multicritical scaling postulated in that reference.

**6. Method**

To obtain an asymptotic expression for  $G_\ell(w_2, \dots, w_\ell, x, q)$  in the limit  $q \rightarrow 1^-$  for arbitrary  $\ell \geq 2$ , one proceeds analogously to [17, 18]. From a simple factorisation argument [16] one obtains the functional equation

$$G_\ell(\mathbf{w}, x, q) = 1 + xG_\ell(\mathbf{w}, x, q) \left( \sum_{k=1}^{\ell} w_k \prod_{j=1}^k G_\ell(\mathbf{w}, q^j x, q) \right). \tag{11}$$

We linearise equation (11) by using the ansatz

$$G_\ell(w_2, \dots, w_\ell, x, q) = \frac{\Phi(qx)}{\Phi(x)}, \tag{12}$$

where  $\Phi(x) \equiv \Phi(w_2, \dots, w_\ell, x, q)$ . The solution of the linearised equation is then given by the  $q$ -hypergeometric series [19]

$$\Phi(x) = \sum_{n=0}^{\infty} \frac{\prod_{j=1}^{\ell-1} (\omega_j; q)_n}{(q; q)_n} (-x)^n q^{n^2-n}, \tag{13}$$

where  $(z; q)_n = \prod_{j=0}^{n-1} (1 - zq^j)$  is the  $q$ -Pochhammer symbol and the parameters  $(\omega_k)_{k=1}^{\ell-1}$  satisfy

$$w_k = (-1)^{k-1} \sum_{j_1=1}^{\ell-1} \sum_{j_2=j_1+1}^{\ell-1} \dots \sum_{j_{k-1}=j_{k-2}+1}^{\ell-1} \prod_{p=1}^{k-1} \omega_{j_p} \quad (2 \leq k \leq \ell). \tag{14}$$

Using the identity

$$\frac{(-1)^{n+1} q^{\binom{n}{2}}}{(q; q)_n (q; q)_\infty} = \text{Res} [(z; q)_\infty^{-1}; z = q^{-n}], \tag{15}$$

we obtain for  $k \in \mathbb{Z}$  the integral expression

$$\Phi(q^k x) = \frac{A}{2\pi i} \int_C \frac{z^{\frac{1}{2}(\log_q(z)+1) - \log_q(x)}}{z^k \left( \prod_{j=1}^{\ell-1} (\omega_j/z; q)_\infty \right) (z; q)_\infty} dz, \tag{16}$$

where the prefactor  $A$  is independent of  $k$ , and the contour  $C$  connects the points  $-i\infty$  and  $i\infty$  such that all zeros of  $(z; q)_\infty$  lie to the right of the contour and all zeros of  $z^k \left( \prod_{j=1}^{\ell-1} (\omega_j/z; q)_\infty \right)$  lie to the left of the contour. Substituting an asymptotic expression for the  $q$ -Pochhammer symbol [20], the above integral satisfies

$$\Phi(q^k x) \sim \frac{A}{2\pi i} \int_C \exp\left(\frac{1}{\epsilon} f(z)\right) \frac{g(z)}{z^k} dz \quad (q = e^{-\epsilon} \rightarrow 1^-), \tag{17}$$

where the functions are

$$f(z) = \log(x) \log(z) - \frac{\log(z)^2}{2} + \text{Li}_2(z) + \sum_{j=1}^{\ell-1} \text{Li}_2\left(\frac{\omega_j}{z}\right),$$

$$g(z) = \left( \frac{z^\ell}{(1-z) \prod_{j=1}^{\ell-1} (z - \omega_j)} \right)^{\frac{1}{2}},$$

and where  $\text{Li}_2(z)$  is the Euler dilogarithm [13]. The saddle points of  $f(z)$  are the zeros of the polynomial

$$\chi(z) = x \left( \prod_{j=1}^{\ell-1} z - \omega_j \right) - z^\ell (1 - z)$$

$$= z^{\ell+1} \left( 1 - \frac{1}{z} + \frac{x}{z^2} + \frac{x}{z} \sum_{k=2}^{\ell} \frac{w_k}{z^k} \right) \quad (z \neq 0).$$

Comparing the bracket in the last expression with equation (11), we see that the solution of equation (11) for  $q = 1$  is equal to the inverse of a saddle point of  $f(z)$ . If we set the weights for  $2 \leq k \leq \ell$  to the ones given in equation (10), then  $\ell + 1$  saddle points coalesce in the point  $z = (\ell + 1)^{-1}$  for  $x = (\ell + 1)^{-1}$ . To obtain an asymptotic expression for  $\Phi(q^k x)$ , we apply a method devised in [21], based on a theorem from [22], from which it follows that if the parameters of the function  $f(z)$  are close to the critical values given in equation (10), then there exists a mapping  $T : u \mapsto z(u)$ , which is analytic and bijective in the vicinity of the point  $z = (\ell + 1)^{-1}$ , such that

$$f(z(u)) = \frac{u^{\ell+2}}{\ell + 2} - \sum_{j=0}^{\ell} \alpha_j u^j = p(u). \tag{18}$$

Moreover, the coefficients  $(\alpha_j)_{j=0}^{\ell}$  are analytic functions of  $(w_j)_{j=2}^{\ell}$  and  $x$  in the region around the point of coalescence of the saddle points. Using the transformation  $T$ , the integral in equation (17) can be rewritten as

$$\Phi(q^k x) = \frac{A}{2\pi i} \int_{C'} \exp\left(\frac{1}{\epsilon} \left[ \frac{u^{\ell+2}}{\ell + 2} - \sum_{j=0}^{\ell} \alpha_j u^j \right]\right) S_k(u) du, \tag{19}$$

where  $q = e^{-\epsilon} \rightarrow 1^-$ ,  $C'$  is the image of the contour  $C$  under  $T^{-1}$  and  $S_k(u) = \frac{g(z(u))}{z(u)^k} \frac{dz}{du}$ . Now one writes

$$S_k(u) = \sum_{j=0}^{\ell} P_j^{(k)} u^k + p'(u) H(u), \tag{20}$$

where the  $(P_j^{(k)})_{j=0}^{\ell}$  are analytic functions of the  $(w_j)_{j=2}^{\ell}$  and  $x$ , and  $H(u)$  is some analytic function of  $u$ . Substituting equation (20) into equation (19), one arrives at the asymptotic expression

$$\Phi(q^k x) = A \sum_{j=0}^{\ell} P_j^{(k)} \epsilon^{\frac{j}{\ell+2}} \Theta_\ell^{(j)}(\alpha_1 \epsilon^{-\frac{\ell+1}{\ell+2}}, \alpha_2 \epsilon^{-\frac{\ell}{\ell+2}}, \dots, \alpha_\ell \epsilon^{-\frac{2}{\ell+2}}), \tag{21}$$

where  $\Theta_\ell^{(0)} = \Theta_\ell$  and  $\Theta_\ell^{(j)} = -\frac{\partial}{\partial s_j} \Theta_\ell$  for  $1 \leq j \leq \ell$ ; see equation (6) for the definition of  $\Theta_\ell$ . By substituting equation (21) into equation (12) for  $k = 0$  and  $k = 1$ , we obtain an asymptotic expression for  $G_\ell(w_2, \dots, w_\ell, x, q)$  which is valid uniformly with respect to the parameters  $(w_j)_{j=2}^\ell$  and  $x$ , in particular in the vicinity of the multicritical point at which  $\ell + 1$  saddle points of the function  $f(z)$  coalesce. Close to the multicritical point, the singular part of  $G_\ell(w_2, \dots, w_\ell, x, q)$  satisfies the scaling relation

$$G_\ell^{\text{sing}}(w_2, \dots, w_\ell, x, e^{-\epsilon}) = \epsilon^\theta F(\alpha_1 \epsilon^{-\phi_1}, \dots, \alpha_\ell \epsilon^{-\phi_\ell}), \quad (22)$$

where  $F$  is expressible via  $\Theta_\ell(s_1, \dots, s_\ell)$ ,  $\theta = \frac{1}{\ell+2}$ , and the  $\phi_j \equiv \phi_j(\ell)$  are given by equation (5).

## ORCID iDs

N Haug  <https://orcid.org/0000-0002-5130-9193>

T Prellberg  <https://orcid.org/0000-0003-0519-6762>

## References

- [1] Cardy J 1996 *Scaling and Renormalization in Statistical Physics (Cambridge Lecture Notes in Physics)* (Cambridge: Cambridge University Press)
- [2] Henkel M 2013 *Conformal Invariance and Critical Phenomena* (Berlin: Springer)
- [3] Wu T T, McCoy B M, Tracy C A and Barouch E 1976 *Phys. Rev. B* **13** 316
- [4] Cardy J 2001 *J. Phys. A: Math. Gen.* **34** L665
- [5] Seifert U 1997 *Adv. Phys.* **46** 13
- [6] Leibler S, Singh R R P and Fisher M E 1987 *Phys. Rev. Lett.* **59** 1989
- [7] Banavar J R, Maritan A and Stella A 1991 *Phys. Rev. A* **43** 5752
- [8] Fisher M E, Guttman A J and Whittington S G 1991 *J. Phys. A: Math. Gen.* **24** 3095
- [9] Jensen I and Guttman A J 1999 *J. Phys. A: Math. Gen.* **32** 4867
- [10] Lawrie I D and Sarbach S 1984 *Phase Transitions and Critical Phenomena* vol 9 ed C Domb and J L Lebowitz (New York: Academic) ch1 pp 2–161
- [11] Richard C, Guttman A J and Jensen I 2001 *J. Phys. A: Math. Gen.* **34** L495
- [12] Richard C, Jensen I and Guttman A J 2004 *J. Stat. Mech.* P08007
- [13] Olver F W J et al ed 2020 *DLMF, Digital Library of Mathematical Functions* <http://dlmf.nist.gov/> Release 1.0.26 of 2020-03-15
- [14] Banderier C and Flajolet P 2002 *Theor. Comput. Sci.* **281** 37
- [15] Le Gall J-F 2005 *Probab. Surv.* **2** 245
- [16] Brak R, Iliev G K and Prellberg T 2011 *J. Stat. Phys.* **145** 669
- [17] Haug N and Prellberg T 2015 *J. Math. Phys.* **56** 043301
- [18] Haug N, Olde Daalhuis A and Prellberg T 2017 *J. Stat. Phys.* **166** 1193
- [19] Gasper G and Rahman M 1990 *Basic Hypergeometric Series (Encyclopedia of Mathematics and its Applications)* vol 96 (Cambridge: Cambridge University Press) p 428
- [20] Prellberg T 1995 *J. Phys. A: Math. Gen.* **28** 1289
- [21] Ursell F 1972 *Math. Proc. Camb. Phil. Soc.* **72** 49
- [22] Levinson N 1961 *Duke Math. J.* **28** 345